



Matrix Product States for Lattice Gauge Theories

Krzysztof Cichy

NIC, DESY Zeuthen, Germany

Adam Mickiewicz University, Poznań, Poland


in collaboration with:

Mari Carmen Bañuls (MPQ Garching)

J. Ignacio Cirac (MPQ Garching)

Karl Jansen (DESY Zeuthen)

Hana Saito (Humboldt Universität, DESY Zeuthen)



Overview



Overview

Seminar outline


Introduction

Results

Summary

The talk I will present differs a bit from typical ETMC talks...

| | typical ETMC | this talk |
|-------------|--------------|-----------------|
| theory | QCD | QED |
| dimension | 3+1 | 1+1 |
| fermions | twisted mass | staggered |
| formulation | Lagrangian | Hamiltonian |
| method | Monte Carlo | tensor networks |



Seminar outline



1. Introduction

- Motivation – Lattice QCD
- Schwinger model
- Hamiltonian approach
 - ★ strong coupling expansion
 - ★ Matrix Product States


2. Results

- Ground state energy
- Vector and scalar mass gap
- Chiral condensate $T = 0$
- Chiral condensate $T > 0$

3. Prospects

Based on:

- K. Cichy, A. Kujawa-Cichy and M. Szyniszewski, “Lattice Hamiltonian approach to the massless Schwinger model: Precise extraction of the mass gap,” *Comput. Phys. Commun.* **184** (2013) 1666, [arXiv:1211.6393 [hep-lat]]
- M. C. Bañuls, K. Cichy, K. Jansen and J. I. Cirac, “The mass spectrum of the Schwinger model with Matrix Product States,” *JHEP* **1311** (2013) 158, [arXiv:1305.3765 [hep-lat]]
- M. C. Bañuls, K. Cichy, J. I. Cirac, K. Jansen and H. Saito, “Matrix Product States for Lattice Field Theories,” *PoS(LATTICE 2013)*332, [arXiv:1310.4118 [hep-lat]]



Lattice QCD



- The most common approach to Lattice QCD simulations consists in sampling the QCD path integral numerically via the Monte Carlo method.

- The QCD path integral: $Z = \int D\bar{\psi} D\psi DU e^{-S_{gauge}[U] - S_{ferm}[\psi, \bar{\psi}, U]}$.

- The fermionic degrees of freedom can be integrated out:


$$Z = \int DU e^{-S_{gauge}[U]} \prod_{f=1}^{N_f} \det(\hat{D}_f[U]),$$

where $\det(\hat{D}_f[U])$ is the determinant of the Dirac operator matrix for fermion flavour f .

- The fermionic determinant $\det(\hat{D}_f[U])$ is by far the highest cost in a MC simulation. But, due to γ_5 -Hermiticity ($\gamma_5 \hat{D}_f \gamma_5 = \hat{D}_f^\dagger$) it is **real**, so MC simulations are **possible**:

$$\det(\gamma_5(\hat{D}_f + m)\gamma_5) = \det(\hat{D}_f^\dagger + m) = \det(\hat{D}_f + m)^\dagger.$$

- First approximation \Rightarrow neglect the determinant (“**quenched approximation**”) – commonly used until early 2000s.
- **Dynamical simulations** \Rightarrow take the determinant into account.



Problems of Lattice QCD



LQCD simulations led to spectacular successes. However, there are some areas where progress is hard to achieve:


- non-vanishing chemical potential μ – if $\mu \neq 0$, the determinant becomes complex:

$$\det \left(\gamma_5 (\hat{D}_f + m + \mu \gamma_0) \gamma_5 \right) = \det \left(\hat{D}_f^\dagger + m - \mu \gamma_0 \right) = \det \left(\hat{D}_f + m - \mu^* \gamma_0 \right)^\dagger,$$

determinant real only if μ taken to be purely imaginary.

Ways to tackle the problem: reweighting, Taylor expansion, analytic continuation from imaginary μ .


- LQCD works in Euclidean space, related to Minkowski space by analytic continuation – hence time is imaginary. Hence, it is **not possible** to simulate real-time phenomena, i.e. non-equilibrium dynamics.



[K. Fukushima, T. Hatsuda, Rep. Prog. Phys. 74 (2011) 14001]

Alternative approaches wanted for these classes of problems!

Tensor Networks?




The way to apply TNS to QCD is a long one.

- **START**: Schwinger model, i.e. an Abelian gauge theory with U(1) gauge group, 1+1 dimensions

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}(i \not{\partial} - g \not{A} - m)\psi$$

- **NATURAL NEXT STEP**: non-Abelian gauge theories (SU(2), SU(3)) in 1+1 dimensions
- **AND ALSO**: go to 2+1 dimensions
- **FINALLY**: go to 3+1 dimensions, non-Abelian gauge group SU(3) for QCD

All these next steps non-trivial and challenging.



The Schwinger model



The Schwinger model is QED in 1+1 dimensions:

[J. S. Schwinger, Phys. Rev. **128** (1962) 2425]

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}(i \not{\partial} - g \not{A} - m)\psi$$

where ψ is a 2-component spinor field. The field strength term is:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu.$$

The coupling g has dimensions of mass (theory super-renormalizable). Using g as the scale of energy, the physical properties of the model are then functions of the dimensionless ratio m/g .

- simplest gauge theory
- but physics still surprisingly rich
- in several aspects resembles much more complex theories (QCD)
- standard toy model for testing lattice techniques

Overview
Seminar outline

Introduction

Lattice QCD

Problems

Road map to QCD
with TNS


Schwinger model

Hamiltonian
approach

Tensor Network
States

Results

Summary



The Schwinger model



[J. S. Schwinger, “Gauge Invariance and Mass. 2.,” *Phys. Rev.* **128** (1962) 2425.]

Abstract: *The possibility that a vector gauge field can imply a nonzero mass particle is illustrated by the exact solution of a one-dimensional model.*

Most prominent feature of the Schwinger model: **non-perturbative generation of mass gap!**

The mass gap can be calculated analytically: $\frac{M_V}{g} = \frac{1}{\sqrt{\pi}} \approx 0.564189584$.

How well can lattice techniques reproduce this number?

- 0.555(25) – MC [O. Martin and S. Otto, *Nucl. Phys. B* **203** (1982) 297]
- 0.560(10) – Hamiltonian approach [D. P. Crewther and C. J. Hamer, *Nucl. Phys. B* **170** (1980) 353]
- 0.565(2) – Hamiltonian approach + renormalization of coupling [A. C. Irving and A. Thomas, *Nucl. Phys. B* **215** (1983) 23]
- 0.56417(2) – Hamiltonian approach + renormalization of coupling [P. Sriganesh, R. Bursill and C. J. Hamer, *Phys. Rev. D* **62** (2000) 034508]
- 0.56419(4) – Hamiltonian approach + DMRG [T. Byrnes, P. Sriganesh, R. J. Bursill and C. J. Hamer, *Phys. Rev. D* **66** (2002) 013002]

Overview
Seminar outline

Introduction

Lattice QCD


Problems
Road map to QCD
with TNS

Schwinger model

Hamiltonian
approach
Tensor Network
States

Results

Summary



$$\mathcal{L} \rightarrow \mathcal{H}$$



The Hamiltonian \mathcal{H} is the Legendre transform of the Lagrangian \mathcal{L} :

$$\mathcal{H} = \pi^\mu \dot{A}_\mu - \mathcal{L},$$

where:

$$\pi^\mu = \frac{\partial \mathcal{L}}{\partial \dot{A}_\mu} = -F^{0\mu}.$$

We choose the time like axial gauge $A_0 = 0$:

$$H = \int dx \left(-i\bar{\psi}\gamma^1(\partial_1 + igA_1)\psi + m\bar{\psi}\psi + \frac{1}{2}E^2 \right).$$

The γ matrices:


$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Going to the lattice:

$$U(n, n+1) = e^{i\theta(n)} = e^{-iagA^1(n)}$$

fermionic fields are associated with lattice sites and gauge fields with lattice links

- Overview
- Seminar outline
- Introduction
- Lattice QCD
- Problems
- Road map to QCD with TNS
- Schwinger model
- Hamiltonian approach**
- Tensor Network States
- Results
- Summary



Staggered discretization



- Overview
- Seminar outline
- Introduction
- Lattice QCD
- Problems
- Road map to QCD with TNS
- Schwinger model
- Hamiltonian approach**
- Tensor Network States
- Results
- Summary

The Hamiltonian becomes:

$$H = -\frac{i}{2a} \sum_{n=0}^{M-1} \left(\phi^\dagger(n) e^{i\theta(n)} \phi(n+1) - \phi^\dagger(n+1) e^{-i\theta(n)} \phi(n) \right) + \\ + m \sum_{n=0}^{M-1} (-1)^n \phi^\dagger(n) \phi(n) + \frac{ag^2}{2} \sum_{n=0}^{M-1} L^2(n),$$

in the Kogut-Susskind discretization:


[T. Banks, L. Susskind and J. B. Kogut, Phys. Rev. D **13** (1976) 1043]

[J. B. Kogut and L. Susskind, Phys. Rev. D **11** (1975) 395.]

$$\phi(n)/\sqrt{a} \rightarrow \begin{cases} \psi_{\text{upper}}(x) & n \text{ even} \\ \psi_{\text{lower}}(x) & n \text{ odd} \end{cases}$$

The correspondence between lattice and continuum fields is:

$$\frac{1}{ag} \theta(n) \rightarrow -A^1(x) \\ gL(n) \rightarrow E(x).$$



Basic ingredients



Overview
Seminar outline

Introduction

Lattice QCD

Problems

Road map to QCD
with TNS

Schwinger model


**Hamiltonian
approach**

Tensor Network
States

Results

Summary

- $\phi(n)$ is a single-component fermion field, defined on each site of a M -site lattice with periodic b.c. and obeying the anticommutation relations: $\{\phi^\dagger(n), \phi(m)\} = \delta_{nm}$, $\{\phi(n), \phi(m)\} = 0$, $\{\phi^\dagger(n), \phi^\dagger(m)\} = 0$
- The gauge field variable $\theta(n)$ is defined on the link between sites n and $n + 1$ and is related to the spatial component of the Abelian vector potential by $\theta(n) = agA(n)$
- The angular momentum variable $L(n)$ is related to the electric field $E(n)$ by the relation $L(n) = E(n)/g$ and to the gauge field by the commutation relations: $[\theta(n), L(m)] = i\delta_{nm}$. The possible values of $L(n)$ are quantized: $L(n)|l\rangle = l|l\rangle$, $l = 0, \pm 1, \pm 2, \dots$. This implies: $e^{\pm i\theta(n)}|l\rangle = |l \pm 1\rangle$
- m – fermion mass
- g – gauge coupling
- a – lattice spacing
- M – lattice size



Jordan-Wigner transformation



Overview
Seminar outline

Introduction

Lattice QCD

Problems

Road map to QCD
with TNS

Schwinger model

**Hamiltonian
approach**

Tensor Network
States

Results

Summary


$$H = -\frac{i}{2a} \sum_{n=0}^{M-1} \left(\phi^\dagger(n) e^{i\theta(n)} \phi(n+1) - \phi^\dagger(n+1) e^{-i\theta(n)} \phi(n) \right) + \\ + m \sum_{n=0}^{M-1} (-1)^n \phi^\dagger(n) \phi(n) + \frac{ag^2}{2} \sum_{n=0}^{M-1} L^2(n),$$

For numerics, it is convenient to perform the Jordan-Wigner transformation: [P. Jordan, E. Wigner, Z. Phys. 47 (1928) 631.]

$$\phi(n) = \prod_{p < n} (i\sigma^3(p)) \sigma^-(n),$$

where $\sigma^i(n)$ are Pauli matrices ($\sigma^\pm = \sigma^1 \pm i\sigma^2$). This gives:

$$H = -\frac{1}{2a} \sum_{n=0}^{M-1} \left(\sigma^+(n) e^{i\theta(n)} \sigma^-(n+1) + \sigma^+(n+1) e^{-i\theta(n)} \sigma^-(n) \right) + \\ + \frac{m}{2} \sum_{n=0}^{M-1} (1 + (-1)^n \sigma^3(n)) + \frac{ag^2}{2} \sum_{n=0}^{M-1} L^2(n).$$



Choice of basis



Rewrite Hamiltonian in a dimensionless form: $W = \frac{2}{ag^2} H_{JW} = W_0 - xV$, with:

$$W_0 = \frac{m}{ag^2} \sum_{n=0}^{M-1} (1 + (-1)^n \sigma^3(n)) + \sum_{n=0}^{M-1} L^2(n),$$

$$V = \sum_{n=0}^{M-1} \left(\sigma^+(n) e^{i\theta(n)} \sigma^-(n+1) + \sigma^+(n+1) e^{-i\theta(n)} \sigma^-(n) \right)$$


$$x \equiv \beta = 1/a^2 g^2.$$

- Natural choice of basis: direct product of Ising basis $\{|i\rangle\}$, acted upon by Pauli spin operators, and the ladder space of states $\{|l\rangle\}$:

$$|i_0 i_1 \dots i_{M-2} i_{M-1}\rangle \otimes |l_{0,1} l_{1,2} \dots l_{M-2,M-1} (l_{M-1,0})\rangle,$$

where $(l_{M-1,0})$ is present if PBC are considered and absent for OBC.

- Formally, the operator W_0 can be treated as an unperturbed part and V as a perturbation. Ground state of W_0 : $|0\rangle = |\downarrow\uparrow\downarrow\uparrow \dots \downarrow\uparrow\rangle \otimes |0000 \dots 00\rangle$,
- The perturbation operator V flips two neighbouring spins and couples them via a gauge field excitation (flux line): $V | \bullet \quad \bullet \rangle = | \uparrow \rightsquigarrow \downarrow \rangle$



Choice of basis



- The gauge degrees of freedom $l_{i,i+1}$ can be eliminated using the Gauss law:


$$L_n - L_{n-1} = \frac{1}{2} (\sigma_n^z + (-1)^n),$$

leaving the basis states as:

$$|i_0 i_1 \dots i_{M-2} i_{M-1}\rangle \otimes |l\rangle,$$

with:

- ★ $l = 0, \pm 1, \pm 2, \dots$ for PBC,
- ★ $l = 0$ (or other constant) for OBC.
- With M -site lattice, $\dim(\text{spin part}) = 2^M$, while for the gauge part the basis is
 - ★ infinite-dimensional for PBC \Rightarrow truncation needed,
 - ★ one-dimensional for OBC.
- Truncation for PBC:
 - ★ at some finite $\pm l_{\max}$, thus reducing the basis to dimension $(2l_{\max} + 1)2^M$,
 - ★ or use strong coupling expansion (SCE):
 - [T. Banks, L. Susskind and J. B. Kogut, Phys. Rev. D **13** (1976) 1043]
 - [J. B. Kogut and L. Susskind, Phys. Rev. D **11** (1975) 395.]



Tensor Network States



Overview
Seminar outline

Introduction

Lattice QCD

Problems

Road map to QCD
with TNS


Schwinger model
Hamiltonian
approach

Tensor Network
States

Results

Summary

- An arbitrary state from a Hilbert space of an N -body interacting system needs in general an **exponential** number of coefficients – thus computational complexity increases very fast and prohibits exact diagonalization of systems larger than e.g.:
 - ★ $\mathcal{O}(20)$ Heisenberg spins (with a naive approach) or
 - ★ $\mathcal{O}(40)$ Heisenberg spins (using symmetries etc.).
- However, physical states (ground states, thermal states) of most systems are far from arbitrary.
- In many cases, they can be described by Tensor Network states that have only a **polynomial** number of parameters.
- In other words, only a small “corner” of the Hilbert space is physically relevant.



Matrix Product States





- A particularly successful and efficient family of Tensor Network states is called Matrix Product States (MPS).
- The MPS ansatz for some state $|\Psi\rangle$ has the following form:


$$|\Psi\rangle = \sum_{i_0 \dots i_{N-1}=1}^d \text{tr} \left(A_0^{i_0} \dots A_{N-1}^{i_{N-1}} \right) |i_0 \dots i_{N-1}\rangle,$$

where:

$|i_k\rangle$ are individual basis states for each site ($k = 0, \dots, d - 1$),
 d – dimension of one-site Hilbert space,
 each A_j^i is a D -dimensional matrix
 and D is called the bond dimension.



- The ground state can be found variationally by successively minimizing the energy $\frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle}$ with respect to each tensor A_j until convergence is achieved.
- Having the ground state, one can find ground state expectation values of any operator of interest.



Excited states




- After having found the ground state of the system, $|\Psi_0\rangle$, we can construct the projector onto the orthogonal subspace, $\Pi_0 = 1 - |\Psi_0\rangle\langle\Psi_0|$.
- The projected Hamiltonian, $\Pi_0 H \Pi_0$, has $|\Psi_0\rangle$ as eigenstate with zero eigenvalue, and the first excited state as eigenstate with energy E_1 .
- Given that $E_1 < 0$, what we can always ensure by adding an appropriate constant to H , the first excitation corresponds then to the state that minimizes the energy of the projected Hamiltonian:

$$E_1 = \min_{|\Psi\rangle} \frac{\langle\Psi|\Pi_0 H \Pi_0|\Psi\rangle}{\langle\Psi|\Psi\rangle} = \frac{\langle\Psi|(H - E_0|\Psi_0\rangle\langle\Psi_0|)|\Psi\rangle}{\langle\Psi|\Psi\rangle}.$$

- This minimization corresponds to finding the ground state of the effective Hamiltonian $H_{\text{eff}}[1] = \Pi_0 H \Pi_0$.
- The procedure can be concatenated to find subsequent energy levels, so that, to find the M -th excited state, we will search for the ground state of the Hamiltonian:

$$H_{\text{eff}}[M] = \Pi_{M-1} \dots \Pi_0 H \Pi_0 \dots \Pi_{M-1} = H - \sum_{k=0}^{M-1} E_k |\Psi_k\rangle\langle\Psi_k|.$$



Earlier works



Overview
Seminar outline

Introduction

Lattice QCD

Problems
Road map to QCD
with TNS

Schwinger model
Hamiltonian
approach

Tensor Network
States

Results


Summary

Density Matrix Renormalization Group approach:

- T. Byrnes, P. Sriganesh, R. J. Bursill and C. J. Hamer, “Density matrix renormalization group approach to the [massive Schwinger model](#),” PRD **66** (2002) 013002, [hep-lat/0202014].
- T. Sugihara, “Density matrix renormalization group in a two-dimensional $\lambda\phi^4$ [Hamiltonian lattice model](#),” JHEP **0405** (2004) 007, [hep-lat/0403008]

Matrix Product States approach:

- T. Sugihara, “Matrix product representation of gauge invariant states in a [Z\(2\) lattice gauge theory](#),” JHEP **0507** (2005) 022, [hep-lat/0506009]
- A. Milsted, J. Haegeman and T. J. Osborne, “Matrix product states and variational methods applied to [critical quantum field theory](#),” ($\lambda\phi^4$ theory) arXiv:1302.5582 [hep-lat]



Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation

Mass gaps


Chiral condensate

Some result

Continuum limit

Summary

Results



SCE+ED, infinite volume extrapolation



Overview
Seminar outline
Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation


Mass gaps

Chiral condensate


Some result

Continuum limit

Summary



$$F(1/M^2) = F_0 + F_2 \frac{1}{M^2} + F_4 \frac{1}{M^4} + F_6 \frac{1}{M^6}$$



SCE+ED, infinite volume extrapolation




- Overview
- Seminar outline
- Introduction

Results


SCE+ED

- MPS
- GS energy
- Excited states
- Dispersion relation
- Mass gaps
- Chiral condensate
- Some result
- Continuum limit

Summary



$$F(1/M^2) = F_0 + F_2 \frac{1}{M^2} + F_4 \frac{1}{M^4} + F_6 \frac{1}{M^6}$$



SCE+ED, continuum extrapolation




- Overview
- Seminar outline
- Introduction

Results


SCE+ED

- MPS
- GS energy
- Excited states
- Dispersion relation
- Mass gaps
- Chiral condensate
- Some result
- Continuum limit

Summary




$$F_0(ag) = F_{00} + F_{01} \cdot ag + F_{02} \cdot (ag)^2$$



SCE+ED, comparison with literature



| | M_S/g | | M_V/g | |
|------------------------------|----------------------|---------------------|---------------------|-----------------------|
| | result | error | result | error |
| exact | 1.12837916710 | – | 0.5641895836 | – |
| this work | 1.12837916719 | $8 \cdot 10^{-9}\%$ | 0.5641895845 | $1.8 \cdot 10^{-7}\%$ |
| [Crewther, Hamer 1980] | 1.120 | 0.7% | 0.560 | 0.7% |
| [Irving, Thomas 1982] | 1.128 | 0.03% | 0.565 | 0.1% |
| [Hamer et al. 1997] (I) | 1.25 | 11% | 0.56 | 0.7% |
| [Hamer et al. 1997] (II) | 1.14 | 1% | 0.57 | 1% |
| [Sriganesh et al. 1999] (I) | 1.11 | 1.6% | 0.563 | 0.2% |
| [Sriganesh et al. 1999] (II) | 1.1284 | 0.002% | 0.56417 | 0.003% |
| [Byrnes et al. 2002] | – | – | 0.56419 | $7 \cdot 10^{-5}\%$ |



Matrix Product States



Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation

Mass gaps

Chiral condensate

Some result

Continuum limit

Summary


We want to find:

- ground state energy
- vector mass gap
- scalar mass gap


for selected values of the fermion mass $m/g = 0, 0.125, 0.25, 0.5$.

Simulate with finite D (bond dimension), N (system size), x (inverse lattice spacing). We want:


- large enough D – check $D \in [20, 140]$,
- $N \rightarrow \infty$ – choose $N \in [100, 850]$ (note that $N \propto x$),
- $x \rightarrow \infty$ – choose $x \in [5, 600]$.




GS energy. Bond dimension




$m/g = 0, x = 100, N = 300$




$m/g = 0.125, x = 100, N = 300$




GS energy. Finite size scaling




$m/g = 0, x = 100$




$m/g = 0.125, x = 100$



GS energy. Continuum extrapolation



$m/g = 0$




$m/g = 0.125$

continuum result:

$-0.318338(22)_{D,N,x \text{ extrapol.}} (24)_{\text{fit ansatz}}$

$-0.318343(96)_{D,N,x \text{ extrapol.}} (25)_{\text{fit ansatz}}$

exact result: $1/\pi \approx -0.318310$



Computing the mass gap




Overview
Seminar outline
Introduction

Results
SCE+ED
MPS
GS energy

Excited states
Dispersion relation
Mass gaps
Chiral condensate
Some result
Continuum limit

Summary

- After having computed the GS energy, we want to compute the masses of the two lightest bound states (“mesons”) of the theory:
 - ★ vector meson,
 - ★ scalar meson.
- Important: we have to recognize the vector and scalar states – use the charge conjugation transformation:
 - ★ PBC – $C = -1 \Rightarrow$ vector state, $C = +1 \Rightarrow$ scalar state,
 - ★ OBC – C no longer an exact symmetry, but “enough” to differentiate vector vs. scalar.
- Note: with OBC translational symmetry is lost – hence we also have momentum excitations of the vector meson *before* we reach the scalar.



Dispersion relation




- Overview
- Seminar outline
- Introduction

- Results

- SCE+ED
- MPS
- GS energy
- Excited states
- Dispersion relation**
- Mass gaps
- Chiral condensate
- Some result
- Continuum limit

- Summary




Continuum momentum operator

$$\hat{P} = \int dx \Psi^\dagger(x) i \partial_x \Psi(x)$$

Lattice spin representation




$$\hat{O}_P = -ix \sum_n (\sigma_n^- \sigma_{n+1}^z \sigma_{n+2}^+ - H.c.)$$






Results for the mass gaps, $m/g = 0$



VECTOR



SCALAR




Truncation

$x = 100, N = 300$

Finite size scaling

$x = 100$




Continuum extrapolation






Results for the mass gaps, $m/g = 0.125$



VECTOR



SCALAR




Truncation

$x = 100, N = 300$

Finite size scaling

$x = 100$

Continuum extrapolation



Results for the mass gaps



| | Vector binding energy exact 0.5641895 | |
|-------|--|-------------|
| m/g | MPS with OBC | DMRG result |
| 0 | 0.56421(9) | 0.56419(4) |
| 0.125 | 0.53953(5) | 0.53950(7) |
| 0.25 | 0.51922(5) | 0.51918(5) |
| 0.5 | 0.48749(3) | 0.48747(2) |


| | Scalar binding energy exact 1.12838 | |
|-------|--|------------|
| m/g | MPS with OBC | SCE result |
| 0 | 1.1279(12) | 1.11(3) |
| 0.125 | 1.2155(28) | 1.22(2) |
| 0.25 | 1.2239(22) | 1.24(3) |
| 0.5 | 1.1998(17) | 1.20(3) |

DMRG result:

[T. Byrnes, P. Sriganesh, R. J. Bursill and C. J. Hamer, Phys. Rev. D **66** (2002) 013002]

SCE result:

[P. Sriganesh, R. Bursill and C. J. Hamer, Phys. Rev. D **62** (2000) 034508]



Chiral condensate



Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation

Mass gaps

Chiral condensate

Some result

Continuum limit

Summary

- The Schwinger model possesses a $U(1)_A$ chiral symmetry, which is broken by the chiral anomaly.
- This symmetry breaking is signaled by a non-zero value of the chiral condensate:


$$\Sigma = \frac{\sqrt{x}}{N} \sum_n (-1)^n \frac{1 + \sigma_n^z}{2}$$

→ compute GS expectation value of Σ .

- The naively computed condensate has a logarithmic divergence $\propto \frac{m}{g} \log ag$. This divergence can be subtracted off by subtracting the free theory contribution (in the infinite volume limit):

$$\Sigma_{\text{free}}^{(\text{bulk})}(m/g, x) = \frac{m}{\pi g} \frac{1}{\sqrt{1 + \frac{m^2}{g^2 x}}} K \left(\frac{1}{1 + \frac{m^2}{g^2 x}} \right),$$




where $K(u)$ is the complete elliptic integral of the first kind.






Results for the chiral condensate



$m/g = 0$



$m/g = 0.25$




Truncation

$x = 100, N = 300$

Finite size scaling

$x = 100$

Continuum extrapolation



Results for the chiral condensate



Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation

Mass gaps

Chiral condensate

Some result

Continuum limit


Summary

| | Subtracted condensate | | |
|-------|-----------------------|----------|----------|
| m/g | MPS with OBC | exact | Hosotani |
| 0 | 0.159930(8) | 0.159929 | - |
| 0.125 | 0.092023(4) | - | 0.0918 |
| 0.25 | 0.066660(11) | - | - |
| 0.5 | 0.042383(22) | - | - |

Exact result: $\frac{\Sigma}{g} = \frac{1}{2\pi^{3/2}} e^{\gamma_E} \approx 0.1599288$.

Hosotani (reduction to a quantum mechanics problem and numerical solution of the resulting Schrödinger equation):

[Y. Hosotani, "Chiral dynamics in weak, intermediate, and strong coupling QED in two-dimensions," In: Nagoya 1996, Perspectives of strong coupling gauge theories, 390-397 [hep-th/9703153].]



Chiral condensate at finite temperature



Analytic prediction for the behaviour of the condensate at finite T :

[I. Sachs, A. Wipf, "Finite Temperature Schwinger Model," *Helv. Phys. Acta* 65, 652 (1992), arXiv:1005.1822 [hep-th]]

$$\frac{\Sigma}{g} = \frac{1}{2\pi^{3/2}} e^{\gamma_E} e^{2I\left(\frac{g}{\sqrt{\pi T}}\right)},$$


where: $I(x) = \int_0^\infty dt (1 - e^{x \cosh(t)})^{-1}$.

- Low-temperature limit:

$$\frac{\Sigma}{g} = \frac{1}{2\pi^{3/2}} e^{\gamma_E} \approx 0.1599288$$

- High-temperature limit:

$$\frac{\Sigma}{g} \approx \frac{2T}{g} e^{-\pi^{3/2} T/g}$$



Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation


Mass gaps

Chiral condensate

Some result

Continuum limit

Summary



Idea of the computation



Given some operator \mathcal{O} , we want to calculate its thermal expectation value:

$$\langle \mathcal{O} \rangle_\beta = \frac{\text{Tr}(\mathcal{O}\rho(\beta))}{\text{Tr}(\rho(\beta))},$$

where $\beta = 1/T$, $\rho(\beta)$ is the thermal density operator.

- $\rho(\beta) = \rho(\beta/2)^\dagger \rho(\beta/2)$ to ensure positivity
- divide the interval $\beta/2$ into $N = \beta/\delta$ steps of length $\delta/2$:

$$\rho(\beta/2) = \underbrace{e^{-\frac{\delta}{2}H} \dots e^{-\frac{\delta}{2}H}}_{N=\beta/\delta \text{ times}}$$

- 2nd order Trotter expansion:

$$e^{-\frac{\delta}{2}H} \approx e^{-\frac{\delta}{4}H_g} \underbrace{e^{-\frac{\delta}{2}(H_{hop} + H_{mass})}}_{\approx e^{-\frac{\delta}{4}H_e} e^{-\frac{\delta}{2}H_o} e^{-\frac{\delta}{4}H_e}} e^{-\frac{\delta}{4}H_g},$$

H_e/H_o – on even/odd sites

where:

$$H = \underbrace{x \sum_{n=0}^{N-2} (\sigma_n^+ e^{i\theta_n} \sigma_{n+1}^- + H.c.)}_{H_{hop}} + \underbrace{\mu \sum_{n=0}^{N-1} (1 + (-1)^n \sigma_n^3)}_{H_{mass}} + \underbrace{\sum_{n=0}^{N-2} \left(l + \frac{1}{2} \sum_{k=0}^n ((-1)^k + \sigma_k^3) \right)^2}_{H_g}$$

Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation


Mass gaps

Chiral condensate

Some result

Continuum limit

Summary



Influence of D and δ




- Overview
- Seminar outline
- Introduction


- Results

- SCE+ED
- MPS
- GS energy
- Excited states
- Dispersion relation
- Mass gaps
- Chiral condensate
- Some result**
- Continuum limit




- Summary




$N = 20$, $x = 16$, $D = 20—160$, $\delta = 0.0001—0.01$



Zoom into high and low T





Towards the continuum limit



Overview
Seminar outline

Introduction

Results

SCE+ED

MPS

GS energy

Excited states

Dispersion relation


Mass gaps

Chiral condensate


Some result

Continuum limit



Summary




fixed $N/\sqrt{x} = 20$, i.e. pretty large volume



Infinite volume and continuum limits at $g\beta = 0.2$



analytic result at $g\beta = 0.2$ [I. Sachs, A. Wipf, 1992]: $\frac{\Sigma}{g} \approx 8.1 \cdot 10^{-12}$



Conclusions



Overview
Seminar outline

Introduction


Results

Summary

Conclusions

Prospects


- Proof of concept – the MPS approach can be used to extract:
 - ★ mass spectrum (GS energy, masses of lightest particles of a theory),
 - ★ ground state expectation values (chiral condensate).
- Precision better or comparable to best results in the literature, in some cases better than **0.01%**.
- **The success of our work so far encourages to look in more detail into the use of Tensor Network methods in lattice gauge theories.**



Prospects




- We would like to look into aspects of lattice gauge theories where the standard methods have problems:
 - ★ thermodynamics at non-zero chemical potential,
 - ★ non-equilibrium properties.



[K. Fukushima, T. Hatsuda, Rep. Prog. Phys. 74 (2011) 14001]

- First attempts already underway: computation of the chiral condensate at finite temperature.
- **Ultimate aim: full QCD**, i.e.:
 - ★ a non-Abelian theory (with $SU(3)$ gauge group),
 - ★ in $3+1$ dimensions.
- Needs **a lot** of work of the Tensor Network + lattice gauge theory community...



Overview
Seminar outline

Introduction

Results

Summary

**Thank you for your
attention!**

Thank you for your attention!