

# Triplet extension of the MSSM: alignment, loop-induced Higgs decays and dark matter

**Germano Nardini**  
(DESY)

*Lab. de Physique Subatomique et de Cosmologie*  
*18/11/2014*

Based on works done with  
C. Arina, A. Delgado, V. Martin-Lozano, M. Quiros

# Outline

## 1 Introduction

- The model

## 2 Higgs Phenomenology

- Features at small  $m_A$
- Features at large  $m_A$ 
  - Without dark matter requests
  - With dark matter requests

## 3 Conclusion

# Motivation

- No clear discrepancies between LHC data and SM predictions with  $m_h \simeq 125 \text{ GeV}$ . And BAU and DM imply just “some new fields”
- If we do not give up with the (Planck - EW) hierarchy problem, **SUSY** is (one of) the favourite UV option
- In the **MinimalSSM**  $m_h \simeq 125 \text{ GeV}$  requires “heavy” stop sector  $\Rightarrow$  **Little Hierarchy Problem**, i.e. some fine-tuning
- Non-minimalSSM models can alleviate this problem as they can enhance the tree-level Higgs mass via
  - D-terms: Extra gauge interactions
  - F-terms: Extra chiral sector (singlets and/or triplets)

## THE $Y = 0$ TRIPLET

Less free parameters than  $Y = \pm 1$  extension, and extra charginos with collider and cosmological effects

(other tripl. extens. in E.J.Chun&al,1209.1303; Z.Kang&al,1301.2204; M.Garcia&al,1308.4025)

# The $Y=0$ Triplet Extension (Espinosa&Quiros'92, Di Chiara&Hsieh'08)

$$\Sigma = \begin{pmatrix} \xi^0/\sqrt{2} & -\xi_2^+ \\ \xi_1^- & -\xi^0/\sqrt{2} \end{pmatrix}, \quad \Delta W = \lambda H_1 \cdot \Sigma H_2 + \frac{1}{2} \mu_\Sigma \text{tr} \Sigma^2 + \mu H_1 \cdot H_2$$

- $T$  parameter bound requires  $\langle \xi^0 \rangle \lesssim 4 \text{ GeV}$  which imposes (unless of fine-tuning)

$$|A_\lambda|, |\mu|, |\mu_\Sigma| \lesssim \frac{m_\Sigma^2 + \lambda^2 v^2 / 2}{10^2 \lambda v}$$

- This hierarchy implies decoupling between  $\xi^0$  and  $H_1, H_2$

$$\text{Mass boost: } V(H_1, H_2) \simeq V_{MSSM} + \lambda^2 |H_1^0 H_2^0|^2$$

$$\begin{aligned} (\text{for } m_A \rightarrow \infty: \quad m_h^2 &= m_Z^2 \cos^2 2\beta + \lambda^2 v^2 \sin^2 2\beta / 2) \\ &(\text{large } \lambda \text{ and small } \tan \beta \text{ preferred}) \end{aligned}$$

# The relevant spectrum

- Heavy scalar triplet [  $\gtrsim 5 \text{ TeV}$  ]
- Minimiz. conditions

$$m_3^2 = m_A^2 \sin \beta \cos \beta , \quad m_Z^2 = \frac{m_2^2 - m_1^2}{\cos 2\beta} - m_A^2 + \frac{\lambda^2}{2} v^2$$

- CP-odd/charged Higgses

$$m_A^2 = m_1^2 + m_2^2 + 2|\mu|^2 + \frac{\lambda^2}{2} v^2 , \quad m_{H^\pm}^2 = m_A^2 + m_W^2 + \frac{\lambda^2}{2} v^2$$

# The relevant spectrum

- CP-even Higgs masses (basis  $h_2, h_1$ )

$$\mathcal{M}_0^2 = \begin{pmatrix} m_A^2 \cos^2 \beta + m_Z^2 \sin^2 \beta & (\lambda^2 v^2 - m_A^2 - m_Z^2) \sin 2\beta/2 \\ (\lambda^2 v^2 - m_A^2 - m_Z^2) \sin 2\beta/2 & m_A^2 \sin^2 \beta + m_Z^2 \cos^2 \beta \end{pmatrix}$$

- After including radiative corrections  $\Delta\mathcal{M}_t^2(h_t)$  and  $\Delta\mathcal{M}_\Sigma^2(\lambda)$  (also in the min.condts) [ $m_h = 125 \text{ GeV}$ ]

$$\begin{pmatrix} h_2 \\ h_1 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} h \\ H \end{pmatrix}$$

Coupling ratios  $r_{\mathcal{H}XX} = g_{\mathcal{H}XX}/g_{hXX}^{\text{SM}}$  ( $\mathcal{H} = h, H$ )

|                        |                        |                                  |                                  |                                   |                                  |
|------------------------|------------------------|----------------------------------|----------------------------------|-----------------------------------|----------------------------------|
| $r_{hVV}^0$            | $r_{HVV}^0$            | $r_{htt}^0$                      | $r_{Htt}^0$                      | $r_{hdd}^0$                       | $r_{Hdd}^0$                      |
| $\sin(\beta - \alpha)$ | $\cos(\beta - \alpha)$ | $\frac{\cos \alpha}{\sin \beta}$ | $\frac{\sin \alpha}{\sin \beta}$ | $-\frac{\sin \alpha}{\cos \beta}$ | $\frac{\cos \alpha}{\cos \beta}$ |

# The relevant spectrum

- CP-even Higgs masses (basis  $h_2, h_1$ )

$$\mathcal{M}_0^2 = \begin{pmatrix} m_A^2 \cos^2 \beta + m_Z^2 \sin^2 \beta & (\lambda^2 v^2 - m_A^2 - m_Z^2) \sin 2\beta/2 \\ (\lambda^2 v^2 - m_A^2 - m_Z^2) \sin 2\beta/2 & m_A^2 \sin^2 \beta + m_Z^2 \cos^2 \beta \end{pmatrix}$$

## SM-LIKE INTERACTIONS WHEN $\alpha \simeq \beta - \pi/2$

Case 1:  $\beta \simeq \beta_c = \pi/4$ ,  $\lambda \simeq \lambda_c = \sqrt{2}m_h/v$  (relevant but accidental)

Case 2:  $m_A \gg m_h$  (robust but obvious)

Coupling ratios  $r_{\mathcal{H}XX} = g_{\mathcal{H}XX}/g_{hXX}^{\text{SM}}$  ( $\mathcal{H} = h, H$ )

| $r_{hVV}^0$            | $r_{HVV}^0$            | $r_{htt}^0$                      | $r_{Htt}^0$                      | $r_{hdd}^0$                       | $r_{Hdd}^0$                      |
|------------------------|------------------------|----------------------------------|----------------------------------|-----------------------------------|----------------------------------|
| $\sin(\beta - \alpha)$ | $\cos(\beta - \alpha)$ | $\frac{\cos \alpha}{\sin \beta}$ | $\frac{\sin \alpha}{\sin \beta}$ | $-\frac{\sin \alpha}{\cos \beta}$ | $\frac{\cos \alpha}{\cos \beta}$ |

Small  $m_A$ 

## Case 1

- Existence of the SM-like point beyond tree-level approx
- Quantifying the “ $\simeq$ ”
- Some phenomenology

Delgado, GN, Quiros, ArXiv:1303.0800

*Similar idea in the MSSM for some radiative corrections:*  
*Carena&al. [ArXiv:1310.2248](#)*

*Long ago for 2HDM: Haber and Gunion, [hep-ph/0207010](#)*



Small  $m_A$ 

(similar idea in Carena&amp;al 1310:2248 for 2HDM)

- CP-even Higgs masses (basis  $h_2, h_1$ )

$$\mathcal{M}_0^2 = \begin{pmatrix} m_A^2 \cos^2 \beta + m_Z^2 \sin^2 \beta & (\lambda^2 v^2 - m_A^2 - m_Z^2) \sin 2\beta/2 \\ (\lambda^2 v^2 - m_A^2 - m_Z^2) \sin 2\beta/2 & m_A^2 \sin^2 \beta + m_Z^2 \cos^2 \beta \end{pmatrix}$$

$$\begin{pmatrix} h_2 \\ h_1 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} h \\ H \end{pmatrix}$$

- Tree-level  $h$  couplings are SM-like if  $\alpha = \beta - \pi/2$ . With  $\mathcal{M}_0^2$ :

$$\beta_c = \frac{\pi}{4}, \quad \lambda_c = \sqrt{2} \frac{m_h}{v}$$

The lightest eigenvalue is INDEPENDENT of  $m_A$



AND BEYOND TREE LEVEL ?

# Small $m_A$

Radiative correction  $\Delta\mathcal{M}_t^2(h_t)$  and  $\Delta\mathcal{M}_\Sigma^2(\lambda)$  included ( $m_Q = m_U = 700$  GeV,  $A_t = 0$ ,  $m_\Sigma = 5$  TeV).

Curves:  $m_A = \infty, 200, 155, 145, 140, 135, 130$  GeV

$\tan\beta(\lambda)$  and  $\sin\alpha(\lambda)$  fixing  $m_h = 125$  GeV



THE SM-LIKE CONDITION HAS MOVED TO THE  
CROSSING POINT

# Small $m_A$

(similar idea in Carena&al 1310:2248 for 2HDM)

- Given a spectrum and its dominant stop and triplet corrections ...

$$\mathcal{M}_1^2 = \begin{pmatrix} m_A^2 \cos^2 \beta + m_{11}^2 \sin^2 \beta & (-m_A^2 - m_{12}^2) \sin 2\beta/2 \\ (-m_A^2 - m_{12}^2) \sin 2\beta/2 & m_A^2 \sin^2 \beta + m_{22}^2 \cos^2 \beta \end{pmatrix}$$

- ... the lightest eigenvalue  $m_h$  is equal to  $\overline{m}_h$  when

$$\mathcal{D} - \overline{m}_h^2 \mathcal{T} + \overline{m}_h^4 = 0$$

$$\Downarrow$$

$$A(\tan \beta, \lambda; \overline{m}_h^2) m_A^2 + B(\tan \beta, \lambda; \overline{m}_h^2) = 0 \quad (\text{no } m_A^4!),$$

- there exists a solution that is INDEPENDENT of  $m_A$  at

$$\begin{cases} A(\tan \beta_c, \lambda_c, \overline{m}_h^2) = 0 \\ B(\tan \beta_c, \lambda_c, \overline{m}_h^2) = 0 \end{cases}$$

- Moreover (analytically) when the lightest eigenvalue is independent of  $m_A$ ,  $\alpha$  is independent as well.
- Since independent of  $m_A$ , the particular case  $m_A \rightarrow \infty$  is included, i.e.  $\alpha = \beta - \pi/2$

Small  $m_A$ Signal strengths  $\mathcal{R}_{\mathcal{H}XX}$ 

$$\mathcal{R}_{\mathcal{H}XX} = \frac{\sigma(pp \rightarrow \mathcal{H}) BR(\mathcal{H} \rightarrow XX)}{[\sigma(pp \rightarrow h) BR(h \rightarrow XX)]_{SM}}$$

$$\mathcal{R}_{\mathcal{H}XX}^{(ggF)} = \mathcal{R}_{\mathcal{H}XX}^{(\mathcal{H}tt)} = \frac{r_{\mathcal{H}tt}^2 r_{\mathcal{H}XX}^2}{\mathcal{D}}, \quad \mathcal{R}_{\mathcal{H}XX}^{(VBF)} = \mathcal{R}_{\mathcal{H}XX}^{(V\mathcal{H})} = \frac{r_{\mathcal{H}WW}^2 r_{\mathcal{H}XX}^2}{\mathcal{D}}$$

$$\begin{aligned} \mathcal{D} = & BR(h \rightarrow b\bar{b})_{SM} r_{\mathcal{H}bb}^2 + BR(h \rightarrow gg, cc)_{SM} r_{\mathcal{H}tt}^2 \\ & + BR(h \rightarrow \tau\tau)_{SM} r_{\mathcal{H}\tau\tau}^2 + BR(h \rightarrow WW, ZZ)_{SM} r_{\mathcal{H}WW}^2 \end{aligned}$$


- No extra inv. width, i.e.  $m_{\chi_0} \gtrsim m_{\mathcal{H}}/2$
- sbottom-gluino may correct  $r_{hbb}$  ( $M_3 = 1 \text{ TeV}, m_{\tilde{g}} = 700 \text{ GeV}$ )

Small  $m_A$ 

$$m_A = 140 \text{ GeV}, \mu = \mu_\Sigma = 250 \text{ GeV}, m_{\chi^\pm} = 104 \text{ GeV}$$

Signal strengths  $\mathcal{R}_{hXX}$  from ggF and tt

$$\mathcal{R}_{\mathcal{H}XX} = \frac{\sigma(pp \rightarrow \mathcal{H}) BR(\mathcal{H} \rightarrow XX)}{[\sigma(pp \rightarrow h) BR(h \rightarrow XX)]_{SM}}$$




Small  $m_A$ 

$$m_A = 140 \text{ GeV}, \mu = \mu_\Sigma = 250 \text{ GeV}, m_{\chi^\pm} = 104 \text{ GeV}$$

Signal strengths  $\mathcal{R}_{hXX}$  from VBF and Vh

$$\mathcal{R}_{\mathcal{H}XX} = \frac{\sigma(pp \rightarrow \mathcal{H}) BR(\mathcal{H} \rightarrow XX)}{[\sigma(pp \rightarrow h) BR(h \rightarrow XX)]_{SM}}$$




Small  $m_A$ 

$$m_A = 140 \text{ GeV}, \mu = \mu_\Sigma = 250 \text{ GeV}, m_{\chi^\pm} = 104 \text{ GeV}$$

Signal strengths  $\mathcal{R}_{HXX}$  from ggF and htt

$$\mathcal{R}_{\mathcal{H}XX} = \frac{\sigma(pp \rightarrow \mathcal{H}) BR(\mathcal{H} \rightarrow XX)}{[\sigma(pp \rightarrow h) BR(h \rightarrow XX)]_{SM}}$$



- Further reduction if  $m_{\chi_0} \lesssim m_H/2$  ( $m_H \sim 138 \text{ GeV}$ )

# Small $m_A$

## Mini Summary

Exp. results on Higgs decays are hinting to SM-like values. If confirmed with much better accuracy, they still don't imply  $m_A \gg m_h$

For instance in the TMSSM there exists a parameter region where

- we have both  $m_h \approx 125$  GeV and SM-like  $h$  decay independently of  $m_A$ ;
- $H$  production is suppressed

To probe the scenario, present LHC studies on  $A$  and  $H^\pm$  decays need to be improved (in the TMSSM these limits seem  $\lesssim$  MSSM or NMSSM ones).

In this direction:

Bandyopadhyay&Huitu&Sabanci'13, Bandyopadhyay&Di Chiara&al,'14



Large  $m_A$ 

## Case 2

- Diphoton enhancement highlighted before
- Correlation with  $h \rightarrow Z\gamma$
- Imposing DM

Delgado, GN, Quiros, ArXiv:1207.6596  
Arina, GN, Martin-Lozano, ArXiv:1403.6434

# Enhancement in $h \rightarrow \gamma\gamma$

Signal strength  $\mathcal{R}_{\gamma\gamma}$   $[= BR(h \rightarrow \gamma\gamma)/BR(h \rightarrow \gamma\gamma)_{SM}]$

$$A_{\tilde{\chi}_{1,2,3}^{\pm}}^{\gamma\gamma} = \left(-\frac{4}{3}\right) \frac{v^2(\lambda^2 M_2 + g^2 \mu_\Sigma) \sin 2\beta}{M_2 \mu \mu_\Sigma - \frac{1}{2} \lambda^2 v^2 (\lambda^2 M_2 + g^2 \mu_\Sigma) \sin 2\beta}$$

- Loop-induced process which is sensitive to new charged particles
- New triplet charged fermion can enhance  $\mathcal{R}_{\gamma\gamma}$  ( $\lesssim 1.2$  via MSSM charginos; e.g. Casas, Moreno, Roliecki, Zaldivar '13)
- As no (large) modifications in the Higgs production exist for  $m_A \rightarrow \infty$ , the diphoton enhancement is (easily implemented from the QED eff.pot. Ellis&al,79; Shifman&al,79; Carena&al,12)

$$\mathcal{R}_{\gamma\gamma} = \left| 1 + \frac{A_{\tilde{\chi}_{1,2,3}^{\pm}}^{\gamma\gamma}}{A_W^{\gamma\gamma} + A_t^{\gamma\gamma}} \right|^2$$

# Enhancement in $h \rightarrow \gamma\gamma$

Signal strength  $\mathcal{R}_{\gamma\gamma}$   $[= BR(h \rightarrow \gamma\gamma)/BR(h \rightarrow \gamma\gamma)_{SM}]$

$$A_{\tilde{\chi}_{1,2,3}^\pm}^{\gamma\gamma} = \left(-\frac{4}{3}\right) \frac{\partial}{\partial \log v} \log(\det M_{\tilde{\chi}^\pm}^{tree})$$

$$\mathcal{M}_{\tilde{\chi}^\pm}^{tree} = \begin{pmatrix} M_2 & gv \sin \beta & 0 \\ gv \cos \beta & \mu & -\lambda v \sin \beta \\ 0 & \lambda v \cos \beta & \mu_\Sigma \end{pmatrix}, \quad UM_{\tilde{\chi}^\pm}V^\dagger = diag$$


- As no (large) modifications in the Higgs production exist for  $m_A \rightarrow \infty$ , the diphoton enhancement is (easily implemented from the QED eff.pot. Ellis&al,79; Shifman&al,79; Carena&al,12)

$$R_{\gamma\gamma} = \left| 1 + \frac{A_{\tilde{\chi}_{1,2,3}^\pm}^{\gamma\gamma}}{A_W^{\gamma\gamma} + A_t^{\gamma\gamma}} \right|^2$$

# Enhancement in $h \rightarrow \gamma\gamma$

Signal strength  $\mathcal{R}_{\gamma\gamma} \quad [= BR(h \rightarrow \gamma\gamma)/BR(h \rightarrow \gamma\gamma)_{SM}]$


$$A_{\tilde{\chi}_{1,2,3}^{\pm}}^{\gamma\gamma} = \left(-\frac{4}{3}\right) \frac{v^2(\lambda^2 M_2 + g^2 \mu_\Sigma) \sin 2\beta}{M_2 \mu \mu_\Sigma - \frac{1}{2} \lambda^2 v^2 (\lambda^2 M_2 + g^2 \mu_\Sigma) \sin 2\beta}$$



$M_2 = 250 \text{ GeV}$

$X_t = 4, 0; m_Q = 700 \text{ GeV}$

$\mu = \mu_\Sigma \rightarrow m_{\tilde{\chi}_1^\pm} \sim 105 \text{ GeV}$



$M_2 = 750 \text{ GeV}$

Correlation  $h \rightarrow \gamma\gamma$  vs  $h \rightarrow Z\gamma$ Signal strength  $\mathcal{R}_{Z\gamma}$   $[ = \mathcal{BR}(h \rightarrow Z\gamma) / \mathcal{BR}(h \rightarrow Z\gamma)_{\text{SM}} ]$ 

$$A_{\tilde{\chi}_{1,2,3}^{\pm}}^{Z\gamma} = \sum_{j,k=1}^3 \frac{g_2 m_{\tilde{\chi}_j^{\pm}}}{g_1 m_Z} g_{Z\tilde{\chi}_j^+ \tilde{\chi}_i^-} f(m_{\tilde{\chi}_j^{\pm}}, m_{\tilde{\chi}_k^{\pm}}, m_{\tilde{\chi}_k^{\pm}}) g_{h\tilde{\chi}_j^+ \tilde{\chi}_i^-}$$


- Weak bounds by LHC but planned improvements
- Typically correlated to  $h \rightarrow \gamma\gamma$
- TMSSM chargino sector should play a role (as for  $R_{\gamma\gamma}$ )
- Less transparent expression because no applicable low-energy limit

$$R_{\gamma\gamma} = \left| 1 + \frac{A_{\tilde{\chi}_{1,2,3}^{\pm}}^{Z\gamma}}{A_W^{Z\gamma} + A_t^{Z\gamma}} \right|^2$$


# Correlation $h \rightarrow \gamma\gamma$ vs $h \rightarrow Z\gamma$ ( $m_{\tilde{\chi}_1^\pm} \gtrsim 100$ GeV)

Signal strength  $\mathcal{R}_{Z\gamma}$  [ $= BR(h \rightarrow Z\gamma)/BR(h \rightarrow Z\gamma)_{SM}$ ]

$$A_{\tilde{\chi}_{1,2,3}^\pm}^{Z\gamma} = \sum_{j,k=1}^3 \frac{g_2 m_{\tilde{\chi}_j^\pm}}{g_1 m_Z} g_{Z\tilde{\chi}_j^+ \tilde{\chi}_i^-} f(m_{\tilde{\chi}_j^\pm}, m_{\tilde{\chi}_k^\pm}, m_{\tilde{\chi}_k^\pm}) g_{h\tilde{\chi}_j^+ \tilde{\chi}_i^-}$$



solid:  $\mu_\Sigma = \mu = M_2 = 230$  GeV  
 dash:  $\mu_\Sigma = \mu = 230$  GeV,  $M_2 = 1$  TeV  
 dot-dash:  $\mu_\Sigma = M_2 = 230$  GeV,  $\mu = 400$  GeV




$A_t = 0$ ;  $m_Q = m_U$  adjusted for  $m_h = 125$  GeV

# Correlation $h \rightarrow \gamma\gamma$ vs $h \rightarrow Z\gamma$ ( $m_{\tilde{\chi}_1^\pm} \gtrsim 100$ GeV)

Signal strength  $\mathcal{R}_{Z\gamma}$  [ $= \overline{BR}(h \rightarrow Z\gamma) / \overline{BR}(h \rightarrow Z\gamma)_{\text{SM}}$ ]


$$A_{\tilde{\chi}_{1,2,3}^\pm}^{Z\gamma} = \sum_{j,k=1}^3 \frac{g_2 m_{\tilde{\chi}_j^\pm}}{g_1 m_Z} g_{Z\tilde{\chi}_j^+ \tilde{\chi}_i^-} f(m_{\tilde{\chi}_j^\pm}, m_{\tilde{\chi}_k^\pm}, m_{\tilde{\chi}_k^\pm}) g_{h\tilde{\chi}_j^+ \tilde{\chi}_i^-}$$



solid:  $\mu_\Sigma = \mu = M_2 = 230$  GeV

dash:  $\mu_\Sigma = \mu = 230$  GeV,  $M_2 = 1$  TeV

dot-dash:  $\mu_\Sigma = M_2 = 230$  GeV,  $\mu = 400$  GeV



$0.01 < \text{BR}(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0) < 0.2$

$0.2 < \text{BR}(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0) < 0.5$


$0.5 < \text{BR}(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0)$

# Correlation $h \rightarrow \gamma\gamma$ vs $h \rightarrow Z\gamma$ ( $m_{\chi_1^\pm} \gtrsim 100$ GeV)

$$\text{BR}(h \rightarrow \chi_1^0 \chi_1^0)$$

$$\Gamma(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0) = \frac{G_F m_W^2}{2\sqrt{2}\pi} m_h \left(1 - \frac{4m_{\tilde{\chi}_1^0}^2}{m_h^2}\right)^{3/2} g_{h11}^2$$

$$\mathcal{M}_{\chi^0} = \begin{pmatrix} M_1 & 0 & -\frac{1}{2}g_1 v_1 & \frac{1}{2}g_1 v_2 & 0 \\ \cdot & M_2 & \frac{1}{2}g_2 v_1 & -\frac{1}{2}g_2 v_2 & 0 \\ \cdot & \cdot & 0 & -\mu & -\frac{1}{2}v_2 \lambda \\ \cdot & \cdot & \cdot & 0 & -\frac{1}{2}v_1 \lambda \\ \cdot & \cdot & \cdot & \cdot & \mu \Sigma \end{pmatrix}$$





$$0.01 < \text{BR}(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0) < 0.2$$

$$0.2 < \text{BR}(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0) < 0.5$$

$$0.5 < \text{BR}(h \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0)$$



Again  $h \rightarrow \gamma\gamma$ 

$$R_{\gamma\gamma} < 1.1, 1.2, 1.3, 1.4 \quad R_{\gamma\gamma} > 1.4$$

If  $\mu$  large,  $R_{\gamma\gamma}$  small


This seems to be strictly linked to Dark Matter!!!

# If DM is the Bino-like neutralino

Framework is like well-tempered neutralino:  $\Omega_{DM}$  relies on SM + ewkinos  
 If  $m_A$  or  $m_L$  are small, there are potential new channels.

| Observable                    | Measured/Limit                                |
|-------------------------------|---|
| $\sigma_{Xe}^{SI}$            | LUX (90% CL)                                  |
| $\Omega_{DM} h^2$             | $0.1186 \pm 0.0031$ (exp) $\pm 20\%$ (theo)   |
| $m_h$                         | $125.85 \pm 0.4$ GeV (exp) $\pm 3$ GeV (theo) |
| $\Gamma_Z^{\text{invisible}}$ | $(166 \pm 2)$ MeV                             |
| $m_{\tilde{t}_1}$             | $> 650$ GeV (LHC 90% CL)                      |
| $m_{\tilde{\chi}_1^+}$        | $> 104$ GeV (LEP 95% CL)                      |

# If DM is the Bino-like neutralino (LUX bound)





(dangerous) SI cross section is dominated by Higgs interchange ( $g_{h\tilde{\chi}_1^0\chi_1^0}$ )

For a given parameter set, LUX  $\Rightarrow$  lower bound on  $\mu$   
 $\Rightarrow$  upper bound on  $R_{\gamma\gamma}$  and  $R_{Z\gamma}$

# If DM is the Bino-like neutralino

# (well-tempered)




 $M_2 = 1 \text{ TeV}$ 
 $\tan \beta = 2.8$ 
 $\lambda = 0.85$ 

Good relic density if

*Branch 1:* Triplino-Bino coannihilation (driven by gauge interactions)

*Branch 2:* Higgs/Z resonance

# If DM is the Bino-like neutralino *(resonance)*





$$M_2 = 1.5 \text{ TeV}$$

$$\tan \beta = 2.8$$

$$\lambda = 0.85$$

LUX goes better when diphoton enhancement is smaller



# If DM is the Bino-like neutralino *(resonance)*



$$M_2 = 1.5 \text{ TeV} \quad \tan \beta = 2.8 \quad \lambda = 0.85$$

Quantitatively, how much  $R_{\gamma\gamma}$  (and  $R_{Z\gamma}$ ) is allowed by LUX?

# If DM is the Bino-like neutralino



$$R_{\gamma\gamma} \lesssim 1.25$$

$$R_{Z\gamma} \lesssim 1.2$$

(correlated!)

# Conclusion

- ❶ Triplet extension alleviates the fine-tuning with respect to the MSSM
- ❷ SM-like Higgs signatures do NOT imply large  $m_A$
- ❸ If the dominant channels are SM-like, TMSSM chargino sector can provide  $R_{\gamma\gamma}$  and  $R_{Z\gamma}$  are large as
  - 1.6 and 1.4 (no DM)
  - 1.3 and 1.2 (with DM)
- ❹  $R_{\gamma\gamma}$  and  $R_{Z\gamma}$  are strongly correlated
- ❺ Concerning well tempering, TMSSM DM easier than MSSM DM from top-down approach
- ❻ Open issues: LHC bounds? EWino composition via ILC?  
De Blas, Delgado, Ostidek, Quiros, '14 Moortgat-Pick, Porto, Rolbiecki, '14