

Guiding stars for physics beyond SM: Higgs boson and dark matter

朱守华, Shou-hua Zhu
ITP, Peking University

January 9, 2008

Brief Introduction to Institute of Theoretical Physics (ITP) of Peking University

String Theory and Cosmology ($>$ or \gg TeV):

① Bin Chen

TeV Physics (LHC/ILC energy scale):

- 1 Chong Sheng Li
- 2 Da-xin Zhang
- 3 Shou-hua Zhu

Hadronic Physics (GeV):

- ① Kuang-ta Chao (director)
- ② Chuan Liu
- ③ Bo-qiang Ma
- ④ Han-qing Zheng
- ⑤ Shi-lin Zhu

Nuclear Physics (MeV):

- 1 Wei-zhen Deng
- 2 Yi-an Lei
- 3 Yu-xin Liu

Condensed Matter Physics (Many-body, eV):

- ① Ding-ping Li
- ② Zhong-shui Ma

Contents

- 1 Introduction
- 2 Invisible SM-like Higgs boson due to X(214) and analysis fault
- 3 Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter
- 4 Higgs boson as the looking glass in mirror model
- 5 Conclusion and discussion

Contents

- 1 Introduction
- 2 Invisible SM-like Higgs boson due to X(214) and analysis fault
- 3 Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter
- 4 Higgs boson as the looking glass in mirror model
- 5 Conclusion and discussion

Contents

- 1 Introduction
- 2 Invisible SM-like Higgs boson due to X(214) and analysis fault
- 3 Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter
- 4 Higgs boson as the looking glass in mirror model
- 5 Conclusion and discussion

Contents

- 1 Introduction
- 2 Invisible SM-like Higgs boson due to X(214) and analysis fault
- 3 Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter
- 4 Higgs boson as the looking glass in mirror model
- 5 Conclusion and discussion

Contents

- 1 Introduction
- 2 Invisible SM-like Higgs boson due to X(214) and analysis fault
- 3 Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter
- 4 Higgs boson as the looking glass in mirror model
- 5 Conclusion and discussion

Based on

- W. s. Li, P. f. Yin and S. H. Zhu, Phys. Rev. D **76**, 095012 (2007) [arXiv:0709.1586 [hep-ph]].
- S. H. Zhu, Chin. Phys. Lett. **24**, 381 (2007), arXiv:hep-ph/0601224.
- S. h. Zhu, Phys. Rev. D **75**, 115004 (2007) [arXiv:hep-ph/0701001].
- P.f. Yin and S. h. Zhu, arXiv:hep-ph/0611270.
- S. h. Zhu, Eur. Phys. J. C **47**, 833 (2006) [arXiv:hep-ph/0512055].

Introduction

- Higgs sector is the least known part in SM
- Dark matter is required by cosmological observations!
- Invisible Higgs boson (漆黑格斯粒子)
- Higgs boson and/or dark matter as the guiding stars!

Introduction

- Higgs sector is the least known part in SM
- Dark matter is required by cosmological observations!
- Invisible Higgs boson (漆黑格斯粒子)
- Higgs boson and/or dark matter as the guiding stars!

1 Not found yet.

2 Direct limit 114 GeV from LEP at CERN.

- Q: What is direct limit?
- A: Limit from experiments in which Higgs boson is supposed to be produced directly.

3 Indirect information from quantum fluctuations and screening theorem: precision observables are only sensitive to $\log(m_H)$ for leading quantum fluctuation effects.

- Q: What is indirect limit?
- A: Limit from experiments in which Higgs boson can't be produced directly, namely shows up only as virtual states.

- ① Not found yet.
- ② Direct limit 114 GeV from LEP at CERN.
 - Q: What is direct limit?
 - A: Limit from experiments in which Higgs boson is supposed to be produced directly.
- ③ Indirect information from quantum fluctuations and screening theorem: precision observables are only sensitive to $\log(mH)$ for leading quantum fluctuation effects.
 - Q: What is indirect limit?
 - A: Limit from experiments in which Higgs boson can't be produced directly, namely shows up only as virtual states.


- 1 Not found yet.
- 2 Direct limit 114 GeV from LEP at CERN.
 - Q: What is direct limit?
 - A: Limit from experiments in which Higgs boson is supposed to be produced directly.
- 3 Indirect information from quantum fluctuations and screening theorem: precision observables are only sensitive to $\log(m_H)$ for leading quantum fluctuation effects.
 - Q: What is indirect limit?
 - A: Limit from experiments in which Higgs boson can't be produced directly, namely shows up only as virtual states.

- ① Not found yet.
- ② Direct limit 114 GeV from LEP at CERN.
 - Q: What is direct limit?
 - A: Limit from experiments in which Higgs boson is supposed to be produced directly.
- ③ Indirect information from quantum fluctuations and screening theorem: precision observables are only sensitive to $\log(m_H)$ for leading quantum fluctuation effects.
 - Q: What is indirect limit?
 - A: Limit from experiments in which Higgs boson can't be produced directly, namely shows up only as virtual states.

- ① Not found yet.
- ② Direct limit 114 GeV from LEP at CERN.
 - Q: What is direct limit?
 - A: Limit from experiments in which Higgs boson is supposed to be produced directly.
- ③ Indirect information from quantum fluctuations and screening theorem: precision observables are only sensitive to $\log(mH)$ for leading quantum fluctuation effects.
 - Q: What is indirect limit?
 - A: Limit from experiments in which Higgs boson can't be produced directly, namely shows up only as virtual states.

- ① Not found yet.
- ② Direct limit 114 GeV from LEP at CERN.
 - Q: What is direct limit?
 - A: Limit from experiments in which Higgs boson is supposed to be produced directly.
- ③ Indirect information from quantum fluctuations and screening theorem: precision observables are only sensitive to $\log(mH)$ for leading quantum fluctuation effects.
 - Q: What is indirect limit?
 - A: Limit from experiments in which Higgs boson can't be produced directly, namely shows up only as virtual states.

Consistent of direct and indirect limits?



Introduction

- Higgs sector is the least known part in SM
- **Dark matter is required by cosmological observations!**
- Invisible Higgs boson (漆黑格斯粒子)
- Higgs boson and/or dark matter as the guiding stars!

- Q: Is dark matter relevant to weak scale physics?
- A: Likely because the relic density of dark matter can be naturally correlated with weak coupling α and weak scale 100 GeV.
- Q: If yes, where to insert dark matter sector?
- A: Most likely in Higgs sector because success of standard model of particle physics permits naturally the additional sector in Higgs part.

- Q: Is dark matter relevant to weak scale physics?
- A: Likely because the relic density of dark matter can be naturally correlated with weak coupling α and weak scale 100 GeV.
- Q: If yes, where to insert dark matter sector?
- A: Most likely in Higgs sector because success of standard model of particle physics permits naturally the additional sector in Higgs part.

- Q: Is dark matter relevant to weak scale physics?
- A: Likely because the relic density of dark matter can be naturally correlated with weak coupling α and weak scale 100 GeV.
- Q: If yes, where to insert dark matter sector?
- A: Most likely in Higgs sector because success of standard model of particle physics permits naturally the additional sector in Higgs part.

- Q: Is dark matter relevant to weak scale physics?
- A: Likely because the relic density of dark matter can be naturally correlated with weak coupling α and weak scale 100 GeV.
- Q: If yes, where to insert dark matter sector?
- A: Most likely in Higgs sector because success of standard model of particle physics permits naturally the additional sector in Higgs part.

Introduction

- Higgs sector is the least known part in SM
- Dark matter is required by cosmological observations!
- **Invisible Higgs boson (漆黑格斯粒子)**
- Higgs boson and/or dark matter as the guiding stars!

- Higgs boson may not be observed (invisible) in the modern detectors at colliders!
- What is the meaning of 'invisible decay'?
- For modern detectors, some particles which do not interact with the detector will appear as invisible signal, for example neutrino in the SM.
- Cold dark matter, which interacts only weakly with usual matter in detector, appears as invisible signals.
- Higgs boson, which are produced at colliders, may decay mainly into dark matter. Thus Higgs boson appears as invisible particle also.

- Higgs boson may not be observed (invisible) in the modern detectors at colliders!
- What is the meaning of 'invisible decay'?
- For modern detectors, some particles which do not interact with the detector will appear as invisible signal, for example neutrino in the SM.
- Cold dark matter, which interacts only weakly with usual matter in detector, appears as invisible signals.
- Higgs boson, which are produced at colliders, may decay mainly into dark matter. Thus Higgs boson appears as invisible particle also.

- Higgs boson may not be observed (invisible) in the modern detectors at colliders!
- What is the meaning of 'invisible decay'?
- For modern detectors, some particles which do not interact with the detector will appear as invisible signal, for example neutrino in the SM.
- Cold dark matter, which interacts only weakly with usual matter in detector, appears as invisible signals.
- Higgs boson, which are produced at colliders, may decay mainly into dark matter. Thus Higgs boson appears as invisible particle also.

- Higgs boson may not be observed (invisible) in the modern detectors at colliders!
- What is the meaning of 'invisible decay'?
- For modern detectors, some particles which do not interact with the detector will appear as invisible signal, for example neutrino in the SM.
- Cold dark matter, which interacts only weakly with usual matter in detector, appears as invisible signals.
- Higgs boson, which are produced at colliders, may decay mainly into dark matter. Thus Higgs boson appears as invisible particle also.

- Higgs boson may not be observed (invisible) in the modern detectors at colliders!
- What is the meaning of 'invisible decay'?
- For modern detectors, some particles which do not interact with the detector will appear as invisible signal, for example neutrino in the SM.
- Cold dark matter, which interacts only weakly with usual matter in detector, appears as invisible signals.
- Higgs boson, which are produced at colliders, may decay mainly into dark matter. Thus Higgs boson appears as invisible particle also.

- Besides dark matter, the Higgs boson can be invisible due to other reasons (see next part)!

Introduction

- Higgs sector is the least known part in SM
- Dark matter is required by cosmological observations!
- Invisible Higgs boson (漆黑格斯粒子)
- Higgs boson and/or dark matter as the guiding stars!

Introduction

Invisible SM-like Higgs boson due to X(214) and analysis fault
Correlation between masses of SM-like Higgs boson and (singlet) scalar
Higgs boson as the looking glass in mirror model
Conclusion and discussion

Higgs sector is the least known part in SM
Dark matter is required by cosmological observations!
Invisible Higgs boson (漆黑格斯粒子)
Higgs boson and/or dark matter as the guiding stars!

The investigation on Higgs boson and/or dark matter can provide crucial information for deepening our understanding of the nature!

Invisible SM-like Higgs boson due to X(214) and analysis fault

- 'SM-like' Higgs boson
- HyperCP three events
- A new pseudoscalar X(214 MeV)?
- All about X(214)
- Consequences on Higgs physics

Invisible SM-like Higgs boson due to X(214) and analysis fault

- 'SM-like' Higgs boson
- HyperCP three events
- A new pseudoscalar X(214 MeV)?
- All about X(214)
- Consequences on Higgs physics

- What is the 'SM-like' Higgs boson?
- A: We name the Higgs boson as 'SM-like' in case that one of the Higgs bosons in physics beyond the SM (for example MSSM) acts like the Higgs boson in the SM.
- Why 'SM-like' Higgs boson?
- Because the SM is extremely success, the existence of one 'SM-like' Higgs boson is the simplest way to coincide the data.
- Other crazy scenarios are, of course, allowed!

- What is the 'SM-like' Higgs boson?
- A: We name the Higgs boson as 'SM-like' in case that one of the Higgs bosons in physics beyond the SM (for example MSSM) acts like the Higgs boson in the SM.
- Why 'SM-like' Higgs boson?
- Because the SM is extremely success, the existence of one 'SM-like' Higgs boson is the simplest way to coincide the data.
- Other crazy scenarios are, of course, allowed!

- What is the 'SM-like' Higgs boson?
- A: We name the Higgs boson as 'SM-like' in case that one of the Higgs bosons in physics beyond the SM (for example MSSM) acts like the Higgs boson in the SM.
- Why 'SM-like' Higgs boson?
- Because the SM is extremely success, the existence of one 'SM-like' Higgs boson is the simplest way to coincide the data.
- Other crazy scenarios are, of course, allowed!

- What is the 'SM-like' Higgs boson?
- A: We name the Higgs boson as 'SM-like' in case that one of the Higgs bosons in physics beyond the SM (for example MSSM) acts like the Higgs boson in the SM.
- Why 'SM-like' Higgs boson?
- Because the SM is extremely success, the existence of one 'SM-like' Higgs boson is the simplest way to coincide the data.
- Other crazy scenarios are, of course, allowed!

- What is the 'SM-like' Higgs boson?
- A: We name the Higgs boson as 'SM-like' in case that one of the Higgs bosons in physics beyond the SM (for example MSSM) acts like the Higgs boson in the SM.
- Why 'SM-like' Higgs boson?
- Because the SM is extremely success, the existence of one 'SM-like' Higgs boson is the simplest way to coincide the data.
- Other crazy scenarios are, of course, allowed!

Invisible SM-like Higgs boson due to X(214) and analysis fault

- 'SM-like' Higgs boson
- HyperCP three events
- A new pseudoscalar X(214 MeV)?
- All about X(214)
- Consequences on Higgs physics

PRL 94, 021801 (2005)

PHYSICAL REVIEW LETTERS

week ending
 21 JANUARY 2005


Evidence for the Decay $\Sigma^+ \rightarrow p\mu^+\mu^-$

H. K. Park,⁸ R. A. Burnstein,⁵ A. Chakravorty,⁵ Y. C. Chen,¹ W. S. Choong,^{2,7} K. Clark,⁹ E. C. Dukes,¹⁰ C. Durandet,¹⁰
 J. Felix,⁴ Y. Fu,⁷ G. Gidal,⁷ H. R. Gustafson,⁸ T. Holmstrom,¹⁰ M. Huang,¹⁰ C. James,³ C. M. Jenkins,⁹ T. Jones,⁷
 D. M. Kaplan,⁵ L. M. Lederman,⁵ N. Leros,⁶ M. J. Longo,^{8,*} F. Lopez,⁸ L. C. Lu,¹⁰ W. Luebke,⁵ K. B. Luk,^{2,7}
 K. S. Nelson,¹⁰ J.-P. Perroud,⁶ D. Rajaram,⁵ H. A. Rubin,⁵ J. Volk,³ C. G. White,⁵ S. L. White,⁵ and P. Zyla⁷

(HyperCP Collaboration)

We report the first evidence for the decay $\Sigma^+ \rightarrow p\mu^+\mu^-$ from data taken by the HyperCP (E871) experiment at Fermilab. Based on three observed events, the branching ratio is $\mathcal{B}(\Sigma^+ \rightarrow p\mu^+\mu^-) = [8.6_{-5.4}^{+6.6}(\text{stat}) \pm 5.5(\text{syst})] \times 10^{-8}$. The narrow range of dimuon masses may indicate that the decay proceeds via a neutral intermediate state, $\Sigma^+ \rightarrow pP^0, P^0 \rightarrow \mu^+\mu^-$ with a P^0 mass of 214.3 ± 0.5 MeV/ c^2 and branching ratio $\mathcal{B}(\Sigma^+ \rightarrow pP^0, P^0 \rightarrow \mu^+\mu^-) = [3.1_{-1.9}^{+2.4}(\text{stat}) \pm 1.5(\text{syst})] \times 10^{-8}$.

HyperCP Spectrometer (built for hyperon CP-violation search)



Salient Features:

- High-rate detectors & DAQ (100k evts/s)
- Alternating "+" & "-" running (with reversed B fields) to minimize systematics
- Simple, low-bias triggers based on hodoscope coincidences
- **Muon-ID system:**
 - 3 layers 80-cm-thick steel
 - 3 layers x & y proportional tubes
 - hodoscopes for triggering
 - μ triggers: $2\mu\text{LR} + 1\mu\text{L} + 10, 1\mu\text{R} + 5$
- **No other particle ID**

Invisible SM-like Higgs boson due to X(214) and analysis fault

- 'SM-like' Higgs boson
- HyperCP three events
- A new pseudoscalar X(214 MeV)?
- All about X(214)
- Consequences on Higgs physics

- X. G. He, J. Tandean and G. Valencia, Phys. Rev. D **72**, 074003 (2005) [arXiv:hep-ph/0506067].
- X. G. He, J. Tandean and G. Valencia, Phys. Lett. B **631**, 100 (2005) [arXiv:hep-ph/0509041].
- N. G. Deshpande, G. Eilam and J. Jiang, Phys. Lett. B **632**, 212 (2006) [arXiv:hep-ph/0509081].

- C. Q. Geng and Y. K. Hsiao, Phys. Lett. B **632**, 215 (2006) [arXiv:hep-ph/0509175].
- D. S. Gorbunov and V. A. Rubakov, Phys. Rev. D **73**, 035002 (2006) [arXiv:hep-ph/0509147].
- S. V. Demidov and D. S. Gorbunov, JETP Lett. **84**, 479 (2007) [arXiv:hep-ph/0610066].
- X. G. He, J. Tandean and G. Valencia, Phys. Rev. D **74**, 115015 (2006) [arXiv:hep-ph/0610274].

- X. G. He, J. Tandean and G. Valencia, Phys. Rev. Lett. **98**, 081802 (2007) [arXiv:hep-ph/0610362].
- G. Hiller, Phys. Rev. D **70**, 034018 (2004) [arXiv:hep-ph/0404220].
- C. H. Chen and C. Q. Geng, Phys. Lett. B **645**, 189 (2007) [arXiv:hep-ph/0612142].
- G. Xiangdong, C. S. Li, Z. Li and H. Zhang, arXiv:0712.0257 [hep-ph].

X(214) is *not*

- Scalar
- Vector

However X(214) can be pseudoscalar or axial vector!

X(214) is *not*

- Scalar
- Vector

However X(214) can be pseudoscalar or axial vector!


Invisible SM-like Higgs boson due to X(214) and analysis fault

- 'SM-like' Higgs boson
- HyperCP three events
- A new pseudoscalar X(214 MeV)?
- **All about X(214)**
- Consequences on Higgs physics


- $m_X = 214.3 \pm 0.5 \text{ MeV}$
- Dominantly decays into $\mu^+\mu^-$, not photons
- $\Delta m \equiv m_X - 2m_\mu \approx 3 \text{ MeV}$
- **Likely neglected** for past experiments, LEP, Tevatron etc.
- Reasons...

Decay: $X \rightarrow \mu\mu$

If X is at rest...




If X is very fast...



*Note H to 2 photon signal is
heavily polluted by π^0 to 2
photons at LHC for this reason!*

If X is very fast and $m_X \sim 2 m_{\mu}$...






FIG. 1: Distribution of ΔR between $\mu^+\mu^-$ at Tevatron for signal $gg \rightarrow h \rightarrow aa \rightarrow 4\mu$ with "Tevatron basic cuts" of Eq. (4). In all figures of this paper, $m_h = 120$ GeV and $m_a = 0.215$ GeV, except indicated otherwise. The solid (dashed) lines in all figures represent signal (background).

- $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ approaches 0 due to the tiny $\Delta m \equiv m_X - 2m_\mu$
- At ATLAS, $\Delta R > 0.01$ is applied in order to suppress fake muon and separate different tracks!
- Similar at other detectors!
- X will be missing due to analysis method!
- Fortunately X(214) can be identified at modern detectors, like CMS at LHC, due to the strong magnetic field.

- $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ approaches 0 due to the tiny $\Delta m \equiv m_X - 2m_\mu$
- At ATLAS, $\Delta R > 0.01$ is applied in order to suppress fake muon and separate different tracks!
- Similar at other detectors!
- X will be missing due to analysis method!
- Fortunately X(214) can be identified at modern detectors, like CMS at LHC, due to the strong magnetic field.

- $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ approaches 0 due to the tiny $\Delta m \equiv m_X - 2m_\mu$
- At ATLAS, $\Delta R > 0.01$ is applied in order to suppress fake muon and separate different tracks!
- Similar at other detectors!
- X will be missing due to analysis method!
- Fortunately X(214) can be identified at modern detectors, like CMS at LHC, due to the strong magnetic field.

- $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ approaches 0 due to the tiny $\Delta m \equiv m_X - 2m_\mu$
- At ATLAS, $\Delta R > 0.01$ is applied in order to suppress fake muon and separate different tracks!
- Similar at other detectors!
- X will be missing due to analysis method!
- Fortunately X(214) can be identified at modern detectors, like CMS at LHC, due to the strong magnetic field.

- $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ approaches 0 due to the tiny $\Delta m \equiv m_X - 2m_\mu$
- At ATLAS, $\Delta R > 0.01$ is applied in order to suppress fake muon and separate different tracks!
- Similar at other detectors!
- X will be missing due to analysis method!
- Fortunately X(214) can be identified at modern detectors, like CMS at LHC, due to the strong magnetic field.



Event ($X \rightarrow \mu^+ \mu^-$) view at CMS detector by Z.C. Yang of Peking University!

Introduction
Invisible SM-like Higgs boson due to X(214) and analysis fault
Correlation between masses of SM-like Higgs boson and (singlet) scalar
Higgs boson as the looking glass in mirror model
Conclusion and discussion

'SM-like' Higgs boson
HyperCP three events
A new pseudoscalar X(214 MeV)?
All about X(214)
Consequences on Higgs physics



Invisible SM-like Higgs boson due to X(214) and analysis fault

- 'SM-like' Higgs boson
- HyperCP three events
- A new pseudoscalar X(214 MeV)?
- All about X(214)
- Consequences on Higgs physics

SM-like Higgs boson may decay dominantly into a pair of X(214)




FIG. 1: $BR(h \rightarrow aa)$ as a function of m_a in the NMSSM with $\tan\beta = 30 \sim 60$, $|\mu| = 100 \sim 300$ GeV, $A_t = 1500$ GeV, $M_{SUSY} = 1000$ GeV and $M_{1,2,3} = 100, 300, 500$ GeV. The $+$ (\times) points indicate $m_h < 114$ (> 114) GeV.

- SM-like Higgs boson h will be missing because X is missing, provided that h decays dominantly into X pair.
- Direct limit 114 GeV should be altered, likely shift to lower than 100 GeV, as indicated by indirect limit!
- LEP/Tevatron data need to be re-analyzed!
- LHC may adjust its searching strategies for Higgs boson!

- SM-like Higgs boson h will be missing because X is missing, provided that h decays dominantly into X pair.
- Direct limit 114 GeV should be altered, likely shift to lower than 100 GeV, as indicated by indirect limit!
- LEP/Tevatron data need to be re-analyzed!
- LHC may adjust its searching strategies for Higgs boson!

- SM-like Higgs boson h will be missing because X is missing, provided that h decays dominantly into X pair.
- Direct limit 114 GeV should be altered, likely shift to lower than 100 GeV, as indicated by indirect limit!
- LEP/Tevatron data need to be re-analyzed!
- LHC may adjust its searching strategies for Higgs boson!

- SM-like Higgs boson h will be missing because X is missing, provided that h decays dominantly into X pair.
- Direct limit 114 GeV should be altered, likely shift to lower than 100 GeV, as indicated by indirect limit!
- LEP/Tevatron data need to be re-analyzed!
- LHC may adjust its searching strategies for Higgs boson!

Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter

- What determines the dark matter mass?
- Singlet scalar as the dark matter: the model
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- Other supports
- Pause

Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter

- What determines the dark matter mass?
- Singlet scalar as the dark matter: the model
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- Other supports
- Pause

- No one knows!
- If dark matter is closely related with weak physics, *why not* the dark matter mass originates from electro-weak symmetry breaking?

- No one knows!
- If dark matter is closely related with weak physics, *why not* the dark matter mass originates from electro-weak symmetry breaking?

Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter

- What determines the dark matter mass?
- **Singlet scalar as the dark matter: the model**
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- Other supports
- Pause

- We require the theory to be renormalizable, thus naturally take the singlet scalar as the dark matter
- Lagrangian is written as

$$L = L_{SM} + \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{\lambda_S}{4} S^4 - \lambda S^2 (\Phi^\dagger \Phi)$$

- L_{SM} is the Lagrangian of the SM and Φ is the weak doublet Higgs field. L is obviously invariant under discrete transformation $S \rightarrow -S$, which ensures S the good candidate of cold dark matter.

- We require the theory to be renormalizable, thus naturally take the singlet scalar as the dark matter
- Lagrangian is written as

$$L = L_{SM} + \frac{1}{2}\partial_\mu S \partial^\mu S - \frac{\lambda_S}{4}S^4 - \lambda S^2(\Phi^\dagger\Phi)$$

- L_{SM} is the Lagrangian of the SM and Φ is the weak doublet Higgs field. L is obviously invariant under discrete transformation $S \rightarrow -S$, which ensures S the good candidate of cold dark matter.

- We require the theory to be renormalizable, thus naturally take the singlet scalar as the dark matter
- Lagrangian is written as

$$L = L_{SM} + \frac{1}{2}\partial_\mu S \partial^\mu S - \frac{\lambda_S}{4}S^4 - \lambda S^2(\Phi^\dagger\Phi)$$

- L_{SM} is the Lagrangian of the SM and Φ is the weak doublet Higgs field. L is obviously invariant under discrete transformation $S \rightarrow -S$, which ensures S the good candidate of cold dark matter.

- $m_0^2 S^2$ is simply omitted, or negligible compared with the contribution arising from electro-weak symmetry breaking!
- After electro-weak symmetry breaking $\langle \Phi \rangle = v = 246$ GeV, the Higgs boson, as in the standard model,

$$m_h^2 = \lambda_h v^2$$

with λ_h the coefficient of $(\Phi^\dagger \Phi)^2$ and

$$m_S^2 = \lambda v^2.$$

- $m_0^2 S^2$ is simply omitted, or negligible compared with the contribution arising from electro-weak symmetry breaking!
- After electro-weak symmetry breaking $\langle \Phi \rangle = v = 246$ GeV, the Higgs boson, as in the standard model,

$$m_h^2 = \lambda_h v^2$$

with λ_h the coefficient of $(\Phi^+ \Phi)^2$ and


$$m_S^2 = \lambda v^2.$$

- It is obvious that coupling λ is determined by m_S and in this model λ is the only extra free parameter relevant to our discussion, besides those in the SM.
- How to determine λ , namely dark matter mass m_S ?

- It is obvious that coupling λ is determined by m_S and in this model λ is the only extra free parameter relevant to our discussion, besides those in the SM.
- How to determine λ , namely dark matter mass m_S ?

Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter

- What determines the dark matter mass?
- Singlet scalar as the dark matter: the model
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- Other supports
- Pause



Schematic Feynman diagram for $SS \rightarrow \text{SM}$ particles. Here f and V represent SM fermions and weak gauge bosons respectively.

- The current relic density of S can be written as

(C. P. Burgess, M. Pospelov and T. ter Veldhuis, Nucl. Phys. B **619**, 709 (2001))

$$\Omega_S h^2 = \frac{(1.07 \times 10^9) x_f}{\sqrt{g_*} M_{pl} [\text{in GeV}] \langle \sigma v_{rel} \rangle},$$

where g_* counts the degrees of freedom in equilibrium at annihilation, x_f is the inverse freeze-out temperature in units of m_S , which can be obtained by solving the equation

$$x_f \simeq \ln \left[\frac{0.038 M_{pl} m_S \langle \sigma v_{rel} \rangle}{\sqrt{g_*} x_f} \right].$$

- Here v_{rel} is the relative velocity of the two incoming dark matter particles, M_{pl} is the Planck mass and $\langle \dots \rangle$ denotes the relevant thermal average.

- σv_{rel} is

$$\begin{aligned}\sigma_{ann} v_{rel} &= \frac{8\lambda^2 v^2}{(4m_S^2 - m_h^2)^2 + m_h^2 \Gamma_h^2} F_X \\ &= \frac{8m_S^4}{v^2 [(4m_S^2 - m_h^2)^2 + m_h^2 \Gamma_h^2]} F_X(1)\end{aligned}$$

- Here v_{rel} is the relative velocity of the two incoming dark matter particles, M_{pl} is the Planck mass and $\langle \dots \rangle$ denotes the relevant thermal average.
- σv_{rel} is

$$\begin{aligned}\sigma_{ann} v_{rel} &= \frac{8\lambda^2 v^2}{(4m_S^2 - m_h^2)^2 + m_h^2 \Gamma_h^2} F_X \\ &= \frac{8m_S^4}{v^2 [(4m_S^2 - m_h^2)^2 + m_h^2 \Gamma_h^2]} F_X(1)\end{aligned}$$

- Here

$$F_X = \lim_{m_{\tilde{h}} \rightarrow 2m_S} \left(\frac{\Gamma_{\tilde{h} \rightarrow X}}{m_{\tilde{h}}} \right). \quad (2)$$

- Γ_h is the Higgs total decay width and $\Gamma_{\tilde{h} \rightarrow X}$ denotes the partial decay width for the virtual \tilde{h} decay into X , $\tilde{h} \rightarrow X$, in the limit $m_{\tilde{h}} \rightarrow 2m_S$. Here X represents SM particles.
- Relic density (within 3σ uncertainty)

$$0.093 < \Omega_{dm} h^2 < 0.129$$

where $h \approx 0.71$ is the normalized Hubble expansion rate.

- Here

$$F_X = \lim_{m_{\tilde{h}} \rightarrow 2m_S} \left(\frac{\Gamma_{\tilde{h} \rightarrow X}}{m_{\tilde{h}}} \right). \quad (2)$$

- Γ_h is the Higgs total decay width and $\Gamma_{\tilde{h} \rightarrow X}$ denotes the partial decay width for the virtual \tilde{h} decay into X , $\tilde{h} \rightarrow X$, in the limit $m_{\tilde{h}} \rightarrow 2m_S$. Here X represents SM particles.
- Relic density (within 3σ uncertainty)

$$0.093 < \Omega_{dm} h^2 < 0.129$$

where $h \approx 0.71$ is the normalized Hubble expansion rate.


- Here

$$F_X = \lim_{m_{\tilde{h}} \rightarrow 2m_S} \left(\frac{\Gamma_{\tilde{h} \rightarrow X}}{m_{\tilde{h}}} \right). \quad (2)$$

- Γ_h is the Higgs total decay width and $\Gamma_{\tilde{h} \rightarrow X}$ denotes the partial decay width for the virtual \tilde{h} decay into X , $\tilde{h} \rightarrow X$, in the limit $m_{\tilde{h}} \rightarrow 2m_S$. Here X represents SM particles.
- Relic density (within 3σ uncertainty)

$$0.093 < \Omega_{dm} h^2 < 0.129$$

where $h \approx 0.71$ is the normalized Hubble expansion rate.




m_S [GeV]	m_h upper limit [GeV]	m_h lower limit [GeV]
50	122	119
55	134	131
60	148	144
65	162	158
70	180	174
75	204	197
80	275	261

Table: Upper and lower limits on Higgs boson for several m_S in order to obtain the correct relic abundance.

Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter


- What determines the dark matter mass?
- Singlet scalar as the dark matter: the model
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- Other supports
- Pause




Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter

- What determines the dark matter mass?
- Singlet scalar as the dark matter: the model
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- **Other supports**
- Pause

Background + signal describe EGRET data! by W. de Boer et al.



Blue: background uncertainty



Blue: WIMP mass uncertainty

Summary from W. De Boer

10σ EGRET excess shows all key features from DM annihilation:

Excess has same shape in all sky directions: everywhere it is perfectly (only?) explainable with superposition of background AND *mono-energetic quarks of 50-100 GeV*
Results consistent with minimal SUPERSYMMETRY

Excess follows expectations from galaxy formation:
cored $1/r^2$ profile with substructure,
visible matter/DM ≈ 0.02

Excess is TRACER OF DM, since it can explain
peculiar shape of rotation curve

Significance
 $>10\sigma$ with
 >1400 indep.
data points

Results model independent, since only KNOWN spectral shapes of signal and background used, NO model dependent calculations of absolute fluxes.

Conventional models CANNOT explain above points SIMULTANEOUSLY,
especially spectrum of gamma rays in all directions, shape of rotation curve,
stability of ring of stars at 14 kpc and ring of H_2 at 4 kpc,...

Correlation between masses of SM-like Higgs boson and (singlet) scalar dark matter

- What determines the dark matter mass?
- Singlet scalar as the dark matter: the model
- Correlation between SM-like Higgs boson and dark matter
- Invisible SM-like Higgs boson decay
- Other supports
- **Pause**

- Seems everything is perfect: Higgs mass bounds can predict scalar CDM value and EGRET prefers the same mass region.
- The Higgs boson is light and may decay dominantly into (~ 60 GeV)DM.
- Detecting invisible Higgs boson at LHC and ILC!

- Seems everything is perfect: Higgs mass bounds can predict scalar CDM value and EGRET prefers the same mass region.
- The Higgs boson is light and may decay dominantly into (~ 60 GeV)DM.
- Detecting invisible Higgs boson at LHC and ILC!

- Seems everything is perfect: Higgs mass bounds can predict scalar CDM value and EGRET prefers the same mass region.
- The Higgs boson is light and may decay dominantly into (~ 60 GeV)DM.
- Detecting invisible Higgs boson at LHC and ILC!

Higgs boson as the looking glass in mirror model

- Broken parity
- Parity restoration: two approaches
- Communication with mirror world: three ways
- Minimal mirror model
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal

Higgs boson as the looking glass in mirror model

- Broken parity
- Parity restoration: two approaches
- Communication with mirror world: three ways
- Minimal mirror model
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal

- For the pioneering work on parity-broken, T.D. Lee and C.N. Yang got Nobel Prize in 1957!
- Basis for the SM construction: left-handed fermions feel $SU(2)$ gauge interaction, while right-handed ones do not.
- Why is parity not conserved solely for weak interaction?

- For the pioneering work on parity-broken, T.D. Lee and C.N. Yang got Nobel Prize in 1957!
- Basis for the SM construction: left-handed fermions feel $SU(2)$ gauge interaction, while right-handed ones do not.
- Why is parity not conserved solely for weak interaction?

- For the pioneering work on parity-broken, T.D. Lee and C.N. Yang got Nobel Prize in 1957!
- Basis for the SM construction: left-handed fermions feel $SU(2)$ gauge interaction, while right-handed ones do not.
- Why is parity not conserved solely for weak interaction?

Higgs boson as the looking glass in mirror model

- Broken parity
- **Parity restoration: two approaches**
- Communication with mirror world: three ways
- Minimal mirror model
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal

L-R models (For a review to see, R. Mohapatra, 'Unification and Supersymmetry')

- Parity is restored at energy scale higher than weak interaction (represented by $m_W \sim 100$ GeV)
- Predict generically W_R and Z_R which are non-singlet fields for right-handed usual SM fermions.
- Gauge structure other than the SM one has not been experimentally established. On the contrary, W_R has been pushed up to 1.6 TeV or higher in minimal L-R model.
- LHC can detect $O(5 \text{ TeV})$ Z_R .

L-R models (For a review to see, R. Mohapatra, 'Unification and Supersymmetry')

- Parity is restored at energy scale higher than weak interaction (represented by $m_W \sim 100$ GeV)
- Predict generically W_R and Z_R which are non-singlet fields for right-handed usual SM fermions.
- Gauge structure other than the SM one has not been experimentally established. On the contrary, W_R has been pushed up to 1.6 TeV or higher in minimal L-R model.
- LHC can detect $O(5 \text{ TeV}) Z_R$.

L-R models (For a review to see, R. Mohapatra, 'Unification and Supersymmetry')

- Parity is restored at energy scale higher than weak interaction (represented by $m_W \sim 100$ GeV)
- Predict generically W_R and Z_R which are non-singlet fields for right-handed usual SM fermions.
- Gauge structure other than the SM one has not been experimentally established. On the contrary, W_R has been pushed up to 1.6 TeV or higher in minimal L-R model.
- LHC can detect $O(5 \text{ TeV})$ Z_R ;

L-R models (For a review to see, R. Mohapatra, 'Unification and Supersymmetry')

- Parity is restored at energy scale higher than weak interaction (represented by $m_W \sim 100$ GeV)
- Predict generically W_R and Z_R which are non-singlet fields for right-handed usual SM fermions.
- Gauge structure other than the SM one has not been experimentally established. On the contrary, W_R has been pushed up to 1.6 TeV or higher in minimal L-R model.
- LHC can detect $O(5 \text{ TeV}) Z_R$.

Mirror models (For a review to see, L.B. Okun, hep-ph/0606202)

- Parity is restored in a general sense, as originated in Lee-Yang' s 1956 paper
- Current experiments disfavor the case that mirror particles are non-singlet under SM gauge group.
- Singlet mirror particles = dark matter particles?

Mirror models (For a review to see, L.B. Okun, hep-ph/0606202)

- Parity is restored in a general sense, as originated in Lee-Yang' s 1956 paper
- Current experiments disfavor the case that mirror particles are non-singlet under SM gauge group.
- Singlet mirror particles = dark matter particles?


Mirror models (For a review to see, L.B. Okun, hep-ph/0606202)

- Parity is restored in a general sense, as originated in Lee-Yang' s 1956 paper
- Current experiments disfavor the case that mirror particles are non-singlet under SM gauge group.
- Singlet mirror particles = dark matter particles?


Okun said: “Mirsy (mirror symmetry) cannot compete with SUSY in the depth of its concept and mathematics. But I believe it can compete in the breadth and diversity of its phenomenological predictions. Certainly, mirror matter is richer than the dark matter of SUSY”

Higgs boson as the looking glass in mirror model


- Broken parity
- Parity restoration: two approaches
- **Communication with mirror world: three ways**
- Minimal mirror model
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal



General structure: via Neutrino




General structure: via $U(1)$ s kinetic mixing



Higgs boson as the looking glass in mirror model

- Broken parity
- Parity restoration: two approaches
- Communication with mirror world: three ways
- **Minimal mirror model**
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal

R. Foot, H. Lew and R. R. Volkas, PLB(1991)



- The minimal gauge group of the new mirror model is $G_{SM} \otimes G' = SU(3) \otimes SU(2) \otimes U(1) \otimes SU(3)' \otimes SU(2)' \otimes U(1)'$.
- The gauge quantum numbers under $G_{SM} \otimes G'$ for the usual and mirror fermion fields are

- The minimal gauge group of the new mirror model is $G_{SM} \otimes G' = SU(3) \otimes SU(2) \otimes U(1) \otimes SU(3)' \otimes SU(2)' \otimes U(1)'$.
- The gauge quantum numbers under $G_{SM} \otimes G'$ for the usual and mirror fermion fields are

$$\begin{aligned} L_L^i &\sim (1, 2, -1)(1, 1, 0) \quad , \quad (L_R')^i \sim (1, 1, 0)(1, 2, -1) \\ e_R^i &\sim (1, 1, -2)(1, 1, 0) \quad , \quad (e_L')^i \sim (1, 1, 0)(1, 1, -2) \\ Q_L^i &\sim (3, 2, \frac{1}{3})(1, 1, 0) \quad , \quad (q_R')^i \sim (1, 1, 0)(3, 2, \frac{1}{3}) \\ u_R^i &\sim (3, 1, \frac{4}{3})(1, 1, 0) \quad , \quad (u_L')^i \sim (1, 1, 0)(3, 1, \frac{4}{3}) \\ d_R^i &\sim (3, 1, -\frac{2}{3})(1, 1, 0) \quad , \quad (d_L')^i \sim (1, 1, 0)(3, 1, -\frac{2}{3}) \end{aligned}$$

with i the family index.

- The Z_2 parity symmetry that we define now is

$$\begin{aligned}\vec{r} &\leftrightarrow -\vec{r}, t \leftrightarrow t, & G^\mu &\leftrightarrow G'_\mu \\ W^\mu &\leftrightarrow W'_\mu, B^\mu \leftrightarrow B'_\mu \\ L_L &\leftrightarrow L'_R, e_R \leftrightarrow e'_L, & Q_L &\leftrightarrow Q'_R, \\ u_R &\leftrightarrow u'_L, d_R \leftrightarrow d'_L.\end{aligned}$$

- One of the advantages of this model is that there are natural candidates of non-baryonic dark matter in addition to restore parity.
- Focus on Higgs sector!

- The Z_2 parity symmetry that we define now is

$$\begin{aligned}\vec{r} &\leftrightarrow -\vec{r}, t \leftrightarrow t, & G^\mu &\leftrightarrow G'_\mu \\ W^\mu &\leftrightarrow W'_\mu, B^\mu \leftrightarrow B'_\mu \\ L_L &\leftrightarrow L'_R, e_R \leftrightarrow e'_L, & Q_L &\leftrightarrow Q'_R, \\ u_R &\leftrightarrow u'_L, d_R \leftrightarrow d'_L.\end{aligned}$$

- One of the advantages of this model is that there are natural candidates of non-baryonic dark matter in addition to restore parity.
- Focus on Higgs sector!

- The Z_2 parity symmetry that we define now is

$$\begin{aligned}\vec{r} &\leftrightarrow -\vec{r}, t \leftrightarrow t, & G^\mu &\leftrightarrow G'_\mu \\ W^\mu &\leftrightarrow W'_\mu, B^\mu \leftrightarrow B'_\mu \\ L_L &\leftrightarrow L'_R, e_R \leftrightarrow e'_L, & Q_L &\leftrightarrow Q'_R, \\ u_R &\leftrightarrow u'_L, d_R \leftrightarrow d'_L.\end{aligned}$$

- One of the advantages of this model is that there are natural candidates of non-baryonic dark matter in addition to restore parity.
- Focus on Higgs sector!

Higgs boson as the looking glass in mirror model

- Broken parity
- Parity restoration: two approaches
- Communication with mirror world: three ways
- Minimal mirror model
- **Two kinds of vacua**
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal

- One assumes that the Higgs potential is invariant under the discrete symmetry $\phi_1 \rightarrow \phi_2$ to keep the parity in a broader sense.
- The Higgs potential is very simply given by

$$\begin{aligned} V(\phi_1, \phi_2) = & -\mu^2 \left(\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 \right) \\ & + \lambda \left(\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 \right)^2 \\ & + \eta \phi_1^\dagger \phi_1 \phi_2^\dagger \phi_2. \end{aligned} \quad (3)$$

- One assumes that the Higgs potential is invariant under the discrete symmetry $\phi_1 \rightarrow \phi_2$ to keep the parity in a broader sense.
- The Higgs potential is very simply given by

$$\begin{aligned} V(\phi_1, \phi_2) = & -\mu^2 \left(\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 \right) \\ & + \lambda \left(\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 \right)^2 \\ & + \eta \phi_1^\dagger \phi_1 \phi_2^\dagger \phi_2. \end{aligned} \quad (3)$$

After electro-weak symmetry breaking, the Higgs fields can be written as

$$\phi_i = \left(\begin{array}{c} \varphi_i \\ \frac{1}{\sqrt{2}}(v_i + H_i + \chi_i) \end{array} \right), \quad (4)$$

where $\varphi_i^\dagger, \chi_i$ are Goldstone bosons, which will be absorbed by corresponding gauge fields.

The vacuum may not be invariant under Z_2 transformation although the Higgs potential is invariant under this discrete transformation. In fact there are two ways of spontaneous symmetry breaking, depending on the choice of the sign of η

(R. Foot, H. Lew, R. R. Volkas, JHEP **032**, 0007 (2000)).

$\eta < 0$: symmetric vacua:

- Vacuum is invariant under transformation

$$\phi_1 \leftrightarrow \phi_2.$$

$$v^2 = v_1^2 = v_2^2 = \frac{2\mu^2}{4\lambda + \eta}. \quad (5)$$

- Define Higgs boson mass eigenstates as H, h (assume H is heavier than h),

$$H_1 = \frac{1}{\sqrt{2}}(H + h) \quad (6)$$

$$H_2 = \frac{1}{\sqrt{2}}(H - h). \quad (7)$$

$\eta < 0$: symmetric vacua:

- Vacuum is invariant under transformation

$$\phi_1 \leftrightarrow \phi_2.$$

$$v^2 = v_1^2 = v_2^2 = \frac{2\mu^2}{4\lambda + \eta}. \quad (5)$$

- Define Higgs boson mass eigenstates as H, h
(assume H is heavier than h),

$$H_1 = \frac{1}{\sqrt{2}}(H + h) \quad (6)$$

$$H_2 = \frac{1}{\sqrt{2}}(H - h). \quad (7)$$

- The Higgs boson mass can be expressed as

$$m_H^2 = (4\lambda + \eta)v^2 \quad (8)$$

$$m_h^2 = -\eta v^2. \quad (9)$$

$\eta > 0$: non-symmetric vacuum:

- Requiring the minimum of Higgs potential is stable, then

$$v_1^2 = \frac{\mu^2}{\lambda}, v_2^2 = 0. \quad (10)$$

- The Higgs boson masses are

$$m_h^2 = \frac{\mu^2}{2}, m_H^2 = \frac{\eta v_1^2}{8} \quad (11)$$

$\eta > 0$: non-symmetric vacuum:

- Requiring the minimum of Higgs potential is stable, then

$$v_1^2 = \frac{\mu^2}{\lambda}, v_2^2 = 0. \quad (10)$$

- The Higgs boson masses are

$$m_h^2 = \frac{\mu^2}{2}, m_H^2 = \frac{\eta v_1^2}{8} \quad (11)$$

- It seems that all the mirror particles must be massless. However mirror particle can obtain tiny mass through mirror QCD condensation, but we don't discuss this case further.

Higgs boson as the looking glass in mirror model

- Broken parity
- Parity restoration: two approaches
- Communication with mirror world: three ways
- Minimal mirror model
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal




FIG. 1: The ellipses indicate the regions in the S, T plane which are allowed by EWPO at 1σ (68%) and 2σ (95%) level respectively [20]. Three curves represent three different m_h at 115, 150 and 200 GeV, and m_H increases from 100 ~ 1000 GeV for each curve.

Lighter Higgs boson (h) decay

$$\begin{aligned} & Br(h \rightarrow b\bar{b}) \\ &= \frac{\Gamma(h \rightarrow b\bar{b})}{\Gamma(h \rightarrow SM) + \Gamma(h \rightarrow Mirror)} \\ &= \frac{1}{2} Br(h_{SM} \rightarrow b\bar{b}), \end{aligned} \tag{12}$$

Heavier Higgs boson (H) decay

$$\begin{aligned}
 & Br(H \rightarrow hh) [Br(H \rightarrow b\bar{b})] \\
 &= \frac{\Gamma(H \rightarrow hh) [\Gamma(H \rightarrow b\bar{b})]}{\Gamma(H \rightarrow SM) + \Gamma(H \rightarrow Mirror) + \Gamma(H \rightarrow hh)} \\
 &= \frac{\Gamma(H \rightarrow hh) [\Gamma(H \rightarrow b\bar{b})]}{2\Gamma(H \rightarrow SM) + \Gamma(H \rightarrow hh)}. \tag{13}
 \end{aligned}$$




FIG. 2: Branching ratios of H as a function of m_H , where $m_h = 115\text{GeV}$.

Higgs bosons production

- For the case $m_H < 2m_h$, each Higgs boson acts like the SM one except with only half production rate in MM model. Moreover each Higgs boson decays into SM matter with branching ratio half of SM case in MM model.
- For the case $m_H > 2m_h$, new decay channel opens up. Thus provides unique signal to investigate such model. (We focus on this scenario!)

Higgs bosons production

- For the case $m_H < 2m_h$, each Higgs boson acts like the SM one except with only half production rate in MM model. Moreover each Higgs boson decays into SM matter with branching ratio half of SM case in MM model.
- For the case $m_H > 2m_h$, new decay channel opens up. Thus provides unique signal to investigate such model. (We focus on this scenario!)

Higgs boson as the looking glass in mirror model

- Broken parity
- Parity restoration: two approaches
- Communication with mirror world: three ways
- Minimal mirror model
- Two kinds of vacua
- Higgs phenomenology for symmetric vacuum
- Detecting $H \rightarrow hh$ signal




FIG. 4: Cross sections [in pb] of $gg \rightarrow H \rightarrow hh$ as a function of m_H at the LHC, where $m_h = 100, 115, 130, 150$ and 180 GeV from top to bottom .

Detail simulation for

$$p p \rightarrow g g \rightarrow H \rightarrow h(\rightarrow b^+b^-) + h_{inv}$$

- The most important irreducible background arises from $Zb\bar{b}$ production, where Z decays into neutrinos.
- Moreover QCD multi-jet production, such as $pp \rightarrow Z(\rightarrow \nu\bar{\nu})jj$, are also the sources of the large backgrounds.
- In our analysis we require two b-tagged jets in order to suppress these backgrounds.
- Other backgrounds can arise from ZZ , WZ , $Wb\bar{b}$, single top and $t\bar{t}$ production.

Detail simulation for

$$p p \rightarrow g g \rightarrow H \rightarrow h(\rightarrow b^+b^-) + h_{inv}$$

- The most important irreducible background arises from $Zb\bar{b}$ production, where Z decays into neutrinos.
- Moreover QCD multi-jet production, such as $pp \rightarrow Z(\rightarrow \nu\bar{\nu})jj$, are also the sources of the large backgrounds.
- In our analysis we require two b-tagged jets in order to suppress these backgrounds.
- Other backgrounds can arise from ZZ , WZ , $Wb\bar{b}$, single top and $t\bar{t}$ production.

Detail simulation for

$$p p \rightarrow g g \rightarrow H \rightarrow h(\rightarrow b^+b^-) + h_{inv}$$

- The most important irreducible background arises from $Zb\bar{b}$ production, where Z decays into neutrinos.
- Moreover QCD multi-jet production, such as $pp \rightarrow Z(\rightarrow \nu\bar{\nu})jj$, are also the sources of the large backgrounds.
- In our analysis we require two b-tagged jets in order to suppress these backgrounds.
- Other backgrounds can arise from ZZ , WZ , $Wb\bar{b}$, single top and $t\bar{t}$ production

Detail simulation for

$$p p \rightarrow g g \rightarrow H \rightarrow h(\rightarrow b^+b^-) + h_{inv}$$

- The most important irreducible background arises from $Zb\bar{b}$ production, where Z decays into neutrinos.
- Moreover QCD multi-jet production, such as $pp \rightarrow Z(\rightarrow \nu\bar{\nu})jj$, are also the sources of the large backgrounds.
- In our analysis we require two b-tagged jets in order to suppress these backgrounds.
- Other backgrounds can arise from ZZ , WZ , $Wb\bar{b}$, single top and $t\bar{t}$ production

Basic cuts:

$$P_T(j_1), P_T(j_2) > 20\text{GeV}, 15\text{GeV} \quad (14)$$

$$|\eta_j| < 2 \quad (15)$$

$$\Delta R(jj) > 0.4 \quad (16)$$

$$m_{jj} > 10\text{GeV}, \quad (17)$$

To suppress the backgrounds from $Wb\bar{b}$, WZ , single top with $t \rightarrow bW$ and $t\bar{t} \rightarrow bW\bar{b}W$. For these backgrounds, the final state charge leptons or jets from W escape from the detection. We suppress these contributions by vetoing events from W decay with follow cuts

$$P_T(j) > 15\text{GeV}, |\eta(j)| < 2.0 \quad (18)$$

$$P_T(l^\pm) > 10\text{GeV}, |\eta(l^\pm)| < 2.5 \quad (19)$$


The numerical results after imposing cuts Eqs. 14-19

Channel	$Zb\bar{b}$	$Zb\bar{c}$	Zbj	$Zc\bar{c}$	Zcj	Zjj
$\sigma(pb)$	3.250	0.011	0.107	0.001	0.027	0.063

Channel	ZZ	$W^-b\bar{b}$	W^-Z	$t\bar{b}$	$t\bar{t}$
$\sigma(pb)$	0.072	0.417	0.032	0.017	0.346

Table: The cross sections (in pb) of backgrounds for $b\bar{b} + \cancel{P}_T$ after basic kinematical cuts Eqs. 14-19 and tagging efficiencies where $j = u, \bar{u}, d, \bar{d}, s, \bar{s}, g$.

The signals and backgrounds for $b\bar{b} + \cancel{P}_T$ as a function of \cancel{P}_T




In order to suppress the backgrounds, we impose the further cuts as following

$$|m_{jj} - m_h| < 15\text{GeV} \quad (20)$$

$$40\text{GeV} < \cancel{P}_T < 80\text{GeV}. \quad (21)$$

The signals and backgrounds as the function of $|\eta_{j_1} - \eta_{j_2}|$



We require

$$|\eta_{j_1} - \eta_{j_2}| < 1.5, \quad (22)$$

and this cut would improve significance of the signals by a factor of 1.2.

In order to suppress the largest $Zb\bar{b}$ background, we can utilize the precise measurement of $Z(\rightarrow \mu^+\mu^-)b\bar{b}$.

$$\sigma_{bkg}^{Zb\bar{b},imp} = \sigma_{bkg}^{Zb\bar{b}} - R \times \sigma_{b\bar{b}\mu^+\mu^-}. \quad (23)$$

In Eq. 23, $\sigma_{b\bar{b}\mu^+\mu^-}$ is the cross section for $Z(\rightarrow \mu^+\mu^-)b\bar{b}$ production which adopts the same kinematical cuts for $Z(\rightarrow \nu\bar{\nu})b\bar{b}$.

R is a ratio which is defined as

$$R = \frac{\sum_i Br(Z \rightarrow \nu_i \bar{\nu}_i)}{Br(Z \rightarrow \mu^+ \mu^-)}, \quad (24)$$

and in our case $R = 5.94$. Note that $\sigma_{bkg}^{Zb\bar{b},imp} \approx 0$ if we can measure all final states $b\bar{b}\mu^+\mu^-$ in any kinematical region.

Cuts	$s(fb)$	$b(fb)$	S/B	$S/\sqrt{B_1}$	$S/\sqrt{B_2}$
basic cuts	26.6	4948	0.0054	1.19	2.07
$ m_{jj} - m_h < 30\text{GeV}$	26.6	1133	0.023	2.50	4.32
$ m_{jj} - m_h < 15\text{GeV}$	26.6	492	0.054	3.79	6.56
$20 < \cancel{P}_T < 120\text{GeV}$	25.0	401	0.062	3.94	6.83
$40 < \cancel{P}_T < 80\text{GeV}$	19.4	202	0.096	4.33	7.49
$ \eta_{j_1} - \eta_{j_2} < 1.5$	15.2	95	0.16	4.93	8.54
improved backg	15.2	18	0.83	11.4	19.8

Table: The significance $S/\sqrt{B_1}$ is for the luminosity of $10fb^{-1}$ and $S/\sqrt{B_2}$ is for the luminosity of $30fb^{-1}$. Here $m_H = 260$ GeV and $m_h = 115$ GeV.

	$m_h = 100\text{GeV}$	$m_h = 115\text{GeV}$	$m_h = 130\text{GeV}$
$m_H = 250 \text{ GeV}$	8.2(40,80)	8.3(10,60)	—
$m_H = 300 \text{ GeV}$	9.0(80,130)	9.6(60,110)	17.5(40,80)
$m_H = 350 \text{ GeV}$	5.5(100,150)	6.6(90,140)	11.6(80,120)

Table: The integrated luminosity [in fb^{-1}], which is required to observe $H \rightarrow hh \rightarrow b\bar{b} + \cancel{P}_T$ with 5σ significance at the LHC. The numbers in bracket are mass window of \cancel{P}_T . Note the Eq. 23 is not applied.

Detail simulation for $gg \rightarrow H \rightarrow hh \rightarrow 4b$.




FIG. 7: The distributions of the signals and backgrounds for $b\bar{b}b\bar{b}$ as a function of invariant mass for $4b$ after applying cuts (see text) and tagging efficiencies.

Conclusion and discussion

- Provided that there are physics beyond the SM, we find them *likely* via the investigation on Higgs boson(s) and/or dark matter.
- The importance of combination of cosmology and particle physics.
- The coming LHC may/should open the new chapter of particle physics!

- Provided that there are physics beyond the SM, we find them *likely* via the investigation on Higgs boson(s) and/or dark matter.
- The importance of combination of cosmology and particle physics.
- The coming LHC may/should open the new chapter of particle physics!

- Provided that there are physics beyond the SM, we find them *likely* via the investigation on Higgs boson(s) and/or dark matter.
- The importance of combination of cosmology and particle physics.
- The coming LHC may/should open the new chapter of particle physics!

Thanks for your attention!