

Neutrinoastrophysics with IceCube

Sebastian Böser
Colloquium | Grenoble | Sept. 8th 2015

A hundred year old puzzle...

1912 Victor Hess


- several balloon flights



A hundred year old puzzle...

1912 Victor Hess

- several balloon flights
- ionization **increases** with height!




A hundred year old puzzle...

1912 Victor Hess

- several balloon flights
- ionization **increases** with height!

“The results are best explained ... by the assumption that some radiation of high penetration power enters the atmosphere from above.”


V.Hess, Phys. Z (1912)



Cosmic rays today

Cosmic ray spectrum

- mostly protons / hadrons
- power law spectrum $F \propto E^{-2.7}$




Cosmic rays today

Cosmic ray spectrum

- mostly protons / hadrons
- power law spectrum $F \propto E^{-2.7}$

Normal energy (< GeV)

- the sun



Cosmic rays today

Cosmic ray spectrum


- mostly protons / hadrons
- power law spectrum $F \propto E^{-2.7}$

Normal energy (< GeV)

- the sun

High energy (GeV-PeV): ?

- **plausible** scenario
 - ▶ supernova remnants



Cosmic rays today

Cosmic ray spectrum

- mostly protons / hadrons
- power law spectrum $F \propto E^{-2.7}$

Normal energy (< GeV)


- the sun

High energy (GeV-PeV): ?

- **plausible** scenario
 - ▶ supernova remnants


Ultra-high energy (> PeV): ???

- **possible** scenarios
 - ▶ Active Galactic Nuclei
 - ▶ Gamma-ray burst
 - ▶ ...



Neutrinos from the sky


Cosmic Neutrino Sources



Gamma Ray Bursts



Active Galactic Nuclei



Supernova Remnants

Source power: up to 10^{45} Joule

Fermi Acceleration: Protons with energies up to 10^{19} eV


Particle interactions: in cosmic ray sources

$$p + p \rightarrow \pi, K, \dots$$

$$\quad \quad \quad \downarrow \pi \rightarrow \mu + \nu_{\mu}$$

$$\quad \quad \quad \quad \quad \downarrow \mu \rightarrow e + \nu_e + \nu_{\mu}$$


Cosmic Neutrino Sources



Gamma Ray Bursts



Active Galactic Nuclei



Supernova Remnants

Source power: up to 10^{45} Joule

Fermi Acceleration: Protons with energies up to 10^{19} eV

Particle interactions: in cosmic ray sources

$$p + p \rightarrow \pi, K, \dots$$

$$\downarrow \pi \rightarrow \mu + \nu_{\mu}$$

$$\downarrow \mu \rightarrow e + \nu_e + \nu_{\mu}$$

The cosmic ray – gamma – neutrino link

Propagation



- charged cosmic rays
 - deflected in magnetic fields
- photons / neutrinos
 - pointing back to source

Particle accelerators:

- electron accelerators
 - photons
- cosmic rays accelerators
 - neutrinos, photons

Neutrino detection


→ “smoking gun” for CR acceleration



Neutrino sources

Astrophysical neutrinos

- $p+\gamma/p$ interaction in the source
- e.g. synchrotron radiation
- ▶ generic spectrum $\sim E^{-2}$




Neutrino sources

Astrophysical neutrinos

- $p+\gamma/p$ interaction in the source
- e.g. synchrotron radiation
- ▶ generic spectrum $\sim E^{-2}$

Cosmogenic neutrinos

- $p+\gamma$ interaction with the CMB
- ▶ *guaranteed* neutrino flux



Neutrino sources

Astrophysical neutrinos


- $p+\gamma/p$ interaction in the source
- e.g. synchrotron radiation
- ▶ generic spectrum $\sim E^{-2}$

Cosmogenic neutrinos

- $p+\gamma$ interaction with the CMB
- ▶ *guaranteed* neutrino flux

Atmospheric neutrinos

- $p+p$ interaction in the atmosphere
- ▶ but π/K interact
- steeper spectrum $\sim E^{-3.7}$
- also *charmed mesons*
- ▶ prompt flux $\sim E^{-2}$



Neutrinos

mass →	$\approx 2.3 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$	$\approx 173.07 \text{ GeV}/c^2$	0	$\approx 126 \text{ GeV}/c^2$
charge →	$2/3$	$2/3$	$2/3$	0	0
spin →	$1/2$	$1/2$	$1/2$	1	0
	u up	c charm	t top	g gluon	H Higgs boson
	$\approx 4.8 \text{ MeV}/c^2$	$\approx 95 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$	0	
	$-1/3$	$-1/3$	$-1/3$	0	
	$1/2$	$1/2$	$1/2$	1	
	d down	s strange	b bottom	γ photon	
	$0.511 \text{ MeV}/c^2$	$105.7 \text{ MeV}/c^2$	$1.777 \text{ GeV}/c^2$	$91.2 \text{ GeV}/c^2$	
	-1	-1	-1	0	
	$1/2$	$1/2$	$1/2$	1	
	e electron	μ muon	τ tau	Z Z boson	
	$< 2.2 \text{ eV}/c^2$	$< 0.17 \text{ MeV}/c^2$	$< 15.5 \text{ MeV}/c^2$	$80.4 \text{ GeV}/c^2$	
	0	0	0	± 1	
	$1/2$	$1/2$	$1/2$	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	

QUARKS

LEPTONS

GAUGE BOSONS

Neutrinos

Fact sheet

- three flavors (ν_e , ν_μ , ν_τ)
- no charge
- tiny mass ($\sim$ eV)
- only interact via **weak** force
 - ▶ extremely low interaction rate

mass →	$\approx 2.3 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$	$\approx 173.07 \text{ GeV}/c^2$	0	$\approx 126 \text{ GeV}/c^2$
charge →	$2/3$	$2/3$	$2/3$	0	0
spin →	$1/2$	$1/2$	$1/2$	1	0
	u up	c charm	t top	g gluon	H Higgs boson
QUARKS	$\approx 4.8 \text{ MeV}/c^2$	$\approx 95 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$	0	
	$-1/3$	$-1/3$	$-1/3$	0	
	$1/2$	$1/2$	$1/2$	1	
	d down	s strange	b bottom	γ photon	
	$0.511 \text{ MeV}/c^2$	$105.7 \text{ MeV}/c^2$	$1.777 \text{ GeV}/c^2$	$91.2 \text{ GeV}/c^2$	
	-1	-1	-1	0	
	$1/2$	$1/2$	$1/2$	1	
	e electron	μ muon	τ tau	Z Z boson	
LEPTONS	$< 2.2 \text{ eV}/c^2$	$< 0.17 \text{ MeV}/c^2$	$< 15.5 \text{ MeV}/c^2$	$80.4 \text{ GeV}/c^2$	
	0	0	0	± 1	
	$1/2$	$1/2$	$1/2$	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	
					GAUGE BOSONS

Neutrinos

Fact sheet

- three flavors (ν_e , ν_μ , ν_τ)
- no charge
- tiny mass ($\sim$ eV)
- only interact via **weak** force
 - ▶ extremely low interaction rate

Advantage

- can escape from extreme environments

mass →	$\approx 2.3 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$	$\approx 173.07 \text{ GeV}/c^2$	0	$\approx 126 \text{ GeV}/c^2$
charge →	$2/3$	$2/3$	$2/3$	0	0
spin →	$1/2$	$1/2$	$1/2$	1	0
	u up	c charm	t top	g gluon	H Higgs boson
QUARKS	$\approx 4.8 \text{ MeV}/c^2$	$\approx 95 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$	0	
	$-1/3$	$-1/3$	$-1/3$	0	
	$1/2$	$1/2$	$1/2$	1	
	d down	s strange	b bottom	γ photon	
	$0.511 \text{ MeV}/c^2$	$105.7 \text{ MeV}/c^2$	$1.777 \text{ GeV}/c^2$	$91.2 \text{ GeV}/c^2$	
	-1	-1	-1	0	
	$1/2$	$1/2$	$1/2$	1	
	e electron	μ muon	τ tau	Z Z boson	
LEPTONS	$< 2.2 \text{ eV}/c^2$	$< 0.17 \text{ MeV}/c^2$	$< 15.5 \text{ MeV}/c^2$	$80.4 \text{ GeV}/c^2$	
	0	0	0	± 1	
	$1/2$	$1/2$	$1/2$	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	
					GAUGE BOSONS

Neutrinos

Fact sheet

- three flavors (ν_e , ν_μ , ν_τ)
- no charge
- tiny mass ($\sim$ eV)
- only interact via **weak** force
 - extremely low interaction rate

Advantage


- can escape from extreme environments

Disadvantage


- extremely hard to detect

mass →	$\approx 2.3 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$	$\approx 173.07 \text{ GeV}/c^2$	0	$\approx 126 \text{ GeV}/c^2$
charge →	$2/3$	$2/3$	$2/3$	0	0
spin →	$1/2$	$1/2$	$1/2$	1	0
	u up	c charm	t top	g gluon	H Higgs boson
	$\approx 4.8 \text{ MeV}/c^2$	$\approx 95 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$	0	
	$-1/3$	$-1/3$	$-1/3$	0	
	$1/2$	$1/2$	$1/2$	1	
	d down	s strange	b bottom	γ photon	
	$0.511 \text{ MeV}/c^2$	$105.7 \text{ MeV}/c^2$	$1.777 \text{ GeV}/c^2$	$91.2 \text{ GeV}/c^2$	
	-1	-1	-1	0	
	$1/2$	$1/2$	$1/2$	1	
	e electron	μ muon	τ tau	Z Z boson	
	$< 2.2 \text{ eV}/c^2$	$< 0.17 \text{ MeV}/c^2$	$< 15.5 \text{ MeV}/c^2$	$80.4 \text{ GeV}/c^2$	
	0	0	0	± 1	
	$1/2$	$1/2$	$1/2$	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	


Neutrino interactions




Detection principle



- Charged secondaries emit Cherenkov light
 - ▶ array of optical sensors




Detection principle




- Charged secondaries emit Cherenkov light
 - ▶ array of optical sensors

shopping list

- ✓ 1km^3 medium (transparent!!!)
- ✓ $\phi(10^3)$ optical sensors (stable!!!)





The IceCube Observatory


South Pole Station

Geographic South Pole

IceCube footprint


Skiway

The IceCube observatory




Digital optical module

- sensor 10'' PMT
Hamamatsu
- digital waveform
 $\delta t = 40\text{ns}$
- ambient
up to 200bar
down to -55°C




Cherenkov light

$$\nu_\mu + N \rightarrow \mu \text{ (100 TeV)}$$





Cherenkov light

$$\nu_\mu + N \rightarrow \mu \text{ (100 TeV)}$$



Ice properties



South Pole ice cap

- temperature profile
-55°C(top) to -10°C (bottom)
- dust layers
(e.g from volcanic ash)
- low radioactivity
 - ▶ $\emptyset(\text{Bq}/\text{m}^3)$





Light propagation

- scattering
- absorption
 - ▶ still very clear in optical

Attenuation length

- $\lambda_{\text{att}} \approx 45\text{m}$

Event reconstruction





Event signatures

Tracks

$$\nu_\mu + N \rightarrow \mu + X$$

- pointing resolution

$$\sigma_\Psi \sim 1^\circ$$





Cascades

$$\nu_e + N \rightarrow e + X$$

$$\nu_f + N \rightarrow \nu_f + X$$

- energy resolution


$$\sigma_E \sim 0.1 \log_{10}(E)$$




As well as...

- muon bundles (cosmic rays)
- double-bang (ν_τ)
- exotic states (monopoles,...)

Simulation




Background





Rates

- atm. μ : 2800 per second
- atm. ν : 0.1 per minute
- need $>10^6$ background rejection!



Background rejection




Rejection techniques

- event direction
- energy
- event topology


Point sources

ν_μ pointing resolution




Pointing resolution

- energy dependent
 - ▶ $\sigma(\theta_\nu) \approx 1^\circ$ at 1 TeV
 - ▶ dominated by ν - μ scattering angle




Pointing test

- moon shadow (13 σ /year)




Point source searches



Dataset

- livetime 1371 days (4 years) → 394,000 events after cuts
 - ▶ 40% up-going
 - ▶ 60% down-going (at high energy)

Significance skymap




Hottest spot

- DEC: 29.25°
- RA: 10.55°
- $-\log_{10}(p) = 5.318$
- $n_s = 43.04$, $\gamma = 2.88$
- no close-by candidates


Analysis

- calculate over-density (p-value) of real data
- scramble event directions
- count accidental occurrences
- 22% have equal or larger max. p-value



Unresolved sources

ν_μ energy spectrum



High Energy Starting Events (HESE)

Active veto layer


- reject events with early hit in veto layer
- keep *starting* events

Atmospheric flux

- **atmo. μ** rejected in veto
- **atmo. ν** rejected by
 - ▶ energy cut
 - ▶ μ companion

Result

- Optimized for 1 PeV region
 - ▶ factor 3 gain in sensitivity
- Sensitive to all flavors



HESE results

Dataset

- 1347 days (4 yrs)
full detector

Events


- 54 events
 - ▶ 15 track-like
 - ▶ 39 cascade-like

Background estimate

- **atmo. μ : 12.6 ± 5.1**
- **atmo. ν : 11.5 ± 5.7**

Significance

- well above 5σ



HESE results

Dataset

- 1347 days (4 yrs)
full detector

Events


- 54 events
 - ▶ 15 track-like
 - ▶ 39 cascade-like

Background estimate

- **atmo. μ : 12.6 ± 5.1**
- **atmo. ν : 11.5 ± 5.7**

Significance


- well above 5σ



HESE results

north vs. south?
→ earth becomes
opaque to neutrinos

correlation with
galactic plane?
→ most likely
band = 7.5°
→ 2.8% probability



hot spot?
→ 7.2% statistical
fluctuation

Flavour analysis

Flavours at source

■ regular decay

$$\begin{aligned}
 p(p/\gamma) &\rightarrow \pi^\pm/\pi^0 + \text{anything} \\
 \pi^\pm &\rightarrow \mu^\pm + \nu_\mu/\bar{\nu}_\mu \\
 &\hookrightarrow e^\pm + \nu_e/\bar{\nu}_e + \nu_\mu/\bar{\nu}_\mu
 \end{aligned}$$


► flavour ratio = 1:2:0

Flavours at earth

■ altered by **oscillations**

■ assume average over many sources

► flavour ratio $\sim 1:1:1$
(bi-maximal mixing)




Flavour analysis


Results

- **regular** decay
 - ▶ p-value = 30%
- **muon-suppressed**
 - ▶ p-value = 60%
- **neutron** decay
 - ▶ excluded at 3.2σ

$\nu_e - \nu_\tau$ degeneracy

- can not resolve ν_τ at TeV energies





Astrophysical neutrinos discovered!

What next?


The High Energy Extension (HEX)

Goal


- increase effective area by factor 5
- 50 events / year
 - ▶ find point sources

Method

- increase string-spacing
 - ▶ raise threshold



The Surface Extension (SurfEX)



Air shower veto


- increase A_{eff} by
 - ▶ 50% if effective at 300 TeV
 - ▶ 100% if effective at 30 TeV

... plus cosmic ray physics

Neutrinos on Earth

IceCube (in operation)

- 78 strings, 125m / 17m spacing
- $E_{\text{thresh}} \sim 100 \text{ GeV}$
 - optimized for astrophysical CR-sources
 - discovered!





IceCube's low-energy ν detector

IceCube (in operation)

- 78 strings, 125m / 17m spacing
- $E_{\text{thresh}} \sim 100 \text{ GeV}$

DeepCore (in operation)

- +8 strings, 72m / 7m spacing
- $E_{\text{thresh}} \sim 10 \text{ GeV}$





IceCube's low-energy ν detector

IceCube (in operation)

- 78 strings, 125m / 17m spacing
- $E_{\text{thresh}} \sim 100 \text{ GeV}$

DeepCore (in operation)

- +8 strings, 72m / 7m spacing
- $E_{\text{thresh}} \sim 10 \text{ GeV}$



Atmospheric neutrinos

Event rates (trigger level)


■ $\nu_\mu \sim 10^5$ / year

■ $\nu_e \sim 10^3$ / year

→ Huge data sample

■ cosmic ray physics

■ ν -oscillation physics



Neutrino Oscillations

Weak interactions couple to superpositions of neutrino mass eigenstates:

$$|\nu_e\rangle = \cos\theta |\nu_1\rangle + \sin\theta |\nu_2\rangle$$

$$|\nu_\mu\rangle = -\sin\theta |\nu_1\rangle + \cos\theta |\nu_2\rangle$$

Time evolution

$$|\nu_{1,2}\rangle \rightarrow e^{-iE_{1,2}t} |\nu_{1,2}\rangle$$

Flavor transition probability

$$\begin{aligned} P_{\mu \rightarrow e}(t) &\equiv |\langle \nu_e | e^{-i\hat{H}t} | \nu_\mu \rangle|^2 \\ &= \left| -\sin\theta \cos\theta e^{-iE_1t} + \sin\theta \cos\theta e^{-iE_2t} \right|^2 \\ &= \sin^2(2\theta) \sin^2\left(\frac{(E_1 - E_2)t}{2}\right) \end{aligned}$$

Neutrino Oscillations

Weak interactions couple to superpositions of neutrino mass eigenstates:

$$|\nu_e\rangle = \cos\theta |\nu_1\rangle + \sin\theta |\nu_2\rangle$$

$$|\nu_\mu\rangle = -\sin\theta |\nu_1\rangle + \cos\theta |\nu_2\rangle$$

Time evolution

$$|\nu_{1,2}\rangle \rightarrow e^{-iE_{1,2}t} |\nu_{1,2}\rangle$$


Flavor transition probability

$$P_{\mu \rightarrow e}(t) \equiv |\langle \nu_e | e^{-i\hat{H}t} | \nu_\mu \rangle|^2$$

$$= \left| -\sin\theta \cos\theta e^{-iE_1t} + \sin\theta \cos\theta e^{-iE_2t} \right|^2$$

$$= \sin^2(2\theta) \sin^2\left(\frac{(E_1 - E_2)t}{2}\right)$$

Sound wave analogy



Neutrino Oscillations

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{(E_1 - E_2)t}{2} \right)$$

Using $m \ll E$ and $t \simeq L$:

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

Neutrino Oscillations

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{(E_1 - E_2)t}{2} \right)$$

Using $m \ll E$ and $t \simeq L$:


$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

$$\Delta m^2 \equiv m_2^2 - m_1^2$$

Neutrino Oscillations

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{(E_1 - E_2)t}{2} \right)$$

Using $m \ll E$ and $t \simeq L$:

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

Neutrino Oscillations

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{(E_1 - E_2)t}{2} \right)$$

Using $m \ll E$ and $t \simeq L$:

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

Neutrino Oscillations

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{(E_1 - E_2)t}{2} \right)$$

Using $m \ll E$ and $t \simeq L$:

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

Neutrino Oscillations

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{(E_1 - E_2)t}{2} \right)$$

Using $m \ll E$ and $t \simeq L$:

$$P_{e \rightarrow \mu}(t) = \sin^2(2\theta) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$


Three neutrino flavours:

- two Δm^2 and two signs (Δm_{21}^2 , Δm_{32}^2)
- three mixing angles (θ_{12} , θ_{23} , θ_{13})

Neutrinos vs. earth

Atmospheric neutrinos


- ν -oscillations in earth
 - ▶ wide range of L/E
- measure only ν_μ
 - ▶ sensitive to θ_{23} , Δm_{32}^2




1st oscillation minimum

■ $L_{\max} = 12 \cdot 10^3 \text{ km}$

■ $E_\nu \sim 20 \text{ GeV}$




Oscillation analysis



Oscillation result

Proof of principle
→ competitive
result!



What next?


Neutrino mass hierarchy

Oscillation parameters

- amplitude
 - ▶ mixing angles
- frequency
 - ▶ |mass splitting|

Order of mass eigenstates?



- matter effects
 - ▶ break degeneracy
- for $\nu_2 \leftrightarrow \nu_1$
 - ▶ solar neutrino oscillations ✓
- for $\nu_3 \leftrightarrow \nu_2$?



Neutrinos vs. earth


Atmospheric neutrinos

- ν -oscillations in earth
 - ▶ wide range of L/E
- matter effects
 - ▶ mass hierarchy



matter effects

- strongest in core
- $E_\nu \sim 3\text{--}15$ GeV





IceCube's low-energy ν extension

IceCube (in operation)


- 78 strings, 125m / 17m spacing
- $E_{\text{thresh}} \sim 100 \text{ GeV}$

PINGU (proposed)

- +40 strings, 25m / 3m spacing
- $E_{\text{thresh}} \sim 1 \text{ GeV}$
- neutrino mass hierarchy




Neutrino Mass Hierarchy



Neutrino Mass Hierarchy

Signal


- matter effects (MSW)
- inverted for $\nu \leftrightarrow \bar{\nu}$



Neutrino Mass Hierarchy

Signal


- matter effects (MSW)
 - ▶ inverted for $\nu \leftrightarrow \bar{\nu}$
- *similar* flux from atmosphere
 $\Phi(\nu)/\Phi(\bar{\nu}) \sim 1 \pm 0.1$



Neutrino Mass Hierarchy

Signal

- matter effects (MSW)
 - ▶ inverted for $\nu \leftrightarrow \bar{\nu}$
 - *similar* flux from atmosphere
 $\Phi(\nu)/\Phi(\bar{\nu}) \sim 1 \pm 0.1$
 - x-sections *differ*
 $\sigma(\nu N)/\sigma(\bar{\nu} N) \sim 2$
- few % effect



Neutrino Mass Hierarchy


Signal

- matter effects (MSW)
 - ▶ inverted for $\nu \leftrightarrow \bar{\nu}$
- *similar* flux from atmosphere
 $\Phi(\nu)/\Phi(\bar{\nu}) \sim 1 \pm 0.1$
- x-sections *differ*
 $\sigma(\nu N)/\sigma(\bar{\nu} N) \sim 2$

→ few % effect

Requirements


- high statistics
- very good control of systematic effects



Signatures in PINGU

Detected events (1-80GeV, after cuts)


- $\sim 50\text{k } \nu_\mu + \text{anti-}\nu_\mu$ per year
- $\sim 38\text{k } \nu_e + \text{anti-}\nu_e$ per year
 - ▶ small but distinct signature



Signatures in PINGU

Detected events (1-80GeV, after cuts)


- $\sim 50\text{k } \nu_\mu + \text{anti-}\nu_\mu$ per year
- $\sim 38\text{k } \nu_e + \text{anti-}\nu_e$ per year
 - ▶ small but distinct signature



Signatures in PINGU

Detected events (1-80GeV, after cuts)

- $\sim 50\text{k } \nu_\mu + \text{anti-}\nu_\mu$ per year
- $\sim 38\text{k } \nu_e + \text{anti-}\nu_e$ per year
 - ▶ small but distinct signature




Signatures in PINGU

Detected events (1-80GeV, after cuts)

- $\sim 50\text{k } \nu_\mu + \text{anti-}\nu_\mu$ per year
- $\sim 38\text{k } \nu_e + \text{anti-}\nu_e$ per year
 - ▶ small but distinct signature

Detector effects

- energy resolution
 - ▶ $\sim 20\%$ above 10GeV
- directional resolution
 - ▶ improving with E



Signatures in PINGU

Detected events (1-80GeV, after cuts)


- $\sim 50\text{k } \nu_\mu + \text{anti-}\nu_\mu$ per year
- $\sim 38\text{k } \nu_e + \text{anti-}\nu_e$ per year
 - ▶ small but distinct signature

Detector effects

- energy resolution
 - ▶ $\sim 20\%$ above 10GeV
- directional resolution
 - ▶ improving with E

Particle identification


- $\nu_\mu(\text{CC}) = \text{tracks}$
- $\nu_e(\text{CC}) + \nu_\chi(\text{NC}) = \text{cascades}$
 - ▶ quite distinct



Results: neutrino mass hierarchy

Analysis results


- for worst case ($\theta_{23} = 42.3^\circ$)
- hierarchy significance
 - ▶ $\sim 3\sigma$ in 3.5 years




Comparison to others

- PINGU could be the **first** detector to make this challenging measurement


Outlook



Outlook



Outlook






Conclusion

High-energy neutrinos can tell us about the sources of cosmic rays
→ IceCube just found first neutrinos from outside solar system

Oscillation of low-energy neutrinos can tell us about their
fundamental properties
→ IceCube can also measure these

IceCube-Gen2 will bring major advances for both of these fields



Thank you!





Performance

Installation


- 5484 optical modules installed
 - ▶ 128 DOMs failed (during installation)
 - ▶ 50% fixed again
- expected statistical lifetime
 - ▶ 14.000 years

Data-taking

- average detector uptime
 - ▶ >99% last years
- used for data analysis
 - ▶ >95% over 5 years



ν_μ pointing resolution




Pointing resolution

- improves with energy
- dominated by ν - μ scattering angle


Energy resolution

- estimated from dE/dx of muons
 - dominated by stochastic losses
- $\rightarrow \sigma(E_\mu) \approx 0.3 \log_{10}(E_\mu)$



Pointing verification

- $\pm 4^\circ$ window around moon
- $\rightarrow 13\sigma$ underfluctuation in IC59 (14 moon cycles)




Combined analysis

Results from multiple searches




- 2 dedicated track searches
- 2 dedicated cascade searches
- high- and medium energy starting event searches
 - ▶ all are consistent

Best fit result

- spectral index
 - ▶ $\gamma = 2.50 \pm 0.09$
- matches predictions for Fermi acceleration
- no indication of break in power law



Geometry studies



Constraints

- South Pole sectors
 - ▶ radial assymetrie
- optical ice properties / bedrock
 - ▶ only slightly thicker detector
- polar assymetrie



Wish-list

- active veto technique
 - ▶ avoid regular grid
- no *channels* for background

Expected performance

Tracks

- same resolution for horizontal events
- ▶ region with
 - low μ -background ✓
 - little earth absorption ✓



Cascades

- effective area grows with
 - ▶ volume
 - ▶ veto efficiency
- additional help from Glashow resonance




Transient sources

Transient sources

Gamma ray bursts

Fireball model (long GRBs)


- collapse of massive star
- total energy release $\sim 10^{45}$ Joules
 - ▶ candidate for cosmic rays
- ultra-relativistic jets
 - ▶ PeV neutrino emission ??



IceCube GRB search

- coincidence analysis
 - ▶ time window ($\Delta T \approx 0.1\text{-}100\text{s}$)
 - ▶ direction ($\Delta\Psi_{\text{IceCube}} \approx 1^\circ$)
- per-alert emission model
 - 5.2 events expected
 - **no** events observed

→ starting to constrain GRB models



Follow-up program

Idea

- trigger follow-up observation by neutrinos in IceCube
→ online neutrino analysis

transient
event
(SN, GRB, ...)





Online alerts

- multiplet trigger
 $N_v \geq 2$, $\Delta\Psi \geq 4^\circ$, $\Delta t \leq 100\text{s}$ separation

Follow-up program

- optical: ROTSE, PTF → first results
- X-rays: SWIFT
- γ -rays: MAGIC, VERITAS

northern
hemisphere
institute





IceCube online neutrino filter

Neutrino multiplets

	N_ν	Measured	Expected
IC 40	2	15	8.53
	3	0	0.003
IC 59	2	19	15.66
	3	0	0.004
IC79	2	32	42.5
	3	0	0.012
IC86	2	24	19.59
	3	0	0.005

general muon
filter

high-quality
up-going events

neutrino
selection

90% atmospheric
neutrinos

Trigger
~3000 Hz

Level 1
~30 Hz

Level 2
~3 Hz

Level 3
~2 mHz

Latency


- 3 minutes

Optical follow-up – first results

Supernova jet model

- Are there GRB – like jets in core-collapse SNe?


[Ando & Beacom (PRL 95/2005)]



First year of optical follow-up

- 34 alerts sent to ROTSE
- 0 SN counterparts observed

[Abbasi et al., A&A 2012]





Summary

Motivation

Neutrinos & Cosmic rays

- Neutrinos from heaven
 - Point sources
 - Transient sources
 - Diffuse flux
- High-Energy extension

Neutrino from earth

- PINGU

