

# 48<sup>ème</sup> Ecole de Gif : La Physique Souterraine

## Low Energy Neutrinos Underground Experiments

**Mathieu BONGRAND**

LAL Orsay

September, 2016



Introduction

Solar Neutrinos

Atmospheric Neutrinos

Reactor Neutrinos

Other Neutrinos Detections

## Introduction

Solar Neutrinos


Atmospheric Neutrinos

Reactor Neutrinos

Other Neutrinos Detections

# Neutrino Experiment

- ▶ To build a neutrino experiment, besides the phenomena we want to study, we need to define:
  - ▶ the expected number of neutrino events from a given source
  - ▶ how they would interact in our detector







$$N(E_\nu) = \phi_\nu(E_\nu) \cdot \sigma_\nu(E_\nu) \cdot \text{target}$$

- ▶ The weak interaction neutrino cross-sections  $\sigma_\nu(E_\nu)$  are tiny:
  - ▶ but  $\sigma_\nu(E_\nu)$  increases with  $E_\nu$
  - ▶ we need intense neutrino sources (huge  $\phi_\nu(E_\nu)$ )
  - ▶ target detector has to be (very) large (few t  $\rightarrow$  kt)



# Neutrino Sources


The neutrino is the 2<sup>nd</sup> most abundant particle in the Universe:  
 $\sim 300 \nu_{relic} \text{ cm}^{-3}$  (1<sup>st</sup> are the CMB photons)



**Neutrino Budget on Earth**

|              |                                         |                                |                     |                                                    |
|--------------|-----------------------------------------|--------------------------------|---------------------|----------------------------------------------------|
| Sun          | $\nu_e$                                 | $1.5 \cdot 10^8 \text{ km}$    | continuous          | $6.5 \cdot 10^{10} \text{ cm}^{-2} \text{ s}^{-1}$ |
| SN 1987A     | $\bar{\nu}_e$                           | $1.5 \cdot 10^{18} \text{ km}$ | $\sim 10 \text{ s}$ | $\sim 5 \cdot 10^8 \text{ cm}^{-2} \text{ s}^{-1}$ |
| Atmospheric  | $\nu_{e\mu\tau} / \bar{\nu}_{e\mu\tau}$ | $10\text{-}10\,000 \text{ km}$ | continuous          | $\sim 1 \text{ cm}^{-2} \text{ s}^{-1}$            |
| Geoneutrinos | $\bar{\nu}_e$                           | $0 - 6370 \text{ km}$          | continuous          | $2.0 \cdot 10^6 \text{ cm}^{-2} \text{ s}^{-1}$    |
| Reactor      | $\bar{\nu}_e$                           | $\sim 1 \text{ km}$            | continuous          | $1.0 \cdot 10^{10} \text{ cm}^{-2} \text{ s}^{-1}$ |
| Accelerators | $\nu_\mu / \bar{\nu}_\mu$               | $200 - 800 \text{ km}$         | pulsed              |                                                    |




# Neutrino Fluxes and Energies on Earth



G. Giacomelli & M. Sioli, arXiv:hep-ex/0211035v1, 2002

# The Discovery of the Neutrino

- ▶ 1934 - E. Fermi theory of  $\beta$ -decay
- ▶ 1934 - H. Bethe and R. Peierls understand that  $\nu$  can invert the  $\beta$ -decay
- ▶ 1950's C. Cowan & F. Reines start the "Projet Poltergeist"
- ▶ delayed coincidence between  $e^+$  annihilation and  $n$  radiative capture
- ▶ 1953 - First experiment at Hanford reactor not conclusive
- ▶ 1955 - Second experiment at Savannah River reactor






- ▶ 1956 - Discovery of the neutrino → telegram to Pauli

# Inverse $\beta$ -decay (IBD)

IBD still used for reactor and low energy  $\bar{\nu}_e$  experiments:

- ▶ very low background thanks to the delayed coincidence
- ▶ the antineutrino is stopped and its energy is given by the energy of the positron:  $E_{\bar{\nu}_e} = E_{e^+} + m_n - m_p$
- ▶ energy threshold:  $E_{\bar{\nu}_e}^{min} = m_n - m_p + m_{e^+} = 1.8 \text{ MeV}$
- ▶ incoming direction of the  $\bar{\nu}_e$  with the forward boost of the  $n$ ?




C. Bemporad, G. Gratta, P. Vogel, Rev. Mod. Phys. 74:297, 2002  
M. Apollonio et al., Eur. Phys. J. C27:331-374, 2003

# Neutrino-Electron Elastic Scattering (ES)


ES is used in most of the neutrino experiments (Solar, SN, atmospheric...):  $\nu_\alpha e^- \rightarrow \nu_\alpha e^-$  or  $\bar{\nu}_\alpha e^- \rightarrow \bar{\nu}_\alpha e^-$

- ▶ very well known cross sections increasing with  $E_\nu$
- ▶ sensitive to all the flavors of  $\nu$  and  $\bar{\nu}$
- ▶  $\sigma_{\nu_e} \simeq 2.4 \sigma_{\bar{\nu}_e}$   $\sigma_{\nu_e} \simeq 6.2 \sigma_{\nu_\mu \nu_\tau}$   $\sigma_{\nu_e} \simeq 7.1 \sigma_{\bar{\nu}_\mu \bar{\nu}_\tau}$
- ▶ incoming  $\nu$  direction is preserved in the  $e^-$  recoil:  $E_e^2 \theta_e^2 < 2m_e$
- ▶ energy threshold due to detection technique



Legend: solid  $\nu_e$ , dashed  $\bar{\nu}_e$ , dotted  $\nu_{\mu,\tau}$ , dash-dotted  $\bar{\nu}_{\mu,\tau}$

C. Giunti & C. W. Kim, Fundamentals of Neutrino Physics and Astrophysics, Oxford, 2007



# Neutrino-Nucleon Scattering

At higher neutrino energies the scattering occurs on nucleons:


$$\text{CC: } \nu_{\alpha} N \rightarrow \alpha^{-} X \quad \bar{\nu}_{\alpha} N \rightarrow \alpha^{+} X$$

$$\text{NC: } \nu_{\alpha} N \rightarrow \nu_{\alpha} X \quad \bar{\nu}_{\alpha} N \rightarrow \bar{\nu}_{\alpha} X$$

$$\alpha = e, \mu, \tau \quad N = p, n \quad X = \text{hadrons}$$


- ▶ In elastic or Quasi-Elastic (QE) scattering  $X = p, n$
- ▶ Nuclear Resonance Production creates excited states  
 $X = N^{*} \quad \text{or} \quad \Delta \rightarrow p \pi^{0} / n \pi^{+}$
- ▶ In Deep Inelastic Scattering (DIS) the nucleon is broken

Correct knowledge of the different  $\sigma$  but improvements are coming



# Neutrino Mixing & Oscillations

- ▶ Neutrino oscillations were thought to solve the long-standing missing solar (R. Davis ~1970) or atmospheric neutrinos (SK ~2000):
  - ▶ experiments are generally sensitive to only one neutrino flavor
  - ▶ if a neutrino flavor change during its propagation from the source to the detector: the neutrino won't be detected



- ▶ Oscillations already observed in particle physics:  $K^0 - \bar{K}^0$ 
  - ▶ occur when flavor eigenstates are not mass eigenstates
- ▶ **So neutrinos are massive particles!**
- ▶ In the late 1950s B. Pontecorvo started to study neutrino oscillations

# Two Flavors Neutrino Oscillations

- ▶ Because of the neutrino mass splitting  $\Delta m_{ji}$ , the fixed neutrino energy  $E$  and baseline  $L$ : most of the experiments are only sensitive to the two flavor neutrino mixing (or an effective mixing)
- ▶ Much simpler to compute and to illustrate the oscillations
- ▶ Two flavors neutrino mixing parametrized by:
  - ▶ a single squared-mass difference:  $\Delta m^2 = \Delta m_{ji}^2 = m_j^2 - m_i^2$
  - ▶ a single mixing angle:  $\theta \in [0, \pi/2]$

$$\begin{pmatrix} \nu_\alpha \\ \nu_\beta \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \nu_i \\ \nu_j \end{pmatrix}$$

- ▶ From this matrix we write the transition probability ( $\alpha \neq \beta$ ):

$$\begin{aligned} P_{\nu_\alpha \rightarrow \nu_\beta}(L, E) &= \frac{1}{2} \sin^2 2\theta \left[ 1 - \cos \left( \frac{\Delta m^2 L}{2E} \right) \right] \\ &= \sin^2 2\theta \sin^2 \left( \frac{\Delta m^2 L}{4E} \right) \end{aligned}$$


- ▶ The survival probability ( $\alpha = \alpha$ ) is given by:  $1 - P_{\nu_\alpha \rightarrow \nu_\beta}(L, E)$



# Two Flavors Neutrino Oscillations Probability

- For experiments it is convenient to use:


$$P_{\nu_\alpha \rightarrow \nu_\beta}(L, E) = \sin^2 2\theta \sin^2 \left( 1.27 \frac{\Delta m^2 [\text{eV}^2] L [\text{m}]}{E [\text{MeV}]} \right)$$




- The peak position gives the squared-mass difference  $\Delta m^2$
- The peak amplitude gives the mixing angle  $\theta$

# Two Kind of Oscillation Experiments

- Appearance experiment: transition probability  $P_{\nu_\alpha \rightarrow \nu_\beta}(L, E)$



- Disappearance experiment: survival probability  $P_{\nu_\alpha \rightarrow \nu_\alpha}(L, E)$



# Three Flavors Neutrino Mixing

3×3 flavors neutrino mixing matrix *PMNS*  
(similar to *CKM* mixing matrix for quarks):

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{13}c_{12}s_{23}e^{+i\delta} & c_{23}c_{12} - s_{13}s_{12}s_{23}e^{+i\delta} & c_{13}s_{23} \\ s_{23}s_{12} - s_{13}c_{12}c_{23}e^{+i\delta} & -s_{23}c_{12} - s_{13}s_{12}c_{23}e^{+i\delta} & c_{13}c_{23} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

$s_{ij} = \sin \theta_{ij}$  and  $c_{ij} = \cos \theta_{ij}$

Pontecorvo, Maki, Nakagawa & Sakata

Three flavors neutrino mixing parametrized by:

- ▶ 2 squared-mass difference:  $\Delta m_{32}^2$ ,  $\Delta m_{21}^2$
- ▶ 3 mixing angles:  $\theta_{12}$ ,  $\theta_{23}$ ,  $\theta_{13}$
- ▶ 1 CP violation phase:  $\delta$
- ▶ 2 Majorana phases:  $\eta_1$ ,  $\eta_2$ ?

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & s_{13}e^{-i\delta} & \\ -s_{13}e^{i\delta} & c_{13} & \\ & & 1 \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & \\ -s_{12} & c_{12} & \\ & & 1 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

atmospheric
reactor/accelerator
solar

Introduction

Solar Neutrinos

Atmospheric Neutrinos

Reactor Neutrinos




Other Neutrinos Detections

# Solar Neutrinos

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix}}_{\text{atmospheric}} \underbrace{\begin{pmatrix} c_{13} & s_{13} e^{-i\delta} & \\ -s_{13} e^{i\delta} & c_{13} & \\ & & 1 \end{pmatrix}}_{\text{reactor/accelerator}} \underbrace{\begin{pmatrix} c_{12} & s_{12} & \\ -s_{12} & c_{12} & \\ & & 1 \end{pmatrix}}_{\text{solar}} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

# Solar Neutrinos Production

Solar neutrinos  $\nu_e$  are produced in the core by nuclear fusion and neutrinos in the  $pp$  chain and the CNO cycles




# Solar Neutrinos Spectrum

Predicted by J. Bahcall from the 1960's

${}^7\text{Be}$  and  $pep$  neutrinos have characteristic energy

$pp$ , CNO,  ${}^8\text{B}$  and  $hep$  have a continuous spectrum





# First Detection of the Solar Neutrinos

R. Davis built Chlorine experiment at Homestake (USA)

400 000 L of  $\text{C}_2\text{Cl}_4$  from 1970 to 1994

$\nu_e \text{ }^{37}\text{Cl} \rightarrow e^- \text{ }^{37}_{18}\text{Ar}$  (threshold 814 keV)





Only  $\sim 30\%$  of Bahcall's solar neutrino flux prediction

(Solar Neutrino Unit:  $1 \text{ SNU} = 10^{-36} \text{ capture/target atom/s}$ )



# The Sudbury Neutrino Observatory (SNO)

1 kt heavy water ( $D_2O$ ) 12 m spherical Cherenkov detector at Creighton mine in Sudbury (Canada) under 6 000 m.w.e.



# SNO Neutrino Detection

Heavy water ( $D_2O$ ) allows neutrino detection through 3 reactions:

$$\text{CC:} \quad \nu_e d \rightarrow p p e^- \quad E_\nu > 1.4 \text{ MeV}$$

$$\text{NC:} \quad \nu_\alpha d \rightarrow p n \nu_\alpha \quad E_\nu > 2.2 \text{ MeV}$$

$$\text{ES:} \quad \nu_\alpha e^- \rightarrow e^- \nu_\alpha \quad E_\nu > 5.5 \text{ MeV}$$

3 different phases for the  $n$  detection:

- ▶ 1999-2001:  $n d \rightarrow {}^3\text{H} \gamma$  (6.3 MeV)
- ▶ 2001-2003:  $n {}^{35}\text{Cl} \rightarrow {}^{36}\text{Cl} \gamma$  (8.6 MeV)
- ▶ 2004-2006:  ${}^3\text{H}$  proportional counters



# SNO Main Result

Sensitive to all neutrino flavors and using 3 interactions SNO reconstructed the full solar neutrino flux and solved the 30 y problem

$$\phi_{ES} = 2.35 \pm 0.22 \pm 0.15 \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

$$\phi_{CC} = 1.68 \pm 0.06 \pm_{-0.09}^{+0.08} \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

$$\phi_{NC} = 4.94 \pm 0.21 \pm_{-0.34}^{+0.38} \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$





The Solar neutrino problem is due to  $\nu_e \rightarrow \nu_{\mu\tau}$  oscillations

Phys. Rev. C72:055502, 2005


# The MSW Effect

S. Mikheyev, A. Smirnov and L. Wolfenstein effect

$$\mathcal{H} = \frac{\Delta m_{21}^2}{4E} \begin{pmatrix} -\cos 2\theta_{12} & \sin 2\theta_{12} \\ \sin 2\theta_{12} & \cos 2\theta_{12} \end{pmatrix} + \sqrt{2} G_F N_e \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$






C. Giunti & C. W. Kim, Fundamentals of Neutrino Physics and Astrophysics, Oxford, 2007



Give the sign of  $\Delta m_{12}^2$   
neutrino mixing  
parameter !

# Super-Kamiokande Experiment





50 kt water Cherenkov - Kamioka in Japan -  $\sim 13\,000$  PMTs 20"



$$\nu_{\mu} N \rightarrow \mu X$$

$$\begin{aligned} \nu_{\alpha} e^{-} &\rightarrow e^{-} \nu_{\alpha} \\ \nu_e N &\rightarrow e^{-} X \end{aligned}$$

# Super-Kamiokande Solar Neutrinos



Phys. Rev. D 78, 032002, 2008 &  
arXiv:1606.07538

# Borexino

300 t liquid scintillator target at LNGS -  $\sim 2200$  PMTs - 500 PE/MeV

$\nu_\alpha e^- \rightarrow \nu_\alpha e^-$  detection is identical to  $\beta$  and  $\gamma$  backgrounds

$\rightarrow$  extremely low background detector

## Borexino Detector






photo: BOREXINO collaboration

# Borexino Solar Neutrinos Results




Very low background and low threshold experiment  $\sim 150$  keV  
First detections of  ${}^7\text{Be}$ ,  $pep$  and  $pp$  neutrinos - limits on  $CNO$



$$\chi^2/\text{d.o.f} = 172.3/147$$

- $pp$  v:  $144 \pm 13$  (free)
- ${}^7\text{Be}$  v:  $46.2 \pm 2.1$  (constrained)
- $pep$  v:  $2.8$  (fixed)
- $CNO$  v:  $5.36$  (fixed)
- ${}^{214}\text{Pb}$ :  $0.06$  (fixed)
- ${}^{210}\text{Po}$ :  $583 \pm 2$  (free)
- ${}^{14}\text{C}$ :  $39.8 \pm 0.9$  (constrained)
- Pile-up:  $321 \pm 7$  (constrained)
- ${}^{210}\text{Bi}$ :  $27 \pm 8$  (free)
- ${}^{85}\text{Kr}$ :  $1 \pm 9$  (free)

Fit range: 165-590 keV







arXiv:1308.0443 & arXiv:1507.02432



# KamLAND

Antineutrino reactor experiment sensitive to solar oscillation parameters



$$\bar{\nu}_e : \langle E_\nu \rangle \sim 3 \text{ MeV} - L \sim \langle 180 \rangle \text{ km}$$



# KamLAND Solar Results

Reactor antineutrinos oscillation  
disappearance spectrum:  $\bar{\nu}_e \rightarrow \bar{\nu}_\alpha$


$$P_{ee}^{3\nu} = \cos^4 \theta_{13} \times \left[ 1 - \sin^2 2\theta_{12M} \sin^2 \left( \frac{\Delta m_{21}^2 L}{4E_\nu} \right) \right] + \sin^4 \theta_{13}$$



Phys. Rev. D83:052002, 2011 & arXiv:1303.4667

# Solar Neutrinos Global Fit

$\theta_{13} \neq 0$  reduced the solar and KamLAND  $\theta_{12}$  tension  
 However a small tension remain in  $\Delta m_{21}^2$




PDG values:  
 $\sin^2 2\theta_{12} = 0.846 \pm 0.021$   
 $\Delta m_{12}^2 = (7.53 \pm 0.18) 10^{-5} \text{ eV}^2$

# Transition between MSW and vacuum solar $\nu$ oscillations

At low energy:  $P_{ee} \rightarrow P_{ee}^{vac}$  and matter effects dominate at high energy

Intermediate region is where MSW resonance takes place

This region transition is determined by  $\Delta m_{21}^2$



Eur. Phys. J. A 52 (2016) 87 & arXiv:1507.05287

Introduction

Solar Neutrinos

Atmospheric Neutrinos


Reactor Neutrinos

Other Neutrinos Detections

# Atmospheric Neutrinos

Started with atmospheric neutrinos measurements but is now dominated by accelerator neutrino beam experiments (MINOS, T2K, No $\nu$ a...)




$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix}}_{\text{atmospheric}} \underbrace{\begin{pmatrix} c_{13} & s_{13} e^{-i\delta} & \\ -s_{13} e^{i\delta} & c_{13} & \\ & & 1 \end{pmatrix}}_{\text{reactor/accelerator}} \underbrace{\begin{pmatrix} c_{12} & s_{12} & \\ -s_{12} & c_{12} & \\ & & 1 \end{pmatrix}}_{\text{solar}} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$



# Atmospheric Neutrinos

1912 - V. F. Hess discovers the cosmic rays in his balloon

During 100 years intensively studied





The ratio between  $\nu_e$  and  $\nu_\mu$  events is well known from the cosmic rays interacting in the atmosphere

$$R = \frac{\phi(\nu_\mu + \bar{\nu}_\mu)}{\phi(\nu_e + \bar{\nu}_e)} \quad R \simeq 2 \text{ for } E_\nu < 1 \text{ GeV and } R > 2 \text{ for } E_\nu > 1 \text{ GeV}$$

# Super-Kamiokande Atmospheric Neutrinos

Super-Kamiokande also found an anomaly between the predicted ratio between  $\nu_e$  and  $\nu_\mu$  events



PDG values:

$$\sin^2 2\theta_{23} = 0.999^{+0.001}_{-0.018}$$

$$|\Delta m_{32}^2| = (2.42 \pm 0.06) 10^{-3} \text{ eV}^2$$

Upward going  $\nu_\mu$  ( $\cos\theta < 0$ ) oscillated into  $\nu_\tau$  but not  $\nu_e$



Introduction

Solar Neutrinos

Atmospheric Neutrinos

Reactor Neutrinos

Other Neutrinos Detections



# Reactor Neutrinos

KamLAND is a reactor neutrino experiment at long baseline measuring solar neutrino oscillations

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & s_{13} e^{-i\delta} & \\ -s_{13} e^{i\delta} & c_{13} & \\ & & 1 \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} \\ -s_{12} & c_{12} \\ & & 1 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

atmospheric
reactor/accelerator
solar

$\theta_{13}$  mixing angle was limited by Chooz (1997-1998) limit  $\sin^2 2\theta_{13} < 0.17$



Important for the search of CP violation ( $\delta$  phase)

3 new generation reactor experiments looking for very low  $\theta_{13}$ : Double Chooz, Daya-Bay & RENO


# Reactor Neutrino Experiments

Multidetector experiments to cancel neutrino flux systematics:

- ▶ near detector measure the non oscillated neutrino flux
- ▶ far detector measure the amplitude of the oscillation

Given atmospheric results on  $\Delta m_{31}^2$  :  $\sim 1$  km baseline is needed to measure  $\theta_{13}$



$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \simeq 1 - \sin^2 2\theta_{13} \sin^2 \left( \frac{\Delta m_{31}^2 L}{4E} \right) - \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2 \left( \frac{\Delta m_{21}^2 L}{4E} \right)$$



# Double Chooz Detector Example

Shallow depth underground laboratories  
( $<1000$  m.w.e.)

8 t Gd-LS  $\bar{\nu}_e$  target



**Outer veto:** plastic scintillator strips

**Shielding:** 15 cm of steel

**Inner veto:**

90 m<sup>3</sup> of liquid scintillator & 78 PMTs  
[40% LAB, 60 % Tetradecane, PPO, Bis-MSB]

**Buffer:**


110 m<sup>3</sup> of non scintillating mineral oil  
& 390 PMTs  
[30% Tetradecane, 70% Oil]

**Gamma catcher:**

22.3 m<sup>3</sup> of liquid scintillator  
[46% Dodecane, 50% Oil, 4% PXE, PPO, Bis-MSB]

**Neutrino target:**

10.3 m<sup>3</sup> of liquid scintillator doped with  
0.1 g/L of Gd  
[80% Dodecane, 20% PXE, PPO, Bis-MSB, Gd]




## Signal:

- ▶ prompt  $e^+$  and delayed  $n$ -capture
- ▶ rate:  $\sim 40$  in far and  $\sim 300$  IBD/d in near




## Backgrounds:

- ▶ accidental:  $\gamma + n$
- ▶ fast- $n$ :  $p$ -recoil +  $n$ -capture
- ▶ stopping  $\mu$ :  $\mu \rightarrow e \nu_e \nu_\mu$
- ▶ ( $\beta, n$ ) emitters:  $^9\text{Li}$  &  $^8\text{B}$
- ▶ rate:  $\sim 1$  in far and  $\sim 10$  in near bkg/d



# Oscillation result in DC

Matsubara ICHEP 2016



Best-fit:  $\sin^2 2\theta_{13} = 0.111 \pm 0.018$  ( $\chi^2/\text{dof} = 128.8/120$ )


Non-zero  $\theta_{13}$  observation at  $5.8 \sigma$  C.L.

Prospects:


Improving systematics to 0.01

Increasing statistics including



$\gamma$ -catcher  $n$ -captures on H





# Daya Bay Experiment



## 20 t Gd-LS $\bar{\nu}_e$ target




# Daya Bay $\theta_{13}$ Result



$\sim 2.6 \times 10^6$  IBD in near hall



$\sim 300 \times 10^3$  IBD in far hall








# Daya Bay $\Delta m_{13}^2$ Result

Thanks to multi-reactors and multi-detectors configuration  
Daya Bay can provide the most precise measurement of  $\Delta m_{13}^2$   
(previously know from atmospheric & accelerators)



# Similar Results from RENO Experiment in Korea





# Anti-neutrino energy spectrum reevaluation


Nuclear reactors are huge isotropic source of neutrinos

- ▶ 200 MeV and 6  $\bar{\nu}_e$  per fission
- ▶ short lived  $\beta$ -emitters  $\rightarrow$  follow the thermal power
- ▶  $\sim 2 \cdot 10^{20} \bar{\nu}_e \text{ s}^{-1} \text{ GW}_{\text{th}}^{-1}$

T. Mueller et al., Phys. Rev. C 83, 054615, 2011





- +3.5% new conversion of ILL  $\beta$  spectra:  $\nearrow \phi_{\bar{\nu}_e}$
- +1.5% off-equilibrium (long-lived isotopes):  $\nearrow \phi_{\bar{\nu}_e}$
- +1.5% neutron live-time  $\tau_n$  ( $880 \pm 1$  s):  $\nearrow \sigma_{\text{IBD}}$



# Reactor anti-neutrino anomaly

This reevaluation of the anti-neutrinos reactor spectrum for reactor experiments  $\rightarrow \sim 6.5\%$  deficit in short baseline experiments



G. Mention et al, PRD 83 (2011) 073006



K. N. Abazajian et al., arXiv:1204.5379

F. P. An et al., Phys. Rev. Lett. 116, 061801 (2016)

Possible explanation: the sterile neutrino  $\nu_s$

Sterile(s) neutrino(s) neutral lepton not sensitive to weak interaction:  
SM right-handed singlet which can have in principle any mass  
Can couple to active neutrinos through Lagrangian mass term





G. Mention et al, PRD 83 (2011) 073006



# Distortion of anti-neutrino spectrum “5 MeV bump”

All 3  $\theta_{13}$  reactor experiments observe an excess between 4 and 6 MeV

Y. Abe et al., JHEP 10 (2014) 086



F. P. An et al., Phys. Rev. Lett. 116, 061801 (2016)

# Mass Hierarchy with JUNO




arXiv:1507.05613

Ranucci Neutrino 2016

20 kt LS detector in China 3% energy resolution

Reactor experiment at 53 km for reactors with  $\sim 2100$  m.w.e.

Wide neutrino physics program



# Summary of Neutrino Oscillation measurements

The three flavor  $e, \mu, \tau$  neutrino mixing in the PMNS  $3 \times 3$  matrix describes well the measurements

**Solar:**  $\nu_e \rightarrow \nu_{\mu\tau}$

$$\sin^2 2\theta_{12} = 0.846 \pm 0.021$$

$$\Delta m_{12}^2 = (7.53 \pm 0.18) 10^{-5} \text{ eV}^2$$



**Atmospheric:**  $\nu_{\mu} \rightarrow \nu_{\tau}$

$$\sin^2 2\theta_{23} = 0.999^{+0.001}_{-0.018}$$

$$|\Delta m_{32}^2| = (2.42 \pm 0.06) 10^{-3} \text{ eV}^2$$

**Reactors:**  $\nu_e \rightarrow \nu_x$  or  $\nu_{\mu} \rightarrow \nu_e$

$$\sin^2 2\theta_{13} = 0.085 \pm 0.005$$



PDG 2014/2016

We still don't know absolute mass scale, mass hierarchy and  $\delta_{CP}$



Introduction

Solar Neutrinos

Atmospheric Neutrinos

Reactor Neutrinos


Other Neutrinos Detections

# Supernova Neutrinos

Supernova neutrinos detection is crucial for astrophysics models

Neutrinos from the supernova SN 1987A have been detected:

- ▶ 3 experiments Baksan 0.2 kt, Kamiokande 2.1 kt & IMB 6.8 kt
- ▶ in coincidence within a  $\sim 10$  s burst
- ▶ 25  $\bar{\nu}_e$  events with an average energy of 12.5 MeV
- ▶ total number of neutrinos emitted  $\sim 8 \cdot 10^{57}$
- ▶ energy released  $\sim 2 \cdot 10^{53}$  erg (90 %) in agreement with models



Actual experiments are ready for the next supernova


Addition of Gd for radiative neutron capture in Super-Kamiokande

Physics Letters B 205, 209-214, 1988

# Geoneutrinos in KamLAND

Half Earth's heat is due to natural radioactivity:





- ▶ total heat  $\sim 47 \pm 2$  TW
- ▶  $^{238}\text{U} \sim 8$  TW,  $^{232}\text{Th} \sim 8$  TW and  $^{40}\text{K} \sim 4$  TW



2005 KamLAND reported the 1<sup>st</sup> detection of geoneutrinos ( $\bar{\nu}_e$ )

Update in 2011 after  
Kashiwazaki–Kariwa shutdown

$106^{+29}_{-28}$  geoneutrino events



# Geoneutrinos in Borexino


arXiv:1506.04610

Borexino also detected geoneutrinos and presented an update in 2015

Less  $\bar{\nu}_e$  reactor background

Radiogenic heat production for U and Th

€ 23-36 TW



$23.7^{+6.5}_{-5.7}(stat) {}^{+0.9}_{-0.6}(sys)$   
geo-neutrino events

