Interpreting LHC Searches for New Physics with Simplified Models

Ursula Laa LPSC & LAPTh








Introduction

The Standard Model of Particle Physics





Open Questions:

- Particle description of dark matter?
- How do neutrinos obtain a mass?


Dark Matter

- Precision cosmology → measurement of relic abundance
- Planck 2015: $\Omega h^2 = 0.1188 \pm 0.0010$
- Naturally obtained for WIMPS ("WIMP miracle")
- WIMPS are predicted in many theories Beyond the Standard Model, e.g. in supersymmetric models





Supersymmetry



- Relating fermions and bosons
- New "superpartner" for all SM particles (same charges, spin differing by 1/2)
- R-parity → lightest partner (LSP) is stable
- SUSY broken → superpartners are heavy!
- Minimal (but general) model MSSM > 100 free parameters

Supersymmetric Dark Matter

Requirements

- Stable
- Invisible
- Reproduce observed relic density
- Consistent with bounds from direct and indirect detection experiments

Candidates

- Scalar neutrino (sneutrino) X
- cannot reproduce observed relic density
- excluded by direct detection experiments
- Partners of neutral electroweak gauge bosons and Higgs
- physical states are mixed
- label them as neutralinos $\tilde{\chi}^0$, with $\tilde{\chi}^0_1$ the lightest physical state
- good dark matter candidate, details depend on the mixing!

The corresponding charged superpartners also mix, we call them charginos. Depending on the nature of the LSP the lighter chargino can be almost mass degenerate with the LSP.

What about neutrino masses?

- Neutrino masses are not included in the MSSM
- Simplest implementation is Dirac mass term
- Requires right-handed (sterile) neutrinos
- Supersymmetric version: MSSM+RN

[Borzumati & Nomura, hep-ph/0007018] [Arkani-Hamed et al., hep-ph/0006312]

We can describe the sneutrino mixing as

$$\begin{pmatrix} \tilde{\nu}_{k_1} \\ \tilde{\nu}_{k_2} \end{pmatrix} = \begin{pmatrix} -\sin\theta_{\tilde{\nu}}^k & \cos\theta_{\tilde{\nu}}^k \\ \cos\theta_{\tilde{\nu}}^k & \sin\theta_{\tilde{\nu}}^k \end{pmatrix} \begin{pmatrix} \tilde{\nu}_L^k \\ \tilde{N}^k \end{pmatrix}$$

where the mixing angle is a free parameter

Mixed, mostly right-handed sneutrino LSP is a good dark matter candidate!

LHC Searches E Interpretation

Searching for Supersymmetry at the LHC (ATLAS & CMS)


Searching for SUSY

- LSP is invisible → "Missing energy" (MET) in signal events
- Concrete signatures depend on the scenario → many different analyses, targeting different types of scenarios
- For example "cut and count": define cuts that select signal events, reject SM background, compare observation to SM expectation
- LHC Run 1 All observations compatible with the SM
- Non-observation puts strong constraints on parameter space of supersymmetric models


Interpretation (Top Down)

- Consider "cut and count" type search
- Choose model and parameters
- Evaluate signal selection efficiency, predict number of signal events
- Compare to expected number of background events and the observation
- How can we interpret the results if even the minimal model has more than 100 free parameters?

Interpretation of LHC searches in constrained models



Interpretation of LHC searches in constrained models



Alternative: Bottom Up Approach


- Can we interpret LHC results in a "model independent" way ?
- Instead of considering a full model, consider Simplified Models

Simplified Model Spectra (SMS)

- Effective approach: only specific superpartners are light and considered for the interpretation
- Other superpartners are too heavy to be produced at the LHC
- Decay branching ratios are fixed
- Free parameters: physical mass and production cross section
- This captures the main features relevant for LHC phenomenology, results can be interpreted in mass planes, giving upper limits on the cross section


Example: Gluino Simplified Model


Production of gluino pair





Gluino decays 100 % as

$$\tilde{g} \to t t \tilde{\chi}_1^0$$











A closer look at the results

- The experiments evaluate an upper limit on the cross sections (for each mass bin)
- Those numbers still apply if the BR < 100 %</p>
- lacksquare We can compare the upper limits directly to σ * BR







[CMS-SUS-13-007]


[ATLAS-SUSY-2013-09]


But full models are complicated!




But full models are complicated!









New Features - made in Grenoble

"Missing Topologies"

SModelS identifies Simplified Model topologies with large cross section, not covered by the results in the database This information can be used to improve coverage of BSM searches and SMS interpretations

Input Checks

Consistency checks for input models specified in SLHA format, in particular do not apply Simplified Model results in scenarios with long-lived particles

LHC signatures of long-lived particles can also be treated within the SMS framework - and included in SModelS [Heisig, Lessa, Quertenmont, arXiv:1509.00473]

Application The MSSM+RN model

[Arina, Cabrera Catalan, Kraml, Kulkarni, UL, arXiv:1503.02960]

Setup

Sneutrino Dark Matter

All points consistent with measured relic abundance and limits from direct detection

Collider Constraints


- Use SModelS to study SMS constraints
- Sort out points with long-lived particles
- Study for example scenarios where chargino is light

We can describe the sneutrino mixing as

$$\begin{pmatrix} \tilde{\nu}_{k_1} \\ \tilde{\nu}_{k_2} \end{pmatrix} = \begin{pmatrix} -\sin\theta_{\tilde{\nu}}^k & \cos\theta_{\tilde{\nu}}^k \\ \cos\theta_{\tilde{\nu}}^k & \sin\theta_{\tilde{\nu}}^k \end{pmatrix} \begin{pmatrix} \tilde{\nu}_L^k \\ \tilde{N}^k \end{pmatrix}$$

where the mixing angle is a free parameter

sneutrino LSP



Setup

Sneutrino Dark Matter

All points consistent with measured relic abundance and limits from direct detection

Collider Constraints


- Use SModelS to study SMS constraints
- Sort out points with long-lived particles
- Study for example scenarios where chargino is light

We can describe the sneutrino mixing as

$$\begin{pmatrix} \tilde{\nu}_{k_1} \\ \tilde{\nu}_{k_2} \end{pmatrix} = \begin{pmatrix} -\sin\theta_{\tilde{\nu}}^k & \cos\theta_{\tilde{\nu}}^k \\ \cos\theta_{\tilde{\nu}}^k & \sin\theta_{\tilde{\nu}}^k \end{pmatrix} \begin{pmatrix} \tilde{\nu}_L^k \\ \tilde{N}^k \end{pmatrix}$$

where the mixing angle is a free parameter

sneutrino LSP




Scenarios with Light Chargino (and Neutralino)


Are sneutrino LSP scenarios constrained by results from LHC Run 1?


Which searches give the most important constraints?


Which Simplified Models are not yet considered but would improve the constraints?


Results — Overview






















→ Slepton search results constrain chargino production


Single lepton + MET is important topology in non-excluded points



Single lepton + MET is important topology in non-excluded points



Single lepton + MET is important topology in non-excluded points



Application The U(1)-extended MSSM

[Belanger, Da Silva, UL, Pukhov, arXiv:1505.06243]


Setup

■ Study of MSSM extension with extra U(1) gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y \times U'(1)$

- lacktriangle Extended particle contend, including in particular a Z'
- Constraints from SUSY searches included in terms of Simplified Model constraints
- Here: scenarios with "Higgs-like" LSP (higgsino, also possible in the MSSM) and light stop
- This differs from most Simplified Model assumptions because the lightest chargino is only slightly heavier than the LSP


Many points with light stops are still allowed

→ Check the missing topologies for non-excluded parameter points



Many points with light stops are still allowed


→ Check the missing topologies for non-excluded parameter points



Presentation of Experimental Results

ATLAS-SUSY-2013-15

CMS-SUS-14-011



Parametrised in terms of BR → this cannot be used to study alternative models

Conservative limit valid for any BR → could be included in SModelS

Conclusion

- Simplified Models
- useful tool for quick test of BSM models against constraints from LHC searches
- survey large parameter spaces
- SModelS
- automated decomposition procedure
- identify constraints, missing topologies, long-lived particles
- Sneutrino dark matter + light chargino
- main constraint from dilepton+MET searches
- important single lepton+MET Simplified Model not yet considered
- Higgsino LSP + light stop
- SMS constraints weak
- mixed topologies not fully considered

Work in Progress

- SModelS
- new verision, using also efficiency map type constraints, in preparation
- database can be extended using recasting tools, generating efficiency maps for SMS not considered by the experiments
- interface SModelS micrOMEGAs
- Testing the SModelS assumptions
- applicability of stop SMS results to fermionic top partners
- Beyond SMS
- use recasting tools to test LHC constraints on model with compressed spectra and/or long cascade decays

Backup

Reinterpretation of the Results

- To interpret the result of a search in an alternative model we need to determine the signal selection efficiency to predict an expected number of signal events
- This can be done using recasting frameworks, e.g. MadAnalysis5, CheckMate

Reinterpretation

Event simulation - MadGraph, Pythia, ...

Parton Level events

(fast) Detector simulation - Delphes

Reconstructed objects

Application of signal selection cuts

Cross section prediction - e.g. NLLfast

Point excluded / compatible with observation

Number of expected and observed events

cuts efficiency Selection efficiency

Reinterpretation of the Results

To interpret the result of a search in an alternative model we need to determine the signal selection efficiency to predict an expected number of signal events

This can be done using recasting frameworks, e.g.

MadAnalysis5, CheckMata

Reinterpretation

Event simulation - MadGr

Slow → not well suited to survey large parameter space

ss section ction - e.g. NLLfast

Parton Level events

(fast) Detector simulation - Delphes

Reconstructed objects

Application of signal selection cuts


Point excluded / compatible with observation

expected and

observed
events

Number of

LH Sneutrino Dark Matter?



The MSSM+RN Model

superpotential for Dirac RH neutrino superfield

$$W = \epsilon_{ij} (\mu \hat{H}_{i}^{u} \hat{H}_{j}^{d} - Y_{l}^{IJ} \hat{H}_{i}^{d} \hat{L}_{j}^{I} \hat{R}^{J} + Y_{\nu}^{IJ} \hat{H}_{i}^{u} \hat{L}_{j}^{I} \hat{N}^{J})$$

additional terms in the soft-breaking potential

$$V_{\text{soft}} = (M_L^2)^{IJ} \tilde{L}_i^{I*} \tilde{L}_i^J + (M_N^2)^{IJ} \tilde{N}^{I*} \tilde{N}^J$$
$$-[\epsilon_{ij} (\Lambda_l^{IJ} H_i^d \tilde{L}_j^I \tilde{R}^J + \Lambda_{\nu}^{IJ} H_i^u \tilde{L}_j^I \tilde{N}^J) + \text{h.c.}]$$

the sneutrino mass eigenstates are then given by


$$\begin{pmatrix} \tilde{\nu}_{k_1} \\ \tilde{\nu}_{k_2} \end{pmatrix} = \begin{pmatrix} -\sin\theta_{\tilde{\nu}}^k & \cos\theta_{\tilde{\nu}}^k \\ \cos\theta_{\tilde{\nu}}^k & \sin\theta_{\tilde{\nu}}^k \end{pmatrix} \begin{pmatrix} \tilde{\nu}_L^k \\ \tilde{N}^k \end{pmatrix}$$
with $\sin 2\theta_{\tilde{\nu}}^k = \sqrt{2} \frac{A_{\tilde{\nu}}^k v \sin\beta}{(m_{\tilde{\nu}}^2 - m_{\tilde{\nu}}^2)}$

SModelS description

Additional assumption Simplified Model results depend mainly on the mass spectrum of the new particles, not on specifics of the model (spin structure, production process, ...)





describe topology by vertex structure and outgoing SM particles in each vertex



[[[t,t]],[[t,t]]]

Signatures in Sneutrino LSP scenarios





gluino decay indistinguishable from neutralino LSP scenario

MSSM+RN Parameters sampled using Multinest

GUT scale Parameters	Prior range	
M_1, M_2	(-4000, 4000) GeV	
$log_{10}(M_3/{ m GeV})$	(-4, 4)	
$log_{10}(m_Q/{ m GeV})$	(2, 5)	
m_L, m_R	(1, 2000) GeV	
m_N	(1, 2000) GeV	
$log_{10}(A_Q/{ m GeV})$	(-5, 5)	
A_L	(-4000, 4000) GeV	
$A_{ ilde{ u}}$	(-1000, 1000) GeV	
$log_{10}(m_H/{ m GeV})$	(1, 5)	
$\tan eta$	(3, 50)	

MSSM + RN Selection Details

Observable	Value / constraint	
m_h	125.85 ± 0.4 (exp) ± 4 (theo) GeV	
$BR(B \to X_s \gamma) \times 10^4$	$3.55 \pm 0.24 \pm 0.09 \text{ (exp)}$	
$BR(B \to \mu^+ \mu^-) \times 10^9$	3.2 (+1.4 -1.2) (stat) (+0.5 -0.3) (sys)	
$\Omega_{ m DM} h^2$	0.1186 ± 0.0031 (exp) ± 20% (theo)	
$\Delta\Gamma_Z^{ m invisible}$	< 2 MeV (95% CL)	
$BR(h \to invisible)$	< 20% (95% CL)	
$m_{ ilde{ au}_1^-}$	> 85 GeV (95% CL)	
$m_{ ilde{\chi}_1^+}, m_{ ilde{e}, ilde{\mu}}$	>101 GeV (95% CL)	
$m_{ ilde{g}}$	> 308 GeV (95% CL)	
σ_n^{SI}	$<\sigma_{\mathrm{LUX}}^{SI}$ (90% CL)	

Application of Slepton Limits to Chargino Production

test this for ATLAS-SUSY-2013-11, using the MadAnalysis 5 implementation (B. Dumont, INSPIRE-1326686)


Compare the corresponding efficiencies in a benchmark scenario with


$$m_{mother} = 270 \text{ GeV}, m_{LSP} = 100 \text{ GeV}$$

Selection efficiencies somewhat different for individual cuts


Final efficiencies are comparable!

Cut	Slepton production	Chargino production	
Common preselection			
Initial number of events	50000	50000	
2 OS leptons	35133	33464	
$m_{ll} > 20 \text{ GeV}$	35038	33337	
τ veto	35007	33318	
ee leptons	35007	33318	
jet veto	20176	19942	
Z veto	19380	18984	
Different m_{T2} regions			
$m_{T2} > 90 \text{ GeV}$	11346	11594	
$m_{T2} > 120 \text{ GeV}$	8520	8828	
$m_{T2} > 150 \text{ GeV}$	5723	5926	







But many points are still allowed




But many points are still allowed




But many points are still allowed



MSSM + RN Cross section for single lepton missing topology




MSSM + RN most frequent missing topologies




Collider Searches and Direct Detection experiments are highly complementary!

Many points which are not at all tested by the SMS results are well in reach of XENON1T


Many parameter points below the neutrino background already excluded by SMS results



MSSM + RN Mixing Angle Dependence




MSSM + RN Long-lived Gluinos




Fraction of Models Excluded

In a more generic 19 - parameter model (pMSSM)




Fraction of Models Excluded

In a more generic 19 - parameter model (pMSSM)



UMSSM - many points with light gluino and LSP remain allowed

→ Check the missing topologies for non-excluded parameter points



UMSSM - many points with light gluino and LSP remain allowed

→ Check the missing topologies for non-excluded parameter points

