

The Leptonic CP Phases

Shao-Feng Ge

(gesf02@gmail.com)

Max-Planck-Institut für Kernphysik, Heidelberg, Germany

2017-5-22

SFG, Duane A. Dicus, Wayne W. Repko, PLB **702**, 220 (2011) [arXiv:1104.0602]

SFG, Duane A. Dicus, Wayne W. Repko, PRL **108**, 041801 (2012) [arXiv:1108.0964]

Andrew D. Hanlon, **SFG**, Wayne W. Repko, PLB **729**, 185-191 (2014) [arXiv:1308.6522]

SFG [arXiv:1406.1985]

Jarah Evslin, **SFG**, Kaoru Hagiwara, JHEP **1602** (2016) 137 [arXiv:1506.05023]

SFG, Pedro Pasquini, M. Tortola, J. W. F. Valle, PRD **95** (2017) No.3, 033005 [arXiv:1605.01670]

SFG, Alexei Smirnov, JHEP **1610** (2016) 138 [arXiv:1607.08513]

SFG [arXiv:1704.08518]

SFG, Manfred Lindner, PRD **95** (2017) No.3, 033003 [arXiv:1608.01618]

Why neutrino mass & oscillation?

 Higgs boson for electroweak symmetry breaking & mass.

 Chiral symmetry breaking for mass.

 **The world seems not affected by the tiny neutrino mass!**

 Neutrino mass $\Rightarrow$ Mixing

 3 Neutrino $\Rightarrow$ possible **CP violation**

 CP violation $\Rightarrow$ Leptogenesis

 Leptogenesis $\Rightarrow$ **Matter-Antimatter Asymmetry**

 There is something left in the Universe.

 Baryogenesis from quark mixing is not enough.

ν Oscillation Data

(for NH)	-1σ	Best Value	$+1\sigma$
$\Delta m_s^2 \equiv \Delta m_{12}^2$ (10^{-5}eV^2)	7.33	7.50	7.69
$ \Delta m_a^2 \equiv \Delta m_{13}^2 $ (10^{-3}eV^2)	2.484	2.524	2.563
$\sin^2 \theta_s$ ($\theta_s \equiv \theta_{12}$)	0.294 (32.81°)	0.306 (33.56°)	0.318 (34.33°)
$\sin^2 \theta_a$ ($\theta_a \equiv \theta_{23}$)	0.4200 (40.4°)	0.441 (41.6°)	0.468 (43.1°)
$\sin^2 \theta_r$ ($\theta_r \equiv \theta_{13}$)	0.02091 (8.41°)	0.02166 (8.46°)	0.02241 (8.61°)
δ_D, δ_{Mi}	?, ??	?, ??	?, ??

Esteban, Gonzalez-Garcia, Maltoni, Martinez-Soler & Schwetz, arXiv:1611.01514

Evidence of μ - τ Symmetry



Two small deviations (1σ level):

$$-3.5^\circ < \theta_a - 45^\circ < 5.8^\circ \quad 8.4^\circ < \theta_r < 9.2^\circ$$

with **Best Fit Value**: $\theta_a - 45^\circ = -3.9^\circ$ & $\theta_r = 8.8^\circ$.



Zeroth Order Approximation:

$$\theta_a \approx 45^\circ, \quad \theta_r \approx 0^\circ.$$

$\Rightarrow$ **CP** & μ - τ **Symmetric** Mass Matrix:

$$M_\nu^{(0)} = \begin{pmatrix} A & \mathbf{B} & \mathbf{B} \\ & \mathbf{C} & D \\ & & \mathbf{C} \end{pmatrix}$$

Mohapatra & Nussinov [hep-ph/9809415], Lam [hep-ph/0104116]

Mass Matrix M_ν invariant under Transformation:

$$G_\nu^T M_\nu G_\nu = M_\nu$$

Diagonalization:

$$V_\nu^T M_\nu V_\nu = D_\nu$$

Rephasing:

$$D_\nu = d_\nu^T D_\nu d_\nu$$

with $d_\nu^2 = I_3 \Rightarrow d_\nu = \text{diag}(\pm, \pm, \pm)$.

Together

$$\begin{aligned} M_\nu &= G_\nu^T M_\nu G_\nu = \underline{G_\nu^T V_\nu^* D_\nu V_\nu^\dagger G_\nu} \\ &= \underline{V_\nu^* D_\nu V_\nu^\dagger} = \underline{V_\nu^* d_\nu^T D_\nu d_\nu V_\nu^\dagger} \end{aligned}$$

Consequence: $V_\nu^\dagger G_\nu = d_\nu V_\nu^\dagger \Leftrightarrow \boxed{G_\nu = V_\nu d_\nu V_\nu^\dagger}$

For Leptons: $\underline{F_\ell = V_\ell d_\ell V_\ell^\dagger}$ with $d_\ell = \text{diag}(e^{i\beta_1}, e^{i\beta_2}, e^{i\beta_3})$.

Two Nontrivial Independent possibilities of $\mathbf{d}_\nu$:

$$\mathbf{d}_\nu^{(1)} = \text{diag}(-1, 1, 1), \quad \mathbf{d}_\nu^{(2)} = \text{diag}(1, -1, 1), \quad \mathbf{d}_\nu^{(3)} = -d_\nu^{(1)} d_\nu^{(2)}.$$

θ_s parameterized in terms of $\mathbf{k}$: $\tan \theta_s = \sqrt{2}/k$

$$V_\nu(k) = \begin{pmatrix} \frac{k}{\sqrt{2+k^2}} & \frac{\sqrt{2}}{\sqrt{2+k^2}} & 0 \\ \frac{1}{\sqrt{2+k^2}} & \frac{\sqrt{2(2+k^2)}}{\sqrt{2+k^2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2+k^2}} & \frac{k}{\sqrt{2(2+k^2)}} & \frac{1}{\sqrt{2}} \end{pmatrix} \quad \begin{array}{ll} \mathbf{k} = 2 & \theta_s = 35.3^\circ \text{ [TBM]} \\ \mathbf{k} = \frac{3}{\sqrt{2}} & \theta_s = 33.7^\circ \\ \mathbf{k} = \sqrt{6} & \theta_s = 30.0^\circ \end{array}$$

Two Independent Symmetry Transformations $\mathbf{G}_i = \mathbf{V}_\nu \mathbf{d}_\nu^{(i)} \mathbf{V}_\nu^\dagger$

$$G_1 = \frac{1}{2+k^2} \begin{pmatrix} 2-k^2 & 2k & 2k \\ & k^2 & -2 \\ & & k^2 \end{pmatrix}, \quad G_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\mathbb{Z}_2^S (\times \overline{\mathbb{Z}}_2^S) \times \mathbb{Z}_2^{\mu\tau} \equiv \mathcal{G} = \{\mathbf{E}, \mathbf{G}_1, \mathbf{G}_2 (\equiv G_1 G_3), \mathbf{G}_3\}$$

Full Symmetries:

$\mathcal{H} \equiv \mathcal{G} \times \mathcal{F}$	$\mathcal{G}$	$\mathcal{F}$
S_4	$\mathbb{Z}_2^S \times \mathbb{Z}_2^{\mu\tau}$	$\mathbb{Z}_3 \equiv \{I, F, F^2\}$
$\{G_1, G_3, F\}$	$G_1(G_2), G_3$	$F \equiv \text{diag}(1, \omega, \omega^2)$

Bottom-Up $\uparrow$

$\downarrow$ Top-Down

See also Smirnov et. al., 1204.0445, 1212.2149, 1510.00344

Residual Symmetries:

$$\nu_i: \quad \mathcal{G} \equiv \mathbb{Z}_2^S(\overline{\mathbb{Z}}_2^S) \times \mathbb{Z}_2^{\mu\tau} \quad \text{for} \quad d_\nu^i = \text{diag}(\pm 1, \pm 1, \pm 1)$$

$$\ell_i: \quad \mathcal{F} \in U(1) \times U(1) \quad \text{for} \quad d_\ell^i = \text{diag}(e^{i\beta_1}, e^{i\beta_2}, e^{i\beta_3})$$

Residual Symmetry as Effective Theory



Full symmetry **HAS TO** be **Broken!**



Fermion needs to acquire mass.



Non-trivial mixing

$$V_{\text{PMNS}} = V_{\ell}^{\dagger} V_{\nu}$$



If mixing is **TRUELY** determined by symmetry, it has to be residual symmetry



VEVs



Yukawa couplings



Residual Symmetry as **Custodial Symmetry**



Gauge symmetry has to be broken. Otherwise, no mixing.



Weak mixing angle is a function of gauge couplings, which **cannot be dictated by gauge symmetry** (and VEV).



Weak mixing angle is related to the physical observables, the **gauge boson masses**, by **custodial symmetry**.

Lepton's Representation:

$$\begin{pmatrix} e_L \\ \mu_L \\ \tau_L \end{pmatrix} \sim \mathbf{3}, \quad \begin{matrix} e_R \sim \mathbf{1} \\ \mu_R \sim \mathbf{1}' \\ \tau_R \sim \mathbf{1}'' \end{matrix}, \quad \begin{pmatrix} \varphi_1 \\ \varphi_2 \\ \varphi_3 \end{pmatrix} \sim \mathbf{3}.$$

A_4 invariant Lagrangian:

$$\begin{aligned} \mathcal{L}_\ell &= \mathbf{y}_1 \bar{e}_R (\mathbf{1} \varphi_1^\dagger e_L + \mathbf{1} \varphi_2^\dagger \tau_L + \mathbf{1} \varphi_3^\dagger \tau_L) \\ &+ \mathbf{y}_2 \bar{\mu}_R (\omega \varphi_1^\dagger e_L + \mathbf{1} \varphi_2^\dagger \tau_L + \omega^2 \varphi_3^\dagger \tau_L) \\ &+ \mathbf{y}_3 \bar{\tau}_R (\omega^2 \varphi_1^\dagger e_L + \mathbf{1} \varphi_2^\dagger \tau_L + \omega \varphi_3^\dagger \tau_L). \end{aligned}$$

Mass term with $\langle \varphi_i \rangle = v_i$:

$$\mathcal{L}_\ell = \begin{pmatrix} \bar{e}_R & \bar{\mu}_R & \bar{\tau}_R \end{pmatrix} \begin{pmatrix} \mathbf{y}_1 & & \\ & \mathbf{y}_2 & \\ & & \mathbf{y}_3 \end{pmatrix} \begin{pmatrix} \mathbf{1} & \mathbf{1} & \mathbf{1} \\ \omega & \mathbf{1} & \omega^2 \\ \omega^2 & \mathbf{1} & \omega \end{pmatrix} \begin{pmatrix} \mathbf{v}_1 & & \\ & \mathbf{v}_2 & \\ & & \mathbf{v}_3 \end{pmatrix} \begin{pmatrix} e_L \\ \mu_L \\ \tau_L \end{pmatrix}.$$


$$\mathbf{v}_1 = \mathbf{v}_2 = \mathbf{v}_3 = \mathbf{v} \Rightarrow U_{\ell,R} = I, \mathbf{U}_{\ell,L}(\omega), m_{\ell,i} = \mathbf{y}_i \mathbf{v}.$$

$$\mathbf{y}_1 = \mathbf{y}_2 = \mathbf{y}_3 = \mathbf{y} \Rightarrow U_{\ell,L} = I, \mathbf{U}_{\ell,R}(\omega), m_{\ell,i} = \mathbf{y} \mathbf{v}_i.$$

Prediction of Large δ_D by $\mathbb{Z}_2^s$ and $\overline{\mathbb{Z}}_2^s$

$$\cos \delta_D = \frac{(s_s^2 - c_s^2 s_r^2)(c_a^2 - s_a^2)}{4c_a s_a c_s s_s s_r}$$

$$\cos \delta_D = \frac{(s_s^2 s_r^2 - c_s^2)(c_a^2 - s_a^2)}{4c_a s_a c_s s_s s_r}$$




1σ Indication for $\delta_D = -74^\circ (-110^\circ)$ [Schwetz et.al. 1108.1376]

Dirac CP Phase Measurement

CP Measurement @ Accelerator Exps

LPSC
LEIPZIG
2017-2018

T2K



LPSC
LEIPZIG
2017-2018

$\text{NO}\nu\text{A}$



LPSC
LEIPZIG
2017-2018

DUNE, T2KII/T2HK/T2KK/T2KO, MOMENT/ADS-CI, Super-PINGU

The Dirac CP Phase δ_D @ Accelerator Exp

LPSC To leading order in $\alpha = \frac{\delta M_{21}^2}{|\delta M_{31}^2|} \sim 3\%$, the oscillation probability relevant to measuring δ_D @ T2(H)K,

$$P_{\nu_\mu \rightarrow \nu_e} \approx 4s_a^2 c_r^2 s_r^2 \sin^2 \phi_{31} - 8c_a s_a c_r^2 s_r c_s s_s \sin \phi_{21} \sin \phi_{31} [\cos \delta_D \cos \phi_{31} \pm \sin \delta_D \sin \phi_{31}]$$

$$P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e}$$

for ν & $\bar{\nu}$, respectively. $[\phi_{ij} \equiv \frac{\delta m_{ij}^2 L}{4E_\nu}]$

LPSC $\nu_\mu \rightarrow \nu_\mu$ Exps measure $\sin^2(2\theta_a)$ precisely, but not $\sin^2 \theta_a$.

LPSC Run both ν & $\bar{\nu}$ modes @ first peak $[\phi_{31} = \frac{\pi}{2}, \phi_{21} = \alpha \frac{\pi}{2}]$,

$$P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e} + P_{\nu_\mu \rightarrow \nu_e} = 2s_a^2 c_r^2 s_r^2,$$

$$P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e} - P_{\nu_\mu \rightarrow \nu_e} = \alpha \pi \sin(2\theta_s) \sin(2\theta_r) \sin(2\theta_a) \cos \theta_r \sin \delta_D.$$

The Dirac CP Phase δ_D @ Accelerator Exp

Accelerator experiment, such as **T2(H)K**, uses off-axis beam to compare ν_e & $\bar{\nu}_e$ appearance @ the oscillation maximum.



Disadvantages:



Efficiency:



Proton accelerators produce ν more efficiently than $\bar{\nu}$ ($\sigma_\nu > \sigma_{\bar{\nu}}$).



The $\bar{\nu}$ mode needs more beam time [**$T_{\bar{\nu}} : T_\nu = 2 : 1$**].



Undercut statistics $\Rightarrow$ Difficult to reduce the uncertainty.



Degeneracy:



Only **sin δ_D** appears in $P_{\nu_\mu \rightarrow \nu_e}$ & $P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e}$.



Cannot distinguish **δ_D** from **$\pi - \delta_D$** .



CP Uncertainty $\frac{\partial P_{\mu e}}{\partial \delta_D} \propto \cos \delta_D \Rightarrow \Delta(\delta_D) \propto \mathbf{1 / \cos \delta_D}$.



Solution:

Measure $\bar{\nu}$ mode with μ^+ decay @ rest (μ DAR)

μ DAR $\bar{\nu}$ Oscillation Experiments


LPSC
A cyclotron produces 800 MeV proton beam @ fixed target.

LPSC
Produce $\pi^\pm$ which stops &

LPSC
 π^- is absorbed,

LPSC
 π^+ decays @ rest: $\pi^+ \rightarrow \mu^+ + \nu_\mu$.

LPSC
 μ^+ stops & decays @ rest: $\mu^+ \rightarrow e^+ + \bar{\nu}_\mu + \nu_e$.



LPSC
 $\bar{\nu}_\mu$ travel in all directions, oscillating as they go.

LPSC
A detector measures the $\bar{\nu}_e$ from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillation.

Accelerator + μ DAR Experiments

Combining $\nu_\mu \rightarrow \nu_e$ @ accelerator [narrow peak @ 550 MeV] & $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ @ μ DAR [wide peak ~ 45 MeV] solves the 2 problems:



Efficiency:



$\bar{\nu}$ @ high intensity, μ DAR is plentiful enough.



Accelerator Exps can devote all run time to the ν mode. With same run time, the statistical uncertainty drops by $\sqrt{3}$.



Degeneracy: (decomposition in propagation basis [1309.3176])



New Proposals

1 μ DAR source + 2 detectors

Advantages:

-  Full (100%) duty factor!
-  **Lower** intensity: $\sim 9\text{mA}$ [$\sim 4\times$ lower than DAE δ ALUS]
-  Not far beyond the current state-of-art technology of cyclotron [2.2mA @ Paul Scherrer Institute]
-  MUCH **cheaper** & technically **easier**.
 -  Only one cyclotron.
 -  Lower intensity.


Disadvantage?

-  A second detector!
 -  μ DAR with Two Scintillators (μ DARTS) [1401.3977]
 -  Tokai 'N Toyama to(2) Kamioka (TNT2K) [1506.05023]

TNT2K

LPSC
LEPSC
GRENoble

$$T2(H)K + \mu SK + \mu HK$$





LPSC
LEPSC
GRENoble

μ DAR is also useful for **material**, **medicine** industries in Toyama

Event Shape @ TNT2K

Evslin, Ge & Hagiwara [1506.05023]




Expected μDAR IBD signal from 6 yrs of running @ SK (15km) & HK (23km) with NH.

Simulated by [NuPro](http://nupro.hepforge.org/), <http://nupro.hepforge.org/>

δ_D Precision @ TNT2K


Evslin, Ge & Hagiwara [1506.05023]



Simulated by [NuPro](http://nupro.hepforge.org/), <http://nupro.hepforge.org/>

δ_D Precision @ TNT2K


Evslin, Ge & Hagiwara [1506.05023]




Simulated by [NuPro](http://nupro.hepforge.org/), <http://nupro.hepforge.org/>

Majorana CP Phase Measurement

Preference of NH $\Rightarrow$ Non-Observation of $0\nu 2\beta$?



Any chance of obtaining some information?



$$\langle m \rangle_{ee} \equiv \vec{L}_1 + \vec{L}_2 + \vec{L}_3,$$

with

$$\vec{L}_1 \equiv m_1 U_{e1}^2 = m_1 c_r^2 c_s^2 e^{i\delta_{M1}},$$

$$\vec{L}_2 \equiv m_2 U_{e2}^2 = \sqrt{m_1^2 + \Delta m_s^2} c_r^2 s_s^2,$$



$$\vec{L}_3 \equiv m_3 U_{e3}^2 = \sqrt{m_1^2 + \Delta m_a^2} s_r^2 e^{i\delta_{M3}}.$$

Determine 2 Majorana Phases Simultaneously





$$|L_1 - L_3| \leq L_2 \leq L_1 + L_3.$$

$$\cos \delta_{M1} = -\frac{L_1^2 + L_2^2 - L_3^2}{2L_1L_2} = -\frac{m_1^2 c_r^4 c_s^4 + m_2^2 c_r^4 s_s^4 - m_3^2 s_r^4}{2m_1 m_2 c_r^4 c_s^2 s_s^2},$$


$$\cos \delta_{M3} = +\frac{L_1^2 - L_2^2 - L_3^2}{2L_2L_3} = +\frac{m_1^2 c_r^4 c_s^4 - m_2^2 c_r^4 s_s^4 - m_3^2 s_r^4}{2m_2 m_3 c_r^2 s_r^2 s_s^2}.$$




Uncertainties from Oscillation Parameters






Uncertainties from Oscillation Parameters



Uncertainties from Oscillation Parameters



Majorana Pyramid



Prey of Leptonic CP Phases



Dirac

Majorana 1

Majorana 2

Majorana Pyramid

Thank You!