# Stem cells long term preservation

M. Chapelier (LSM), **F. Piquemal (LP2i Bordeaux**), G. Warot (LSM), M.Zampaolo (LSM) F. Chretien, P. Rocheteau (Pasteur Institute)

Interdisciplinary project between : Physics – Biologie - Medicine

Funded by CNRS – MITI (FAIDORA Program) and Pasteur Institute





## The genesis of the project

View from physicist:

Stem cells are stored in liquid nytrogen for tens of years

Stem cells are inert (no chemical activity)

Integrity of stem cells exposed to natural radioactivity including cosmic-rays for tens of years ?

After several years, meeting with Fabrice Chretien Professor and Pierre Rocheteau biologist at Pasteur Institute

Interested to collaborate to make a first experiment with mouse muscle stem cell


Support of MITI (Mission interdisciplinaire CNRS) – FAIDORA Program (30 k€ over 2 years)







### Radiation effects

### It is well know since long time that radiation can lead to DNA double strand break or ROS (Reactive Oxygen Species)



The DNA of a living cell is repaired several thousand times a day due to chemical effects and natural radiation.












### Radiation effects










## Why the preservation of stem cells is important

By 2050, 2/3 of diseases treated by stem cells







## Results of the studies with mouse stem cells

The objectives were to measure Double Strand Breaks of stem cells from mouse muscles in different conditions of natural radiation exposure and to do some in vivo tests. We used 3 types of Stem Cells : Mesenchymal SC and Hematopoietic SC

Co	ndition	Depth (meter of water equivalent)	Cosmic ray dose (mSv/ year)	Total dose (mSv/year)	Neutrons flux (neutron/ cm <sup>2</sup> .s)
I	Standard Pasteur Institute 10		0.04	0.52–1.77 (calculated)	3.10-3
Ш	LSM surface + sa	nd <mark>0</mark>	1.17	76.65 (measured)	1.10 <sup>-2</sup>
Ш	LSM Undergrour	nd <b>4,800</b>	<2.7 $ imes$ 10 <sup>-6</sup>	$<2.7 \times 10^{-6}$ (calculated)	3.10-6
IV	LSM surface	0	1.17	1.65-2.91 (measured)	1.10-2


Condition II: Increase exposure to CR flux and gamma rays by surrounding cells with sand (Thorium) corresponding to a daily dose of 0.21 mSv/day

Condition III : cells placed underground are shielded by lead







### Experimental setup



Underground Cryostat

### Condition II





Radiopurity measurements of all materials





Cryopreserved Stem Cells Incur Damages Due To Terrestrial Cosmic Rays Impairing Their Integrity Upon Long-Term Storage Cell Transplantation Volume 31: 1–15 (0) The Author(s) 2002 Article rease guidelines: mergaph.com/journale-permissions DOI: 10.1177/09434897211070239 Journals.megoluc.com/home/cll SAGE

P. Rocheteau<sup>1</sup>, G. Warot<sup>2</sup>, M. Chapellier<sup>2</sup>, M. Zampaolo<sup>2</sup>, F. Chretien<sup>1</sup>, and F. Piquemal<sup>3</sup>

### Abstract

Stem cells have the capacity to ensure the renewal of tissues and organs. They could be used in the future for a wide range of therapeutic purposes and are preserved at liquid nitrogen temperature to prevent any chemical or biological activity up to several decades before their use. We show that the cryogenized cells accumulate damages coming from natural radiations, potentially inducing DNA double-strand breaks (DSBs). Such DNA damage in stem cells could lead to either mortality of the cells upon thawing or a mutation diminishing the therapeutic potential of the treatment. Many studies show how seem cells react to different levels of radiation; the effect of terrestrial cosmic rays being key, it is thus also important to investigate the effect of the natural radiation on the cryogenserved stem cell behavior over time. Our study showed that the cryostored stem cells totally shielded from cosmic rays had less DSBs upon long-term storage. This could have important implications on the long-term cryostorage strategy and quality control of different cell banks.

### Keywords

stem cell biology, radiation, mesenchymal stem cells, hematopoietic stem cells, muscle regeneration

### Introduction

The long-term preservation of stem cells is a major challenge for therapeutic applications in the future. However, some studies have shown a decrease in cell quality or viability correlating with the time of cryostorage<sup>1-4</sup>. In the connext of this study, we want to determine whether the cryoconservation of sem cells can lead to an accumulation of low radioactive doses (in the order of mGy) coming from the natural radioactive environment, leading to this decrease in cell quality. Indeed, at the surface of the earth, any biological sample is continuously exposed to radiations coming from telluric radioactivity (TR), radon radioactivity, and terrestrial cosmic trays (TCE).

The TR is defined as the radioactivity of natural radioisotopes coming from potassium, uranium, or thorium decay chains. TR (including radon) doses will depend on the location and the composition of the surrounding materials.

Terrestrial cosmic radiations correspond to particles produced in the earth atmosphere by primary cosmic rays (CR) continuously bombarding the earth atmosphere. These primary CR take their origin from astrophysical processes and are mainly high-energy ions (protons, helium), electrons, and photons. The primary CR interact in altitude with the atoms in the air, leading to the creation of a large number of secondary particles. The secondary particles reaching the earth's surface constitute the TCR flux. They consist of various particles (pions, muons, protons, neutrons, electrons) with a wide range of energy. The secondary particle flux will depend on the location on the earth and solar activity<sup>2</sup>. The only possibility to decrease the entire TCR flux is to protect

<sup>1</sup> Human Hittopathology and Animal Models, Department of Infaction & Epidemiology, Institut, Fataure, Park, France <sup>2</sup> Laboratorie de Physique Subatomique et Corpusculaire, UMR 5821, Univenité Granobie Apez, Centre National de la Racharche Scientifique, Granobie Institute of Technology (Institute of Englissering University Granobie Institute of Technology (Institute of Englissering University Granobie Apez, LPSC-IN2P3, Granobie, France <sup>1</sup> Centre d'Eudee Navelaires de Sordaus Grangian, UMR 5379, Centree National de la Racharche Scientifique and Université de Bordeaux, Grandgran, France

Submitted: May 10, 2021. Revised: December 6, 2021. Accepted: December 7, 2021.


Corresponding Author: F. Reusmal, Contre d'Eutode Nucklaires de Bordeaux Gradignan, UMR 5797, Contre National de la Racherche Scientifique and Université de Bordeaux, F-33170 Gradignan, France. Emait Pajeumail@contbji/10/31/r


Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (http://creativecommons.org/licensex/by-no/4.0) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (http://ut.gapub.com/an.utham/open access=14.ge).


### Published in Cell Transplantation Journal






















e Percentage of cells displaying at least one foci



















Quality of the muscle repair seems better after nngraftment with stem cells protected from radiations







Condition II : CR (neutrons) + Gamma

Condition III : Underground

Condition IV : CR (Neutrons)

Double strand Break seems to be related to CR





## Outlook

- > Still a lot of work to do to
- Long term preservation of biological materials is an issue for many applications : Fundamental research, Medical research, illness treatment, fertility, biodiversity,...
- European patent : WO2019077048 HIGH-FIDELITY LONG-TERM STORAGE OF BIOLOGICAL MATERIAL
- ➤ US patent in progress
- Startup PROCELTECH has been created in particular for fund raising to continue research work
- > For Pasteur Institute, we anticipate a public health problem if treatment with stem cells are used in the future
- > More studies with physicians and biologists in progress : preservation of gametes, seeds,....



