

Quantum Information at Colliders

Kazuki Sakurai
(University of Warsaw)




2024/10/21 LSPC Grenoble

Entanglement and other quantum properties are crucial in:


- developing **quantum technology/devices**
- understanding **QFT** and quantum **gravity**



IBM Q system




Spin




$$F_z \propto \partial_z (\vec{s} \cdot \vec{B}) \propto s_z B'_z$$

- **Z-component** of spin has only two possible values: **+1** and **-1**



- If spin is in the **Z**-direction, **X**, **Y**-component should be zero! (?)




- If spin is in the **Z**-direction, **X**-component should be zero! (?)




- If spin is in the **Z**-direction, **X**-component should be zero! (?)



- If spin is in the **Z**-direction, **X**-component should be zero! (?)



Is there a **definite** s_z **value** of **this particle** before the s_z measurement?




Is probabilistic outcomes in experiments **fundamental** or **due to our ignorance** about the physical state/system?

Is probabilistic outcomes in experiments **fundamental** or **due to our ignorance** about the physical state/system?

Can't be fundamental!

God does not play dice!




Albert Einstein

Is probabilistic outcomes in experiments **fundamental** or **due to our ignorance** about the physical state/system?

Can't be fundamental!

God does not play dice!

Can be tested experimentally :-)




Albert Einstein




John Bell

Alice



Spin-0 particle




Bob




$1/2$




$1/2$




Alice




Spin-0 particle




Bob



Alice




Spin-0 particle




Bob



50%



50%



Alice

+

+

-

+

-

-

+

+


+

-


+

-


Alice



Spin-0 particle




Bob




Alice	+	+	-	+	-	-	+	+	+	-	+	-
Bob	-	-	+	-	+	+	-	-	-	+	-	+



Alice



Spin-0 particle




Bob




Alice	+	+	-	+	-	-	+	+	+	-	+	-
Bob	-	-	+	-	+	+	-	-	-	+	-	+



Alice



Spin-0 particle




Bob




Alice	+	+	-	+	-	-	+	+	+	-	+	-
Bob	-	-	+	-	+	+	-	-	-	+	-	+




Alice




Spin-0 particle




Bob



Alice	+	+	-	+	-	?	+	+	-	+	-
Bob	-	-	+	-	+	?	-	-	+	-	+




	Alice			Bob		
prob.	z	x	θ	z	x	θ
P_1	+	+	+	-	-	-
P_2	-	+	+	+	-	-
P_3	+	-	+	-	+	-
P_4	+	+	-	-	-	+
P_5	+	-	-	-	+	+
P_6	-	+	-	+	-	+
P_7	-	-	+	+	+	-
P_8	-	-	-	+	+	+




	Alice			Bob		
prob.	z	x	θ	z	x	θ
P_1	+	+	+	-	-	-
P_2	-	+	+	+	-	-
P_3	+	-	+	-	+	-
P_4	+	+	-	-	-	+
P_5	+	-	-	-	+	+
P_6	-	+	-	+	-	+
P_7	-	-	+	+	+	-
P_8	-	-	-	+	+	+

$$P(+z, +x) = ?$$



	Alice			Bob		
prob.	z	x	θ	z	x	θ
P ₁	+	+	+	-	-	-
P ₂	-	+	+	+	-	-
P ₃	+	-	+	-	+	-
P ₄	+	+	-	-	-	+
P ₅	+	-	-	-	+	+
P ₆	-	+	-	+	-	+
P ₇	-	-	+	+	+	-
P ₈	-	-	-	+	+	+


$$P(+z, +x) = P_3 + P_5$$



	Alice			Bob		
prob.	z	x	θ	z	x	θ
P_1	+	+	+	-	-	-
P_2	-	+	+	+	-	-
P_3	+	-	+	-	+	-
P_4	+	+	-	-	-	+
P_5	+	-	-	-	+	+
P_6	-	+	-	+	-	+
P_7	-	-	+	+	+	-
P_8	-	-	-	+	+	+

$$P(+z, +x) = P_3 + P_5$$


$$P(+z, +\theta) = ?$$



	Alice			Bob		
prob.	z	x	θ	z	x	θ
P ₁	+	+	+	-	-	-
P ₂	-	+	+	+	-	-
P ₃	+	-	+	-	+	-
P ₄	+	+	-	-	-	+
P ₅	+	-	-	-	+	+
P ₆	-	+	-	+	-	+
P ₇	-	-	+	+	+	-
P ₈	-	-	-	+	+	+

$$P(+z, +x) = P_3 + P_5$$

$$P(+z, +\theta) = P_4 + P_5$$




	Alice			Bob		
prob.	z	x	θ	z	x	θ
P ₁	+	+	+	-	-	-
P ₂	-	+	+	+	-	-
P ₃	+	-	+	-	+	-
P ₄	+	+	-	-	-	+
P ₅	+	-	-	-	+	+
P ₆	-	+	-	+	-	+
P ₇	-	-	+	+	+	-
P ₈	-	-	-	+	+	+

$$P(+z, +x) = P_3 + P_5$$

$$P(+z, +\theta) = P_4 + P_5$$

$$P(+\theta, +x) = ?$$




	Alice			Bob		
prob.	z	x	θ	z	x	θ
P ₁	+	+	+	-	-	-
P ₂	-	+	+	+	-	-
P ₃	+	-	+	-	+	-
P ₄	+	+	-	-	-	+
P ₅	+	-	-	-	+	+
P ₆	-	+	-	+	-	+
P ₇	-	-	+	+	+	-
P ₈	-	-	-	+	+	+

$$P(+z, +x) = P_3 + P_5$$

$$P(+z, +\theta) = P_4 + P_5$$

$$P(+\theta, +x) = P_3 + P_7$$




	Alice			Bob		
prob.	z	x	θ	z	x	θ
P ₁	+	+	+	-	-	-
P ₂	-	+	+	+	-	-
P ₃	+	-	+	-	+	-
P ₄	+	+	-	-	-	+
P ₅	+	-	-	-	+	+
P ₆	-	+	-	+	-	+
P ₇	-	-	+	+	+	-
P ₈	-	-	-	+	+	+

$$P(+z, +x) = P_3 + P_5$$

$$P(+z, +\theta) = P_4 + P_5$$

$$P(+\theta, +x) = P_3 + P_7$$



	Alice			Bob		
prob.	z	x	θ	z	x	θ
P ₁	+	+	+	-	-	-
P ₂	-	+	+	+	-	-
P ₃	+	-	+	-	+	-
P ₄	+	+	-	-	-	+
P ₅	+	-	-	-	+	+
P ₆	-	+	-	+	-	+
P ₇	-	-	+	+	+	-
P ₈	-	-	-	+	+	+

$$P(+z, +x) = P_3 + P_5$$

$$P(+z, +\theta) = P_4 + P_5$$

$$P(+\theta, +x) = P_3 + P_7$$

Bell inequality


$$P(+z, +\theta) + P(+\theta, +x) \geq P(+z, +x)$$

$$B(\theta) = 4 \times [P(z+, \theta+) + P(\theta+, x+) - P(z+, x+)]$$

Bell inequality

$$B(\theta) \geq 0$$

← for any θ




$$B(\theta) = 4 \times [P(z+, \theta+) + P(\theta+, x+) - P(z+, x+)]$$

Bell inequality

$$B(\theta) \geq 0$$

← for any θ




$$|\Psi^{(0,0)}\rangle = \frac{|+, -\rangle_z - |-, +\rangle_z}{\sqrt{2}} \xrightarrow{\text{QM}} B(\theta) = 1 - \sqrt{2} \cos(\theta - \pi/4)$$

$$B(\theta) = 4 \times [P(z+, \theta+) + P(\theta+, x+) - P(z+, x+)]$$

Bell inequality

$$B(\theta) \geq 0$$

← for any θ



$$|\Psi^{(0,0)}\rangle = \frac{|+, -\rangle_z - |-, +\rangle_z}{\sqrt{2}} \xrightarrow{\text{QM}} B(\theta) = 1 - \sqrt{2} \cos(\theta - \pi/4)$$


$$B(\theta = \pi/4) \simeq -0.414 < 0 !!$$

$$B(\theta) = 4 \times [P(z+, \theta+) + P(\theta+, x+) - P(z+, x+)]$$

Bell inequality


$$B(\theta) \geq 0$$

← for any θ



$$|\Psi^{(0,0)}\rangle = \frac{|+, -\rangle_z - |-, +\rangle_z}{\sqrt{2}}$$

QM



$$B(\theta) = 1 - \sqrt{2} \cos(\theta - \pi/4)$$

$$B(\theta = \pi/4) \simeq -0.414 < 0 !!$$



**Violation of Bell inequality
has been experimentally
confirmed!**



NOBELPRISET I FYSIK 2022
THE NOBEL PRIZE IN PHYSICS 2022



KUNGL.
VETENSKAPS-
AKADEMIEN
THE ROYAL SWEDISH ACADEMY OF SCIENCES



Alain Aspect
Université Paris-Saclay &
École Polytechnique, France



John F. Clauser
J.F. Clauser & Assoc.,
USA



Anton Zeilinger
University of Vienna,
Austria

"för experiment med sammanflätade fotoner som påvisat brott mot Bell-olikheter och banat väg för kvantinformationsvetenskap"

"for experiments with entangled photons, establishing the violation of Bell inequalities and pioneering quantum information science"

#nobelprize



Qubit

- The qubit system has two basis kets: $|0\rangle$ and $|1\rangle$. The general state is


$$|\Psi\rangle = c_0|0\rangle + c_1|1\rangle \quad (c_i \in \mathbb{C}, |c_0|^2 + |c_1|^2 = 1)$$

- The n -qubit system has 2^n basis kets:

$$|\psi_1\psi_2\cdots\psi_n\rangle = |\psi_1\rangle \otimes |\psi_2\rangle \otimes \cdots \otimes |\psi_n\rangle \quad |\psi_i\rangle \in \{|0\rangle, |1\rangle\}$$

Ex.) The general state in the **2-qubit** system:

$$|\Psi\rangle = c_{00}|00\rangle + c_{01}|01\rangle + c_{10}|10\rangle + c_{11}|11\rangle \quad \left(c_{ij} \in \mathbb{C}, \sum_{i,j} |c_{ij}|^2 = 1 \right)$$



Separable or Entangled

Q.) Can we write

$$\begin{aligned} |\Psi\rangle_{AB} &= c_{00}|00\rangle + c_{01}|01\rangle + c_{10}|10\rangle + c_{11}|11\rangle \\ &\stackrel{?}{=} \underbrace{(\alpha_0|0\rangle + \alpha_1|1\rangle)}_{|\Psi\rangle_A} \otimes \underbrace{(\beta_0|0\rangle + \beta_1|1\rangle)}_{|\Psi\rangle_B} \end{aligned}$$

Separable or Entangled

Q.) Can we write


$$\begin{aligned} |\Psi\rangle_{AB} &= c_{00}|00\rangle + c_{01}|01\rangle + c_{10}|10\rangle + c_{11}|11\rangle \\ &\stackrel{?}{=} \underbrace{(\alpha_0|0\rangle + \alpha_1|1\rangle)}_{|\Psi\rangle_A} \otimes \underbrace{(\beta_0|0\rangle + \beta_1|1\rangle)}_{|\Psi\rangle_B} \end{aligned}$$

If the answer is $\left\{ \begin{array}{l} \text{Yes} \rightarrow |\Psi\rangle_{AB} \text{ is } \textit{separable} \\ \text{No} \rightarrow |\Psi\rangle_{AB} \text{ is } \textit{entangled} \end{array} \right.$

$$|\Psi\rangle_{AB} = c_{00}|00\rangle + c_{01}|01\rangle + c_{10}|10\rangle + c_{11}|11\rangle$$

is separable iff: $c_{00}c_{11} - c_{01}c_{10} = 0$

→ Separable states are **rare!**



Mixed States

- We are uncertain about the quantum state and only know it is $|\Psi_i\rangle$ with probability p_i .
($i = 1, \dots, N$)
→ We say the state is *mixed* and describe it by the *density operator*:

$$\rho = \sum_{i=1}^N p_i |\Psi_i\rangle \langle \Psi_i| \quad \left(p_i \geq 0, \sum_i p_i = 1 \right)$$

Mixed States

- We are uncertain about the quantum state and only know it is $|\Psi_i\rangle$ with probability p_i .
($i = 1, \dots, N$)
→ We say the state is *mixed* and describe it by the *density operator* :

$$\rho = \sum_{i=1}^N p_i |\Psi_i\rangle \langle \Psi_i| \quad \left(p_i \geq 0, \sum_i p_i = 1 \right)$$

- The density operators are *Hermitian*, *Tr = 1*, *positive definite* :

$$\rho = \rho^\dagger, \quad \text{Tr} \rho = 1, \quad \rho \geq 0$$

- The expectation value of an observable $\hat{\mathcal{O}}$ is can be computed as:

$$\langle \hat{\mathcal{O}} \rangle = \text{Tr} [\hat{\mathcal{O}} \rho]$$

- The mixed state ρ_{AB} of the *A-B* system is said to be *separable*, if it's possible to write

$$\rho_{AB} = \sum_i q_i \cdot \rho_A^i \otimes \rho_B^i \quad \left(q_i \geq 0, \sum_i q_i = 1 \right)$$

Otherwise, the state is said to be *entangled*.


Entanglement as a Resource

- **Separable states will never be entangled** by Local Operation and Classical Communication (LOCC)
- LOCC introduces an **order** among states

⇒ Entanglement is **defined** s.t. it decreases under non-invertible LOCC:

maximally entangled $\rho_{\max} \xrightarrow{\text{LOCC}} \rho \xrightarrow{\text{LOCC}} \rho_{\text{sep}}$ separable state

$$\frac{|00\rangle \pm |11\rangle}{\sqrt{2}}, \frac{|01\rangle \pm |10\rangle}{\sqrt{2}}$$



ex) $|00\rangle + |11\rangle \xrightarrow{\text{local measurement}} |00\rangle$


Entanglement as a Resource

- **Separable states will never be entangled** by Local Operation and Classical Communication (LOCC)
- LOCC introduces an **order** among states

⇒ Entanglement is **defined** s.t. it decreases under non-invertible LOCC:

maximally entangled $\rho_{\max} \xrightarrow{\text{LOCC}} \rho \xrightarrow{\text{LOCC}} \rho_{\text{sep}}$ separable state

$$\frac{|00\rangle \pm |11\rangle}{\sqrt{2}}, \frac{|01\rangle \pm |10\rangle}{\sqrt{2}}$$



ex) $|00\rangle + |11\rangle \xrightarrow{\text{local measurement}} |00\rangle$

- **Entanglement measures** (monotones), $E(\rho)$
 - monotonically decreases under LOCC
 - $E(\rho) = 0$ if ρ is separable and $E(\rho) > 0$ otherwise
 - $E(\rho) = 1$ if ρ is maximally entangled states

Entanglement Measures

- Entanglement measures are often defined nicely for **pure** states $|\Psi\rangle_{AB}$

Von Neumann Entropy: $S_V(|\Psi\rangle_{AB}) = -\text{Tr}[\rho_A \log_2 \rho_A]$

Linear Entropy: $S_L(|\Psi\rangle_{AB}) = 2[1 - \text{Tr}(\rho_A^2)]$

$$\rho_A = \text{Tr}_B (|\Psi\rangle\langle\Psi|_{AB})$$

Entanglement Measures

- Entanglement measures are often defined nicely for **pure** states $|\Psi\rangle_{AB}$

Von Neumann Entropy: $S_V(|\Psi\rangle_{AB}) = -\text{Tr}[\rho_A \log_2 \rho_A]$

$$\rho_A = \text{Tr}_B (|\Psi\rangle\langle\Psi|_{AB})$$

Linear Entropy: $S_L(|\Psi\rangle_{AB}) = 2[1 - \text{Tr}(\rho_A^2)]$

- Entanglement measures for **mixed** states $\rho_{AB} = \sum_i p_i |\Psi_i\rangle\langle\Psi_i|$

Entanglement of Formation:

$$E_F(\rho_{AB}) = \inf_{p_k, |\Psi_k\rangle} [p_k S_V(\Psi_k)]$$

Concurrence:

$$\mathcal{C}(\rho_{AB}) = \inf_{p_k, |\Psi_k\rangle} [p_k \sqrt{S_L(\Psi_k)}]$$

minimising over all possible decompositions

Entanglement Measures

- Entanglement measures are often defined nicely for **pure** states $|\Psi\rangle_{AB}$

Von Neumann Entropy: $S_V(|\Psi\rangle_{AB}) = -\text{Tr}[\rho_A \log_2 \rho_A]$

$$\rho_A = \text{Tr}_B(|\Psi\rangle\langle\Psi|_{AB})$$

Linear Entropy: $S_L(|\Psi\rangle_{AB}) = 2[1 - \text{Tr}(\rho_A^2)]$

- Entanglement measures for **mixed** states $\rho_{AB} = \sum_i p_i |\Psi_i\rangle\langle\Psi_i|$

Entanglement of Formation:

$$E_F(\rho_{AB}) = \inf_{p_k, |\Psi_k\rangle} [p_k S_V(\Psi_k)]$$

minimising over all possible decompositions

Concurrence:

$$\mathcal{C}(\rho_{AB}) = \inf_{p_k, |\Psi_k\rangle} [p_k \sqrt{S_L(\Psi_k)}]$$

-
- For **2-qubit systems**, the concurrence admits the **analytical** expression: [Wootters '98]


$$\mathcal{C}[\rho] = \max(0, \eta_1 - \eta_2 - \eta_3 - \eta_4) \in [0, 1]$$

$\eta_1 \geq \eta_2 \geq \eta_3 \geq \eta_4$ are eigenvalues of $\sqrt{\rho \tilde{\rho}}$ with $\tilde{\rho} \equiv (\sigma_y \otimes \sigma_y) \rho^* (\sigma_y \otimes \sigma_y)$.

measures axis: $x \in \{\vec{n}\}$

outcome: $a \in \{-1, +1\}$

probability for a : $p_A(a | x)$



$y \in \{\vec{n}\}$

$b \in \{-1, +1\}$


$p_B(b | y)$

Experimental data is described by the *joint distribution* $p(a, b | x, y)$

measures axis: $x \in \{\vec{n}\}$

outcome: $a \in \{-1, +1\}$

probability for a : $p_A(a | x)$



$y \in \{\vec{n}\}$

$b \in \{-1, +1\}$

$p_B(b | y)$

Experimental data is described by the *joint distribution* $p(a, b | x, y)$

Local theories

$$p_L(a, b | x, y) = \sum_{\lambda} q_\lambda \cdot p_A(a | x, \lambda) \cdot p_B(b | y, \lambda)$$


probability for λ

measures axis: $x \in \{\vec{n}\}$

outcome: $a \in \{-1, +1\}$

probability for a : $p_A(a | x)$

$$p_A(a | x, \lambda)$$


$$q_\lambda$$


$$p_B(b | y, \lambda)$$



$y \in \{\vec{n}\}$

$b \in \{-1, +1\}$

$p_B(b | y)$

Experimental data is described by the *joint distribution* $p(a, b | x, y)$

Local theories

$$p_L(a, b | x, y) = \sum_{\lambda} q_\lambda \cdot p_A(a | x, \lambda) \cdot p_B(b | y, \lambda)$$

probability for λ

Quantum Mechanics

$$p_Q(a, b | x, y) = \text{Tr} \left[\rho_{AB} \left(M_{a|x} \otimes M_{b|y} \right) \right]$$

density operator

For projective measurement:


$$M_{a|x} = |a_x\rangle\langle a_x|$$
$$\hat{s}_x |a_x\rangle = a_x |a_x\rangle$$

measures axis: $x \in \{\vec{n}\}$

outcome: $a \in \{-1, +1\}$

probability for a : $p_A(a | x)$

$$p_A(a | x, \lambda)$$


$$q_\lambda$$


$$p_B(b | y, \lambda)$$

$y \in \{\vec{n}\}$

$b \in \{-1, +1\}$

$p_B(b | y)$

Experimental data is described by the *joint distribution* $p(a, b | x, y)$

Local theories

$$p_L(a, b | x, y) = \sum_{\lambda} q_\lambda \cdot p_A(a | x, \lambda) \cdot p_B(b | y, \lambda)$$

probability for λ

Quantum Mechanics

$$p_Q(a, b | x, y) = \text{Tr} \left[\rho_{AB} \left(M_{a|x} \otimes M_{b|y} \right) \right]$$

density operator

For projective measurement:

$$M_{a|x} = |a_x\rangle\langle a_x|$$
$$\hat{s}_x |a_x\rangle = a_x |a_x\rangle$$

For separable quantum states:


$$\rho_{AB} = \sum_{\lambda} q_\lambda \cdot \rho_A^\lambda \otimes \rho_B^\lambda \Rightarrow p_{Q_{\text{sep}}}(a, b | x, y) = \sum_{\lambda} q_\lambda \cdot \text{Tr} [\rho_A^\lambda M_{a|x}] \cdot \text{Tr} [\rho_B^\lambda M_{a|x}]$$

measures axis: $x \in \{\vec{n}\}$

outcome: $a \in \{-1, +1\}$

probability for a : $p_A(a|x)$

$$p_A(a|x, \lambda)$$


$$q_\lambda$$


$$p_B(b|y, \lambda)$$



$y \in \{\vec{n}\}$

$b \in \{-1, +1\}$

$p_B(b|y)$

Experimental data is described by the *joint distribution* $p(a, b|x, y)$

Local theories

$$p_L(a, b|x, y) = \sum_{\lambda} q_\lambda \cdot p_A(a|x, \lambda) \cdot p_B(b|y, \lambda)$$


Quantum Mechanics

$$p_Q(a, b|x, y) = \text{Tr} \left[\overset{\text{density operator}}{\rho_{AB}} \left(M_{a|x} \otimes M_{b|y} \right) \right]$$

For separable quantum states:

Local

$$\rho_{AB} = \sum_{\lambda} q_\lambda \cdot \rho_A^\lambda \otimes \rho_B^\lambda \Rightarrow p_{Q_{\text{sep}}}(a, b|x, y) = \sum_{\lambda} q_\lambda \cdot \text{Tr} \left[\rho_A^\lambda M_{a|x} \right] \cdot \text{Tr} \left[\rho_B^\lambda M_{b|y} \right]$$



$$\text{Quantum} \supset \text{Local} \supset \text{Separable}$$

Local theories

$$p_L(a, b | x, y) = \sum_{\lambda} q_{\lambda} \cdot p_A(a | x, \lambda) \cdot p_B(b | y, \lambda)$$


Quantum Mechanics

$$p_Q(a, b | x, y) = \text{Tr} \left[\overset{\text{density operator}}{\rho_{AB}} \left(M_{a|x} \otimes M_{b|y} \right) \right]$$

For separable quantum states:

Local

$$\rho_{AB} = \sum_{\lambda} q_{\lambda} \cdot \rho_A^{\lambda} \otimes \rho_B^{\lambda} \Rightarrow p_{Q_{\text{sep}}}(a, b | x, y) = \sum_{\lambda} q_{\lambda} \cdot \text{Tr} \left[\rho_A^{\lambda} M_{a|x} \right] \cdot \text{Tr} \left[\rho_B^{\lambda} M_{a|x} \right]$$



Quantum $\supset$ **Local** $\supset$ **Separable**



Bell-nonlocal $\subset$ **Entanglement**

Local theories

$$p_L(a, b | x, y) = \sum_{\lambda} q_{\lambda} \cdot p_A(a | x, \lambda) \cdot p_B(b | y, \lambda)$$

Quantum Mechanics

$$p_Q(a, b | x, y) = \text{Tr} \left[\overset{\text{density operator}}{\rho_{AB}} \left(M_{a|x} \otimes M_{b|y} \right) \right]$$

For separable quantum states:

Local

$$\rho_{AB} = \sum_{\lambda} q_{\lambda} \cdot \rho_A^{\lambda} \otimes \rho_B^{\lambda} \Rightarrow p_{Q_{\text{sep}}}(a, b | x, y) = \sum_{\lambda} q_{\lambda} \cdot \text{Tr} \left[\rho_A^{\lambda} M_{a|x} \right] \cdot \text{Tr} \left[\rho_B^{\lambda} M_{a|x} \right]$$


- Nonlocal states in QM does not violate causality

$$p(a|x, y) \equiv \sum_b p(a, b|x, y)$$

Condition for no causality violation: **No-Signalling** [Cirel'son(1980), Popescu, Rohrlich(1994)]

$$\forall a, b, x, x', y, y' \begin{cases} p(a|x, y) = p(a|x, y') & \text{Alice's dist. is indep. of Bob's choice for meas. axis} \\ p(b|x, y) = p(b|x', y) & \text{Bob's dist. is indep. of Alice's choice for meas. axis} \end{cases}$$

No-Signalling $\supset$ **Quantum** $\supset$ **Local** $\supset$ **Separable**



CHSH Inequalities

- Bell-type inequalities (in general) are the inequalities that separate different types of distributions (No-signalling, Quantum, Local).

- Define the correlator $C_{xy} = \langle A_x B_y \rangle \equiv \sum_{a,b} abp(a, b | x, y)$

- CHSH quantity [Clauser-Horne-Shimony-Holt(1969)]

For $a, b \in \{\pm 1\}$, $x \in \{\mathbf{n}_1, \mathbf{n}_2\}$, $y \in \{\mathbf{e}_1, \mathbf{e}_2\}$

$$S_{\text{CHSH}} \equiv C_{\mathbf{n}_1, \mathbf{e}_1} + C_{\mathbf{n}_1, \mathbf{e}_2} + C_{\mathbf{n}_2, \mathbf{e}_1} - C_{\mathbf{n}_2, \mathbf{e}_2}$$

CHSH Inequalities

- Bell-type inequalities (in general) are the inequalities that separate different types of distributions (**No-signalling**, **Quantum**, **Local**).

- Define the correlator $C_{xy} = \langle A_x B_y \rangle \equiv \sum_{a,b} abp(a, b | x, y)$


- CHSH quantity [Clauser-Horne-Shimony-Holt(1969)]

For $a, b \in \{\pm 1\}$, $x \in \{\mathbf{n}_1, \mathbf{n}_2\}$, $y \in \{\mathbf{e}_1, \mathbf{e}_2\}$


$$S_{\text{CHSH}} \equiv C_{\mathbf{n}_1, \mathbf{e}_1} + C_{\mathbf{n}_1, \mathbf{e}_2} + C_{\mathbf{n}_2, \mathbf{e}_1} - C_{\mathbf{n}_2, \mathbf{e}_2}$$

$$S_{\text{CHSH}} \leq \begin{cases} 2 & \text{Local theories} & [\text{CHSH}(1969)] \\ 2\sqrt{2} & \text{Quantum Mechanics} & [\text{Tsirelson}(1987)] \\ 4 & \text{No-signalling} & [\text{Popescu, Rohrlich}(1994)] \end{cases}$$

No-signalling $\supset$ **Quantum** $\supset$ **Local** $\supset$ **Separable**





High Energy Test of QM




- At very short-distances (high-energies), QM might be modified.
 - sense of locality may change (again)
 - QM might be modified to be married with gravity
- Possible modification of QM?
 - No-signalling theories: $\langle \mathcal{B} \rangle_{\text{NS}} \leq 4$ [Cirel'son (1980), Popescu, Rohrlich (1994)]
 - Non-linear extensions of QM: [Weinberg (1989), Polchinski (1991), D.E.Kaplan, S.Rajendran, (2021)]
- Quantum Information quantities (entanglement, CHSH quantity) are often **sensitive** to **BSM** and **non-perturbative effects**. They might open a **door to “new” physics**.


Entangled pairs at Colliders



Entangled pairs at Colliders




how to measure spin?



Particles with weak decays are their own polarimeters

e.g.) For $\tau^- \rightarrow \pi^- + \nu_\tau$ (τ^- rest frame), the spin of τ^- is measured in the direction of π^- ($\vec{\pi}$) and the outcome is +1.

measurement axis




The diagram shows a blue circle labeled τ^- . A blue arrow labeled $\mathbf{m}$ points from the circle towards the upper right. A black arrow labeled $\mathbf{s}$ points upwards from the circle, with two parallel vertical lines below it indicating a specific spin state.

$$p(+|\mathbf{m}) = |\langle +_{\mathbf{m}} | +_{\mathbf{s}} \rangle|^2$$
$$= \frac{1 + (\mathbf{m} \cdot \mathbf{s})}{2}$$


Particles with weak decays are their own polarimeters

e.g.) For $\tau^- \rightarrow \pi^- + \nu_\tau$ (τ^- rest frame), the spin of τ^- is measured in the direction of π^- ($\vec{\pi}$) and the outcome is +1.

measurement axis



$$p(+|\mathbf{m}) = |\langle +_{\mathbf{m}} | +_s \rangle|^2 = \frac{1 + (\mathbf{m} \cdot \mathbf{s})}{2}$$




$$\frac{d\Gamma}{d\Omega} = \frac{1 + (\vec{\pi} \cdot \mathbf{s})}{2}$$


Particles with weak decays are their own polarimeters

e.g.) For $\tau^- \rightarrow \pi^- + \nu_\tau$ (τ^- rest frame), the spin of τ^- is measured in the direction of π^- ($\vec{\pi}$) and the outcome is +1.

measurement axis



$$p(+|\mathbf{m}) = |\langle +_{\mathbf{m}} | +_s \rangle|^2 = \frac{1 + (\mathbf{m} \cdot \mathbf{s})}{2}$$



$$\frac{d\Gamma}{d\Omega} = \frac{1 + (\vec{\pi} \cdot \mathbf{s})}{2}$$

Weakly decaying particles in the SM:

$$\tau, t, W^\pm, Z^0$$

Particles with weak decays are their own polarimeters

More generally,

$$\frac{d\Gamma}{d\Omega} = \frac{1 + \alpha_x \cdot (\vec{x} \cdot \mathbf{s})}{2}$$

$\alpha_x \in [-1, +1]$: **spin analyzing power**

- tau decay

$\alpha_x = 1$ for ($x = \pi^-$ in $\tau^- \rightarrow \pi^- \nu$)

- top decay

decay product x	α_x
b	$-0.3925(6)$
W^+	$0.3925(6)$
ℓ^+ (from a W^+)	$0.999(1)$
$\bar{d}, \bar{s}$ (from a W^+)	$0.9664(7)$
u, c (from a W^+)	$-0.3167(6)$

Spin correlation:


$$C_{\mathbf{n}, \mathbf{n}'} \equiv \langle (\mathbf{s}_A \cdot \mathbf{n})(\mathbf{s}_B \cdot \mathbf{n}') \rangle = \frac{9}{\alpha_x \alpha_y} \langle (\vec{x} \cdot \mathbf{n})(\vec{y} \cdot \mathbf{n}') \rangle$$

$\mathbf{n}, \mathbf{n}'$: spin measurement axes

$\vec{x}, \vec{y}$: direction of decay products

$$S_{\text{CHSH}} \equiv C_{\mathbf{n}_1, \mathbf{n}'_1} + C_{\mathbf{n}_1, \mathbf{n}'_2} + C_{\mathbf{n}_2, \mathbf{n}'_1} - C_{\mathbf{n}_2, \mathbf{n}'_2} \begin{cases} > 2 & \rightarrow \text{Bell-nonlocal} \\ > 2\sqrt{2} & \rightarrow \text{QM violation} \end{cases}$$



$$D \equiv \text{Tr}[C]/3 < -\frac{1}{3} \rightarrow \text{sufficient cond. for entanglement}$$





Recent activities to look into entanglements, etc. in HEP

❖ Prospects to measure entanglement and Bell-nonlocality @ LHC

- $pp \rightarrow t\bar{t}$ Y. Afik and J. R. M. de Nova '21, '22, M. Fabbrichesi, R. Floreanini, G. Panizzo '21
Z. Dong, D. Gonçalves, K. Kong, A. Navarro '23
- $H \rightarrow WW, ZZ$ A. J. Barr '21, J.A. Aguilar-Saavedra, A. Bernal, J.A. Casas, J.M. Moreno '22,
 $pp \rightarrow WW, ZZ$ A. Bernal, P. Caban, J. Rembieliński '23, M. Fabbrichesi, R. Floreanini, E.
Gabrielli, Luca Marzola '23
- $H \rightarrow \tau^+\tau^-$ (@ e^+e^- colliders) M. Fabbrichesi, R. Floreanini, E. Gabrielli 22, M. Altakach,
P. Lamba, F. Maltoni, K. Mawatari, KS '22, K. Ma, T. Li '23




Entanglement Observation at LHC $pp \rightarrow t\bar{t}$




$$\frac{1}{\sigma} \frac{d\sigma}{d\cos\varphi} = \frac{1}{2} (1 - D \cos\varphi)$$

$$\rightarrow D = -3 \cdot \langle \cos\varphi \rangle = \frac{\text{tr}[\mathbf{C}]}{3} < -\frac{1}{3}$$



(sufficient condition for entanglement)



$$\mathcal{L} = \mathcal{L}_{\text{SM}} - \frac{1}{2}\phi(\partial^2 + M_\phi^2)\phi + c_y \frac{y_t}{\sqrt{2}} \phi \bar{t} (\cos \alpha + i\gamma^5 \sin \alpha) t.$$



$$|\phi\rangle = \frac{|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle}{\sqrt{2}}$$




Effect of BSM $pp \rightarrow t\bar{t}$

$$\mathcal{O}_{tG} = g_S \bar{Q} T_A \tilde{\varphi} \sigma^{\mu\nu} t G_{\mu\nu}^A$$


$$\mathcal{O}_{tq}^8 = \sum_{f=1}^2 (\bar{q}_f \gamma_\mu T_A q_f) (\bar{t} \gamma^\mu T^A t)$$

[Aoude Madge Maltoni Mantani (2022)]



β : top velocity in the $t\bar{t}$ rest frame

[Severi Vryonidou (2023)]



$$\Delta^\pm = \pm(C_{kk} + C_{rr}) - C_{nn} - 1$$

$$H \rightarrow \tau^+ \tau^- @ e^+ e^- \text{ colliders}$$

- ✦ M. Altakach, F. Maltoni, K. Mawatari, P. Lamba, KS, [2211.10513] (PRD)

3-particle entanglement

- ✦ KS, M. Spannowsky [2310.01477] (PRL)

$$H \rightarrow \tau^+ \tau^- @ e^+ e^- \text{ colliders}$$

- ◆ M. Altakach, F. Maltoni, K. Mawatari, P. Lamba, KS, [2211.10513] (PRD)

3-particle entanglement


- ◆ KS, M. Spannowsky [2310.01477] (PRL)

Experimental Challenge in $H \rightarrow \tau^+ \tau^-$

Among weakly decaying particles, τ , t , $W^\pm$, Z^0 , the tau-lepton is special because $m_\tau \ll m_H$

One has to measure the direction of pions at the rest frame of each tau.

→ Reconstruction of the tau rest frames (i.e. neutrino reconstruction) is necessary




Experimental Challenge in $H \rightarrow \tau^+ \tau^-$

Among weakly decaying particles, τ , t , $W^\pm$, Z^0 , **the tau-lepton is special** because $m_\tau \ll m_H$

One has to measure the direction of pions at the rest frame of each tau.


→ **Reconstruction of the tau rest frames (i.e. neutrino reconstruction) is necessary**




$H \rightarrow \tau^+ \tau^-$ @ lepton colliders


- For precise event reconstruction and for much smaller background, we consider lepton colliders.

LHC



ILC






$$(P_H^{\text{reco}})^\mu \equiv P_{e^+e^-}^\mu - P_{Z \rightarrow x\bar{x}}^\mu \quad M_{\text{recoil}}^2 \equiv (P_H^{\text{reco}})^2$$

Event selection: $|M_{\text{recoil}} - 125 \text{ GeV}| < 5 \text{ GeV}$

	ILC	FCC-ee
energy (GeV)	250	240
luminosity (ab^{-1})	3	5
beam resolution e^+ (%)	0.18	0.83×10^{-4}
beam resolution e^- (%)	0.27	0.83×10^{-4}
$\sigma(e^+e^- \rightarrow HZ)$ (fb)	240.1	240.3
# of signal ($\sigma \cdot \text{BR} \cdot L \cdot \epsilon$)	385	663
# of background ($\sigma \cdot \text{BR} \cdot L \cdot \epsilon$)	20	36

$$e^+e^- \rightarrow Z + (Z^*/\gamma^*) \rightarrow f\bar{f} + \tau^+\tau^-$$

- Generate the SM events $(\kappa, \delta) = (1,0)$ with **MadGraph5**.
- **100 pseudo-experiments** to estimate the statistical uncertainties



$$(P_H^{\text{reco}})^\mu \equiv P_{e^+e^-}^\mu - P_{Z \rightarrow x\bar{x}}^\mu \quad M_{\text{recoil}}^2 \equiv (P_H^{\text{reco}})^2$$

Event selection: $|M_{\text{recoil}} - 125 \text{ GeV}| < 5 \text{ GeV}$

$$e^+e^- \rightarrow Z + (Z^*/\gamma^*) \rightarrow f\bar{f} + \tau^+\tau^-$$

	ILC	FCC-ee
energy (GeV)	250	240
luminosity (ab^{-1})	3	5
beam resolution e^+ (%)	0.18	0.83×10^{-4}
beam resolution e^- (%)	0.27	0.83×10^{-4}
$\sigma(e^+e^- \rightarrow HZ)$ (fb)	240.1	240.3
# of signal ($\sigma \cdot \text{BR} \cdot L \cdot \epsilon$)	385	663
# of background ($\sigma \cdot \text{BR} \cdot L \cdot \epsilon$)	20	36

- Generate the SM events $(\kappa, \delta) = (1,0)$ with **MadGraph5**.
- **100 pseudo-experiments** to estimate the statistical uncertainties

- To determine the tau momenta, we have to reconstruct the unobserved neutrino momenta $(p_x^\nu, p_y^\nu, p_z^\nu)$, $(p_x^{\bar{\nu}}, p_y^{\bar{\nu}}, p_z^{\bar{\nu}})$.
- **6** unknowns can be constrained by **2** mass-shell conditions and **4** energy-momentum conservation.

$$m_\tau^2 = (p_{\tau^+})^2 = (p_{\pi^+} + p_{\bar{\nu}})^2$$

$$m_\tau^2 = (p_{\tau^-})^2 = (p_{\pi^-} + p_{\nu})^2$$

$$(p_{ee} - p_Z)^\mu = p_H^\mu = [(p_{\pi^-} + p_{\nu}) + (p_{\pi^+} + p_{\bar{\nu}})]^\mu$$


=> 2-fold solutions.

$$|\Psi_{\tau^+\tau^-}^{\text{SM}}\rangle = \frac{|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle}{\sqrt{2}}$$

$$C_{ij}^{\text{SM}} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix}$$


$$C_{\text{SM}}[\rho] = 1$$

$$S_{\text{CHSH}} = 2\sqrt{2}$$



reproduced very accurately in the simulation

→ we found that false solutions also give the correct correlations! (?)



Effect of momentum mismeasurement

$$E_i^{\text{true}} \rightarrow E_i^{\text{obs}} = (1 + \sigma_E \cdot \omega) \cdot E_i^{\text{true}} \quad \sigma_E = 0.03 \quad (i = \pi^\pm, e^\pm, \mu^\pm, j)$$



 random number drawn from the normal distribution

	ILC	FCC-ee
C_{ij}	$\begin{pmatrix} -0.600 \pm 0.210 & 0.003 \pm 0.125 & 0.020 \pm 0.149 \\ 0.003 \pm 0.125 & -0.494 \pm 0.190 & 0.007 \pm 0.128 \\ 0.048 \pm 0.174 & 0.0007 \pm 0.156 & 0.487 \pm 0.193 \end{pmatrix}$	$\begin{pmatrix} -0.559 \pm 0.143 & -0.010 \pm 0.095 & -0.014 \pm 0.122 \\ -0.010 \pm 0.095 & -0.494 \pm 0.139 & -0.002 \pm 0.111 \\ 0.012 \pm 0.124 & 0.020 \pm 0.105 & 0.434 \pm 0.134 \end{pmatrix}$
$\mathcal{C}[\rho]$	0.030 ± 0.071	0.005 ± 0.023
$S_{\text{CHSH}}/2$	0.769 ± 0.189	0.703 ± 0.134

$$C_{ij}^{\text{SM}} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix} \quad C_{\text{SM}}[\rho] = 1 \quad S_{\text{CHSH}}^{\text{SM}}/2 = \sqrt{2}$$

Momentum smearing spoils the previous good result...

Use impact parameter information



Goal:


$$E_i^{\text{true}} \rightarrow E_i^{\text{obs}} \rightarrow E_i^{\text{true}} \quad (i = \pi^\pm, e^\pm, \mu^\pm, j)$$

What we do:

- modify E_i^{obs} for some amount by δ

$$E_i^{\text{obs}} \rightarrow E_i(\delta_i) = (1 + \delta_i \sigma_E) \cdot E_i^{\text{obs}}$$

Use impact parameter information



Goal:

$$E_i^{\text{true}} \rightarrow E_i^{\text{obs}} \rightarrow E_i^{\text{true}} \quad (i = \pi^\pm, e^\pm, \mu^\pm, j)$$

What we do:

- modify E_i^{obs} for some amount by δ


$$E_i^{\text{obs}} \rightarrow E_i(\delta_i) = (1 + \delta_i \sigma_E) \cdot E_i^{\text{obs}}$$

- solve tau direction $\mathbf{e}_{\tau^\pm}(\delta)$

→ lets us calculate $\vec{b}_\pm$ as functions of δ

$$\vec{b}_\pm^{\text{reco}}(\mathbf{e}_{\tau^\pm}) = |\vec{b}_\pm| \cdot [\mathbf{e}_{\tau^\pm} \cdot \sin^{-1} \Theta_\pm - \mathbf{e}_{\pi^\pm} \cdot \tan^{-1} \Theta_\pm]$$

Use impact parameter information



Goal:

$$E_i^{\text{true}} \rightarrow E_i^{\text{obs}} \rightarrow E_i^{\text{true}} \quad (i = \pi^\pm, e^\pm, \mu^\pm, j)$$

What we do:

- modify E_i^{obs} for some amount by δ

$$E_i^{\text{obs}} \rightarrow E_i(\delta_i) = (1 + \delta_i \sigma_E) \cdot E_i^{\text{obs}}$$

- solve tau direction $\mathbf{e}_{\tau^\pm}(\delta)$

→ lets us calculate $\vec{b}_\pm$ as functions of δ

$$\vec{b}_\pm^{\text{reco}}(\mathbf{e}_{\tau^\pm}) = |\vec{b}_\pm| \cdot [\mathbf{e}_{\tau^\pm} \cdot \sin^{-1} \Theta_\pm - \mathbf{e}_{\pi^\pm} \cdot \tan^{-1} \Theta_\pm]$$

- compare the calculated $\vec{b}_\pm^{\text{reco}}(\delta)$ and measured $\vec{b}_\pm^{\text{obs}}$ and construct the likelihood function

2 fold solutions: $i_s = 1, 2$

$$\vec{\Delta}_{b_\pm}^{i_s}(\delta) \equiv \vec{b}_\pm - \vec{b}_\pm^{\text{reco}}(\mathbf{e}_{\tau^\pm}^{i_s}(\delta))$$

$$L_\pm^{i_s}(\delta) = \frac{[\Delta_{b_\pm}^{i_s}(\delta)]_x^2 + [\Delta_{b_\pm}^{i_s}(\delta)]_y^2}{\sigma_{b_T}^2} + \frac{[\Delta_{b_\pm}^{i_s}(\delta)]_z^2}{\sigma_{b_z}^2} + \delta_{\pi^+}^2 + \delta_{\pi^-}^2 + \delta_x^2 + \delta_{\bar{x}}^2.$$

minimizing $L^{i_s}(\delta)$ would give us the correct set of δ_s and solution i_s

Result

2211.10513

	ILC	FCC-ee
C_{ij}	$\begin{pmatrix} 0.830 \pm 0.176 & 0.020 \pm 0.146 & -0.019 \pm 0.159 \\ -0.034 \pm 0.160 & 0.981 \pm 0.1527 & -0.029 \pm 0.156 \\ -0.001 \pm 0.158 & -0.021 \pm 0.155 & -0.729 \pm 0.140 \end{pmatrix}$	$\begin{pmatrix} 0.925 \pm 0.109 & -0.011 \pm 0.110 & 0.038 \pm 0.095 \\ -0.009 \pm 0.110 & 0.929 \pm 0.113 & 0.001 \pm 0.115 \\ -0.026 \pm 0.122 & -0.019 \pm 0.110 & -0.879 \pm 0.098 \end{pmatrix}$
$\mathcal{C}[\rho]$	0.778 ± 0.126	0.871 ± 0.084
$S_{\text{CHSH}}/2$	1.103 ± 0.163	1.276 ± 0.094

$$C_{ij}^{\text{SM}} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix} \qquad C_{\text{SM}}[\rho] = 1 \qquad S_{\text{CHSH}}^{\text{SM}}/2 = \sqrt{2}$$

Result

2211.10513

	ILC	FCC-ee
C_{ij}	$\begin{pmatrix} 0.830 \pm 0.176 & 0.020 \pm 0.146 & -0.019 \pm 0.159 \\ -0.034 \pm 0.160 & 0.981 \pm 0.1527 & -0.029 \pm 0.156 \\ -0.001 \pm 0.158 & -0.021 \pm 0.155 & -0.729 \pm 0.140 \end{pmatrix}$	$\begin{pmatrix} 0.925 \pm 0.109 & -0.011 \pm 0.110 & 0.038 \pm 0.095 \\ -0.009 \pm 0.110 & 0.929 \pm 0.113 & 0.001 \pm 0.115 \\ -0.026 \pm 0.122 & -0.019 \pm 0.110 & -0.879 \pm 0.098 \end{pmatrix}$
$\mathcal{C}[\rho]$	$0.778 \pm 0.126 \quad \sim 5\sigma$	$0.871 \pm 0.084 \quad \gg 5\sigma$
$S_{\text{CHSH}}/2$	1.103 ± 0.163	$1.276 \pm 0.094 \quad \sim 3\sigma$

$$C_{ij}^{\text{SM}} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix} \quad C_{\text{SM}}[\rho] = 1 \quad S_{\text{CHSH}}^{\text{SM}}/2 = \sqrt{2}$$

Result

2211.10513

	ILC	FCC-ee
C_{ij}	$\begin{pmatrix} 0.830 \pm 0.176 & 0.020 \pm 0.146 & -0.019 \pm 0.159 \\ -0.034 \pm 0.160 & 0.981 \pm 0.1527 & -0.029 \pm 0.156 \\ -0.001 \pm 0.158 & -0.021 \pm 0.155 & -0.729 \pm 0.140 \end{pmatrix}$	$\begin{pmatrix} 0.925 \pm 0.109 & -0.011 \pm 0.110 & 0.038 \pm 0.095 \\ -0.009 \pm 0.110 & 0.929 \pm 0.113 & 0.001 \pm 0.115 \\ -0.026 \pm 0.122 & -0.019 \pm 0.110 & -0.879 \pm 0.098 \end{pmatrix}$
$\mathcal{C}[\rho]$	$0.778 \pm 0.126 \quad \sim 5\sigma$	$0.871 \pm 0.084 \quad \gg 5\sigma$
$S_{\text{CHSH}}/2$	1.103 ± 0.163	$1.276 \pm 0.094 \quad \sim 3\sigma$

$$C_{ij}^{\text{SM}} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix} \quad C_{\text{SM}}[\rho] = 1 \quad S_{\text{CHSH}}^{\text{SM}}/2 = \sqrt{2}$$

Superiority of FCC-ee over ILC is due to a better beam resolution

	ILC	FCC-ee
energy (GeV)	250	240
luminosity (ab ⁻¹)	3	5
beam resolution e^+ (%)	0.18	$0.83 \cdot 10^{-4}$
beam resolution e^- (%)	0.27	$0.83 \cdot 10^{-4}$

CP measurement

- Under CP, the spin correlation matrix transforms: $C \xrightarrow{CP} C^T$
- This can be used for a *model-independent* test of CP violation. We define:

$$A \equiv (C_{rn} - C_{nr})^2 + (C_{nk} - C_{kn})^2 + (C_{kr} - C_{rk})^2 \geq 0$$

- Observation of $A \neq 0$ immediately confirms CP violation.
- From our simulation, we observe

$$A = \begin{cases} 0.204 \pm 0.173 & (\text{ILC}) \\ 0.112 \pm 0.085 & (\text{FCC-ee}) \end{cases} \quad \longleftarrow \text{consistent with absence of CPV}$$

- This model independent bounds can be translated to the constraint on the CP-phase δ

$$\mathcal{L}_{\text{int}} \propto H \bar{\psi}_\tau (\cos \delta + i \gamma_5 \sin \delta) \psi_\tau \quad \longrightarrow \quad C_{ij} = \begin{pmatrix} \cos 2\delta & \sin 2\delta & 0 \\ -\sin 2\delta & \cos 2\delta & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \longrightarrow \quad A(\delta) = 4 \sin^2 2\delta$$

CP measurement

- Focusing on the region near $|\delta| = 0$, we find the 1- σ bounds:

$$|\delta| < \begin{cases} 8.9^\circ & (\text{ILC}) \\ 6.4^\circ & (\text{FCC-ee}) \end{cases}$$


- Other studies:

$$\Delta\delta \sim 11.5^\circ \quad (\text{HL-LHC}) \quad [\text{Hagiwara, Ma, Mori 2016}]$$


$$\Delta\delta \sim 4.3^\circ \quad (\text{ILC}) \quad [\text{Jeans and G. W. Wilson 2018}]$$

3-particle entanglement


$$2 \rightarrow 222$$



$$0 \rightarrow 222$$



$$22 \rightarrow 223$$




[Subba, Rahaman, 2404.03292]

3-particle entanglement


$$2 \rightarrow 222$$

Top decay




[KS, Spannowsky, 2310.01477]

$$0 \rightarrow 222$$



[Morales, 2403.18023]

$$22 \rightarrow 223$$



[Subba, Rahaman, 2404.03292]

Three qubit system: 222

- $2^3 = 8$ basis kets:

$$|\Psi_{ABC}\rangle = c_{000}|000\rangle + c_{001}|001\rangle + c_{010}|010\rangle + \dots$$



- Entanglement among **2-individual** particles:

$$\mathcal{C}_{AB}[|\Psi_{ABC}\rangle] = \mathcal{C}[\rho_{AB}] \quad \rho_{AB} = \text{Tr}_C |\Psi\rangle\langle\Psi|_{ABC}$$

- Entanglement among **one-to-other**:

$$\begin{aligned} |\Psi_{ABC}\rangle = & |0\rangle_A \otimes (c_{000}|00\rangle_{BC} + c_{001}|01\rangle_{BC} + \dots) \\ & + |1\rangle_A \otimes (c_{100}|00\rangle_{BC} + c_{101}|01\rangle_{BC} + \dots) \end{aligned}$$

$$\mathcal{C}_{A(BC)}[|\Psi_{ABC}\rangle] = \sqrt{2[1 - \text{Tr}\rho_{BC}^2]} \quad \rho_{BC} = \text{Tr}_A |\Psi\rangle\langle\Psi|_{ABC}$$



Classification

Fully-separable: $|\phi^{\text{fs}}\rangle_{A|B|C} = |\alpha\rangle_A \otimes |\beta\rangle_B \otimes |\gamma\rangle_C$, $\mathcal{C}_{ij} = \mathcal{C}_{i(jk)} = 0$

Bi-separable: $|\phi^{\text{bs}}\rangle_{A|BC} = |\alpha\rangle_A \otimes |\delta\rangle_{BC}$

$$|\phi^{\text{bs}}\rangle_{B|AC} = |\beta\rangle_B \otimes |\delta\rangle_{AC}$$

$$|\phi^{\text{bs}}\rangle_{C|AB} = |\gamma\rangle_C \otimes |\delta\rangle_{AB}$$

$$\mathcal{C}_{A(BC)} = 0$$

$$\mathcal{C}_{BC}, \mathcal{C}_{B(AC)}, \mathcal{C}_{C(AB)} \neq 0$$

Classification

Fully-separable: $|\phi^{\text{fs}}\rangle_{A|B|C} = |\alpha\rangle_A \otimes |\beta\rangle_B \otimes |\gamma\rangle_C$,

$$\mathcal{C}_{ij} = \mathcal{C}_{i(jk)} = 0$$

Bi-separable: $|\phi^{\text{bs}}\rangle_{A|BC} = |\alpha\rangle_A \otimes |\delta\rangle_{BC}$

$$|\phi^{\text{bs}}\rangle_{B|AC} = |\beta\rangle_B \otimes |\delta\rangle_{AC}$$

$$|\phi^{\text{bs}}\rangle_{C|AB} = |\gamma\rangle_C \otimes |\delta\rangle_{AB}$$

$$\mathcal{C}_{A(BC)} = 0$$

$$\mathcal{C}_{BC}, \mathcal{C}_{B(AC)}, \mathcal{C}_{C(AB)} \neq 0$$

complement




**Genuinely Multipartite
Entangled (GME):**

$$|GHZ_3\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$$

$$|W_3\rangle = \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle)$$

maximally GME



Classification

Fully-separable: $|\phi^{\text{fs}}\rangle_{A|B|C} = |\alpha\rangle_A \otimes |\beta\rangle_B \otimes |\gamma\rangle_C$, $\mathcal{C}_{ij} = \mathcal{C}_{i(jk)} = 0$

Bi-separable: $|\phi^{\text{bs}}\rangle_{A|BC} = |\alpha\rangle_A \otimes |\delta\rangle_{BC}$

$$|\phi^{\text{bs}}\rangle_{B|AC} = |\beta\rangle_B \otimes |\delta\rangle_{AC}$$

$$|\phi^{\text{bs}}\rangle_{C|AB} = |\gamma\rangle_C \otimes |\delta\rangle_{AB}$$

$$\mathcal{C}_{A(BC)} = 0$$

$$\mathcal{C}_{BC}, \mathcal{C}_{B(AC)}, \mathcal{C}_{C(AB)} \neq 0$$

complement



**Genuinely Multipartite
Entangled (GME):**

$$|GHZ_3\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$$

$$|W_3\rangle = \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle)$$


maximally GME

- All GME states can be classified either **GHZ** or **W** classes! [Dur, Vidal, Cirac 2000]

$$|\phi^{\text{GME}}\rangle \longrightarrow \hat{A} \otimes \hat{B} \otimes \hat{C} |\phi^{\text{GME}}\rangle = \begin{cases} |GHZ_3\rangle \\ |W_3\rangle \end{cases}$$

$$\hat{A} \in I(\mathcal{H}_A), \hat{B} \in I(\mathcal{H}_B), \hat{C} \in I(\mathcal{H}_C)$$

$I(\mathcal{H})$: set of intertible operators in $\mathcal{H}$



[Guhne,
Toch 2009]

Monogamy




- All 3-qubit pure states can be transformed by a local unitary to

$$|\psi\rangle = \lambda_0|000\rangle + \lambda_1 e^{i\theta}|100\rangle + \lambda_2|101\rangle + \lambda_3|110\rangle + \lambda_4|111\rangle, \quad \lambda_i \geq 0, \sum_i \lambda_i^2 = 1 \text{ and } \theta \in [0; \pi]$$

- 1-2** and **1-1** entanglements are related to by the **monogamy** relations: [Coffman, Kundu, Wootters '99]

$$\mathcal{C}_{A(BC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{AC}^2 + \tau \quad \tau = 4\lambda_0^2\lambda_4^2 \geq 0 \quad \leftarrow \text{3-tangle}$$



Monogamy



- All 3-qubit pure states can be transformed by a local unitary to


$$|\psi\rangle = \lambda_0|000\rangle + \lambda_1 e^{i\theta}|100\rangle + \lambda_2|101\rangle + \lambda_3|110\rangle + \lambda_4|111\rangle, \quad \lambda_i \geq 0, \sum_i \lambda_i^2 = 1 \text{ and } \theta \in [0; \pi]$$

- 1-2** and **1-1** entanglements are related to by the **monogamy** relations: [Coffman, Kundu, Wootters '99]

$$\mathcal{C}_{A(BC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{AC}^2 + \tau \quad \tau = 4\lambda_0^2\lambda_4^2 \geq 0 \quad \leftarrow \text{3-tangle}$$

$$\mathcal{C}_{B(AC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{BC}^2 + \tau$$

$$\mathcal{C}_{C(AB)}^2 = \mathcal{C}_{AC}^2 + \mathcal{C}_{BC}^2 + \tau$$



Monogamy



- All 3-qubit pure states can be transformed by a local unitary to

$$|\psi\rangle = \lambda_0|000\rangle + \lambda_1 e^{i\theta}|100\rangle + \lambda_2|101\rangle + \lambda_3|110\rangle + \lambda_4|111\rangle, \quad \lambda_i \geq 0, \sum_i \lambda_i^2 = 1 \text{ and } \theta \in [0; \pi]$$


- 1-2** and **1-1** entanglements are related to by the **monogamy** relations: [Coffman, Kundu, Wootters '99]

$$\mathcal{C}_{A(BC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{AC}^2 + \tau$$

$$\tau = 4\lambda_0^2\lambda_4^2 \geq 0 \quad \leftarrow \text{3-tangle}$$

$$\mathcal{C}_{B(AC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{BC}^2 + \tau$$

$$\mathcal{C}_{C(AB)}^2 = \mathcal{C}_{AC}^2 + \mathcal{C}_{BC}^2 + \tau$$



Monogamy



- All 3-qubit pure states can be transformed by a local unitary to


$$|\psi\rangle = \lambda_0|000\rangle + \lambda_1 e^{i\theta}|100\rangle + \lambda_2|101\rangle + \lambda_3|110\rangle + \lambda_4|111\rangle, \quad \lambda_i \geq 0, \sum_i \lambda_i^2 = 1 \text{ and } \theta \in [0; \pi]$$

- 1-2** and **1-1** entanglements are related to by the **monogamy** relations: [Coffman, Kundu, Wootters '99]

$$\mathcal{C}_{A(BC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{AC}^2 + \tau \quad \tau = 4\lambda_0^2\lambda_4^2 \geq 0 \quad \leftarrow \text{3-tangle}$$

$$\mathcal{C}_{B(AC)}^2 = \mathcal{C}_{AB}^2 + \mathcal{C}_{BC}^2 + \tau$$

$$\mathcal{C}_{C(AB)}^2 = \mathcal{C}_{AC}^2 + \mathcal{C}_{BC}^2 + \tau$$




- 1-2 concurrence inequalities

$$\mathcal{C}_{A(BC)}^2 + \mathcal{C}_{B(AC)}^2 \geq \mathcal{C}_{C(AB)}^2$$




$$\mathcal{C}_{A(BC)} + \mathcal{C}_{B(AC)} \geq \mathcal{C}_{C(AB)}$$



GME measure

Genuine Multi-particle Entanglement (GME) measure: [Dur, Vidal, Cirac '00, Ma, Chen, Chen, Spengler, Gabriel, Huber '11, Xie, Eberly '21]




The measure should satisfy:

- (1) vanishes for all fully- and bi-separable states
- (2) positive for all GME states
- (3) non-increasing under LOCC

- The **area** of the “**concurrence triangle**” satisfies (1), (2), (3) !

[Jin, Tao, Gui, Fei, Li-Jost, Qiao (2023)]



$$F_3 \equiv \left[\frac{16}{3} Q (Q - C_{A(BC)}) (Q - C_{B(AC)}) (Q - C_{C(AB)}) \right]^{\frac{1}{2}} \in [0, 1]$$

$$Q \equiv \frac{1}{2} [C_{A(BC)} + C_{B(AC)} + C_{C(AB)}]$$

3-body decay: $\psi_0 \rightarrow \psi_1 \bar{\psi}_2 \psi_3$

[KS, M.Spannowsky 2310.01477]

Assumptions:

- all particles have spin 1/2
- all final particles 1,2,3 are massless

Kinematics:

- rest frame of the initial particle 0
- p_1 is in the z -axis
- decay is in the x - z plane


$$p_1^\mu = p_1(1, 0, 0, 1)$$

$$p_2^\mu = p_2(1, \sin \theta_2, 0, \cos \theta_2)$$

$$p_3^\mu = p_3(1, -\sin \theta_3, 0, \cos \theta_3)$$

$\mathbf{n}(\theta, \phi)$: polarisation of initial spin

$\lambda_1, \lambda_2, \lambda_3 \in (+, -)$: helicities of 1,2,3



3-body decay: $\psi_0 \rightarrow \psi_1 \bar{\psi}_2 \psi_3$

[KS, M.Spannowsky 2310.01477]

Assumptions:

- all particles have spin 1/2
- all final particles 1,2,3 are massless

Kinematics:

- rest frame of the initial particle 0
- p_1 is in the z -axis
- decay is in the x - z plane


$$p_1^\mu = p_1(1, 0, 0, 1)$$

$$p_2^\mu = p_2(1, \sin \theta_2, 0, \cos \theta_2)$$

$$p_3^\mu = p_3(1, -\sin \theta_3, 0, \cos \theta_3)$$

$\mathbf{n}(\theta, \phi)$: polarisation of initial spin

$\lambda_1, \lambda_2, \lambda_3 \in (+, -)$: helicities of 1,2,3



initial state

$$|\mathbf{n}(\theta, \phi)\rangle$$

3-body decay: $\psi_0 \rightarrow \psi_1 \bar{\psi}_2 \psi_3$

[KS, M.Spannowsky 2310.01477]

Assumptions:

- all particles have spin 1/2
- all final particles 1,2,3 are massless

Kinematics:

- rest frame of the initial particle 0
- p_1 is in the z -axis
- decay is in the x - z plane


$$p_1^\mu = p_1(1, 0, 0, 1)$$

$$p_2^\mu = p_2(1, \sin \theta_2, 0, \cos \theta_2)$$

$$p_3^\mu = p_3(1, -\sin \theta_3, 0, \cos \theta_3)$$

$\mathbf{n}(\theta, \phi)$: polarisation of initial spin

$\lambda_1, \lambda_2, \lambda_3 \in (+, -)$: helicities of 1,2,3



initial state

$$\hat{1} = \sum_{\lambda_1, \lambda_2, \lambda_3} |\lambda_1, \lambda_2, \lambda_3\rangle \langle \lambda_1, \lambda_2, \lambda_3|$$

$$|\mathbf{n}(\theta, \phi)\rangle = \sum_{\lambda_1, \lambda_2, \lambda_3} \mathcal{M}_{\lambda_1, \lambda_2, \lambda_3}^{\mathbf{n}} |\lambda_1, \lambda_2, \lambda_3\rangle + \dots$$

amplitude

final state

3-body decay: $\psi_0 \rightarrow \psi_1 \bar{\psi}_2 \psi_3$

[KS, M.Spannowsky 2310.01477]

Assumptions:

- all particles have spin 1/2
- all final particles 1,2,3 are massless

Kinematics:

- rest frame of the initial particle 0
- p_1 is in the z -axis
- decay is in the x - z plane


$$p_1^\mu = p_1(1, 0, 0, 1)$$

$$p_2^\mu = p_2(1, \sin \theta_2, 0, \cos \theta_2)$$

$$p_3^\mu = p_3(1, -\sin \theta_3, 0, \cos \theta_3)$$

$\mathbf{n}(\theta, \phi)$: polarisation of initial spin

$\lambda_1, \lambda_2, \lambda_3 \in (+, -)$: helicities of 1,2,3



initial state

$$|\mathbf{n}(\theta, \phi)\rangle$$

$$\hat{1} = \sum_{\lambda_1, \lambda_2, \lambda_3} |\lambda_1, \lambda_2, \lambda_3\rangle \langle \lambda_1, \lambda_2, \lambda_3|$$

amplitude

$$\mathcal{M}_{\lambda_1, \lambda_2, \lambda_3}^{\mathbf{n}} = \langle \lambda_1, \lambda_2, \lambda_3 | \mathbf{n}(\theta, \phi) \rangle$$

$$\sum_{\lambda_1, \lambda_2, \lambda_3}$$

$$\mathcal{M}_{\lambda_1, \lambda_2, \lambda_3}^{\mathbf{n}} |\lambda_1, \lambda_2, \lambda_3\rangle$$

$$= |\Psi\rangle$$

pure (entangled)
3-spin state

Interaction

- Consider **most general** Lorentz invariant 4-fermion interactions

$$\mathcal{L}_{\text{int}} = (\bar{\psi}_1 \Gamma_A \psi_0)(\bar{\psi}_3 \Gamma_B \psi_2)$$

$$\Gamma_{A/B} \in \{1, \gamma^5, \gamma^\mu, \gamma^\mu \gamma^5, \sigma^{\mu\nu}\}$$

$$\psi_0 \rightarrow \psi_1 \bar{\psi}_2 \psi_3$$

❖ Scalar-type

$$[\bar{\psi}_1 (c_S + i c_A \gamma_5) \psi_0][\bar{\psi}_3 (d_S + i d_A \gamma_5) \psi_2]$$

$$c \equiv c_S + i c_A = e^{i\delta_1}$$

$$d \equiv d_S + i d_A = e^{i\delta_2}$$

❖ Vector-type

$$[\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0][\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2]$$

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$

$$c_L, c_R, d_L, d_R \in \mathbb{R}$$

❖ Tensor-type

$$[\bar{\psi}_1 (c_M + i c_E \gamma_5) \sigma^{\mu\nu} \psi_0][\bar{\psi}_3 (d_M + i d_E \gamma_5) \sigma_{\mu\nu} \psi_2]$$

$$c \equiv c_M + i c_E = e^{i\omega_1}$$

$$d \equiv d_M + i d_E = e^{i\omega_2}$$

Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2]$$

$$c \equiv c_S + ic_A = e^{i\delta_1}$$

$$d \equiv d_S + id_A = e^{i\delta_2}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+-+\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

Scalar


$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2]$$

$$c \equiv c_S + ic_A = e^{i\delta_1}$$

$$d \equiv d_S + id_A = e^{i\delta_2}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+-+\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3




Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2] \quad \begin{aligned} c &\equiv c_S + ic_A = e^{i\delta_1} \\ d &\equiv d_S + id_A = e^{i\delta_2} \end{aligned}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+--\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3

$$= [ce^{i\phi} s_{\frac{\theta}{2}} |-\rangle_1 + c^* c_{\frac{\theta}{2}} |+\rangle_1] \otimes \frac{1}{\sqrt{2}} [d |--\rangle_{23} - d^* |++\rangle_{23}] \quad \text{bi-separable}$$



Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2]$$

$$c \equiv c_S + ic_A = e^{i\delta_1}$$

$$d \equiv d_S + id_A = e^{i\delta_2}$$


$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+--\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3

$$= [ce^{i\phi} s_{\frac{\theta}{2}} |-\rangle_1 + c^* c_{\frac{\theta}{2}} |+\rangle_1] \otimes \frac{1}{\sqrt{2}} [d |--\rangle_{23} - d^* |++\rangle_{23}]$$

bi-separable

$$\Rightarrow F_3 = 0$$



Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2]$$

$$c \equiv c_S + ic_A = e^{i\delta_1}$$

$$d \equiv d_S + id_A = e^{i\delta_2}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+-+\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3

$$= [ce^{i\phi} s_{\frac{\theta}{2}} |-\rangle_1 + c^* c_{\frac{\theta}{2}} |+\rangle_1] \otimes \frac{1}{\sqrt{2}} [d|--\rangle_{23} - d^*|++\rangle_{23}]$$

bi-separable

$$\Rightarrow F_3 = 0$$

♣ **1** is **not entangled** with **2** and **3** in any way: $\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{1(23)} = 0$

Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2]$$

$$c \equiv c_S + ic_A = e^{i\delta_1}$$

$$d \equiv d_S + id_A = e^{i\delta_2}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+-+\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3

$$= [ce^{i\phi} s_{\frac{\theta}{2}} |-\rangle_1 + c^* c_{\frac{\theta}{2}} |+\rangle_1] \otimes \frac{1}{\sqrt{2}} [d |--\rangle_{23} - d^* |++\rangle_{23}]$$

bi-separable

$$\Rightarrow F_3 = 0$$

♣ **1** is **not entangled** with **2** and **3** in any way: $\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{1(23)} = 0$

♣ **2** and **3** are **maximally entangled**: $\mathcal{C}_{23} = 1$

Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2] \quad \begin{aligned} c &\equiv c_S + ic_A = e^{i\delta_1} \\ d &\equiv d_S + id_A = e^{i\delta_2} \end{aligned}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+-+\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3

$$= [ce^{i\phi} s_{\frac{\theta}{2}} |-\rangle_1 + c^* c_{\frac{\theta}{2}} |+\rangle_1] \otimes \frac{1}{\sqrt{2}} [d |--\rangle_{23} - d^* |++\rangle_{23}] \quad \text{bi-separable}$$

$$\Rightarrow F_3 = 0$$

♣ **1** is **not entangled** with **2** and **3** in any way: $\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{1(23)} = 0$

♣ **2** and **3** are **maximally entangled**: $\mathcal{C}_{23} = 1$

♣ Due to **monogamy**, **2** and **3** must be **maximally entangled** with the rest:

$$\begin{array}{cc} \mathcal{C}_{2(13)}^2 = \mathcal{C}_{12}^2 + \mathcal{C}_{23}^2 + \tau & \mathcal{C}_{3(12)}^2 = \mathcal{C}_{13}^2 + \mathcal{C}_{23}^2 + \tau \\ \parallel & \parallel \\ 0 & 1 \end{array}$$

Scalar

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_S + ic_A\gamma_5)\psi_0][\bar{\psi}_3(d_S + id_A\gamma_5)\psi_2] \quad \begin{aligned} c &\equiv c_S + ic_A = e^{i\delta_1} \\ d &\equiv d_S + id_A = e^{i\delta_2} \end{aligned}$$

$$\Rightarrow |\Psi\rangle = \frac{cd}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |---\rangle - \frac{cd^*}{\sqrt{2}} \cdot e^{i\phi} s_{\frac{\theta}{2}} |-++\rangle + \frac{c^*d}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+-+\rangle - \frac{c^*d^*}{\sqrt{2}} \cdot c_{\frac{\theta}{2}} |+++ \rangle$$

independent of final state momenta θ_2, θ_3

$$= [ce^{i\phi} s_{\frac{\theta}{2}} |-\rangle_1 + c^* c_{\frac{\theta}{2}} |+\rangle_1] \otimes \frac{1}{\sqrt{2}} [d |--\rangle_{23} - d^* |++\rangle_{23}] \quad \text{bi-separable}$$

$$\Rightarrow F_3 = 0$$

♣ **1** is **not entangled** with **2** and **3** in any way: $\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{1(23)} = 0$

♣ **2** and **3** are **maximally entangled**: $\mathcal{C}_{23} = 1$

♣ Due to **monogamy**, **2** and **3** must be **maximally entangled** with the rest:

$$\begin{array}{cc} \mathcal{C}_{2(13)}^2 = \mathcal{C}_{12}^2 + \mathcal{C}_{23}^2 + \tau & \mathcal{C}_{3(12)}^2 = \mathcal{C}_{13}^2 + \mathcal{C}_{23}^2 + \tau \\ \parallel & \parallel \\ 0 & 1 \end{array} \Rightarrow \begin{cases} \mathcal{C}_{2(13)} = \mathcal{C}_{3(12)} = 1 \\ \tau = 0 \end{cases}$$

[KS, M.Spannowsky
2310.01477]

Vector

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0][\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2]$$

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$
$$c_L, c_R, d_L, d_R \in \mathbb{R}$$

Vector

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0] [\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2]$$

$$c_L, c_R, d_L, d_R \in \mathbb{R}$$

$$\Rightarrow |\Psi\rangle = M_{LL} | - + - \rangle + M_{LR} | - - + \rangle + M_{RL} | + + - \rangle + M_{RR} | + - + \rangle$$


Vector

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0] [\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2] \quad c_L, c_R, d_L, d_R \in \mathbb{R}$$

$$\Rightarrow |\Psi\rangle = M_{LL} | - + - \rangle + M_{LR} | - - + \rangle + M_{RL} | + + - \rangle + M_{RR} | + - + \rangle$$

$$\begin{aligned} \propto & c_L d_L s \frac{\theta_3}{2} \left[c \frac{\theta}{2} c \frac{\theta_2}{2} + e^{i\phi} s \frac{\theta}{2} s \frac{\theta_2}{2} \right] | - + - \rangle + c_L d_R s \frac{\theta_2}{2} \left[c \frac{\theta}{2} c \frac{\theta_3}{2} + e^{i\phi} s \frac{\theta}{2} s \frac{\theta_3}{2} \right] | - - + \rangle \\ & + c_R d_L s \frac{\theta_2}{2} \left[c \frac{\theta}{2} s \frac{\theta_3}{2} - e^{i\phi} s \frac{\theta}{2} c \frac{\theta_3}{2} \right] | + + - \rangle + c_R d_R s \frac{\theta_3}{2} \left[c \frac{\theta}{2} s \frac{\theta_2}{2} - e^{i\phi} s \frac{\theta}{2} c \frac{\theta_2}{2} \right] | + - + \rangle \end{aligned}$$



Vector

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$


$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0] [\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2] \quad c_L, c_R, d_L, d_R \in \mathbb{R}$$

$$\Rightarrow |\Psi\rangle = M_{LL} | - + - \rangle + M_{LR} | - - + \rangle + M_{RL} | + + - \rangle + M_{RR} | + - + \rangle$$

$$\begin{aligned} \propto & c_L d_L s \frac{\theta_3}{2} \left[c \frac{\theta}{2} c \frac{\theta_2}{2} + e^{i\phi} s \frac{\theta}{2} s \frac{\theta_2}{2} \right] | - + - \rangle + c_L d_R s \frac{\theta_2}{2} \left[c \frac{\theta}{2} c \frac{\theta_3}{2} + e^{i\phi} s \frac{\theta}{2} s \frac{\theta_3}{2} \right] | - - + \rangle \\ & + c_R d_L s \frac{\theta_2}{2} \left[c \frac{\theta}{2} s \frac{\theta_3}{2} - e^{i\phi} s \frac{\theta}{2} c \frac{\theta_3}{2} \right] | + + - \rangle + c_R d_R s \frac{\theta_3}{2} \left[c \frac{\theta}{2} s \frac{\theta_2}{2} - e^{i\phi} s \frac{\theta}{2} c \frac{\theta_2}{2} \right] | + - + \rangle \end{aligned}$$

❖ Individual 2-party entanglement:

$$\mathcal{C}_{12} = \mathcal{C}_{13} = 0, \quad \mathcal{C}_{23} = 2|M_{LL}M_{LR}^* + M_{RL}M_{RR}^*|$$



Vector

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0] [\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2] \quad c_L, c_R, d_L, d_R \in \mathbb{R}$$

$$\Rightarrow |\Psi\rangle = M_{LL} | - + - \rangle + M_{LR} | - - + \rangle + M_{RL} | + + - \rangle + M_{RR} | + - + \rangle$$

$$\begin{aligned} \propto & c_L d_L s_{\frac{\theta_3}{2}} \left[c_{\frac{\theta}{2}} c_{\frac{\theta_2}{2}} + e^{i\phi} s_{\frac{\theta}{2}} s_{\frac{\theta_2}{2}} \right] | - + - \rangle + c_L d_R s_{\frac{\theta_2}{2}} \left[c_{\frac{\theta}{2}} c_{\frac{\theta_3}{2}} + e^{i\phi} s_{\frac{\theta}{2}} s_{\frac{\theta_3}{2}} \right] | - - + \rangle \\ & + c_R d_L s_{\frac{\theta_2}{2}} \left[c_{\frac{\theta}{2}} s_{\frac{\theta_3}{2}} - e^{i\phi} s_{\frac{\theta}{2}} c_{\frac{\theta_3}{2}} \right] | + + - \rangle + c_R d_R s_{\frac{\theta_3}{2}} \left[c_{\frac{\theta}{2}} s_{\frac{\theta_2}{2}} - e^{i\phi} s_{\frac{\theta}{2}} c_{\frac{\theta_2}{2}} \right] | + - + \rangle \end{aligned}$$


❖ Individual 2-party entanglement:

$$\mathcal{C}_{12} = \mathcal{C}_{13} = 0, \quad \mathcal{C}_{23} = 2|M_{LL}M_{LR}^* + M_{RL}M_{RR}^*|$$

❖ one-to-other entanglement:

$$\mathcal{C}_{2(13)} = \mathcal{C}_{3(12)} = 2\sqrt{(|M_{LL}|^2 + |M_{RL}|^2)(|M_{LR}|^2 + |M_{RR}|^2)}$$

$$\mathcal{C}_{1(23)} = 2|M_{RR}M_{LL} - M_{LR}M_{RL}|$$



Vector

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$

$$c_L, c_R, d_L, d_R \in \mathbb{R}$$

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0] [\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2]$$

$$\Rightarrow |\Psi\rangle = M_{LL} | - + - \rangle + M_{LR} | - - + \rangle + M_{RL} | + + - \rangle + M_{RR} | + - + \rangle$$

$$\begin{aligned} \propto & c_L d_L s \frac{\theta_3}{2} \left[c \frac{\theta}{2} c \frac{\theta_2}{2} + e^{i\phi} s \frac{\theta}{2} s \frac{\theta_2}{2} \right] | - + - \rangle + c_L d_R s \frac{\theta_2}{2} \left[c \frac{\theta}{2} c \frac{\theta_3}{2} + e^{i\phi} s \frac{\theta}{2} s \frac{\theta_3}{2} \right] | - - + \rangle \\ & + c_R d_L s \frac{\theta_2}{2} \left[c \frac{\theta}{2} s \frac{\theta_3}{2} - e^{i\phi} s \frac{\theta}{2} c \frac{\theta_3}{2} \right] | + + - \rangle + c_R d_R s \frac{\theta_3}{2} \left[c \frac{\theta}{2} s \frac{\theta_2}{2} - e^{i\phi} s \frac{\theta}{2} c \frac{\theta_2}{2} \right] | + - + \rangle \end{aligned}$$

❖ Individual 2-party entanglement:

$$\mathcal{C}_{12} = \mathcal{C}_{13} = 0, \quad \mathcal{C}_{23} = 2|M_{LL}M_{LR}^* + M_{RL}M_{RR}^*|$$

❖ one-to-other entanglement:


$$\mathcal{C}_{2(13)} = \mathcal{C}_{3(12)} = 2\sqrt{(|M_{LL}|^2 + |M_{RL}|^2)(|M_{LR}|^2 + |M_{RR}|^2)}$$

$$\mathcal{C}_{1(23)} = 2|M_{RR}M_{LL} - M_{LR}M_{RL}|$$

❖ 3-tangle

$$\begin{array}{cc} 0 & 0 \\ \parallel & \parallel \end{array}$$

$$\tau = \mathcal{C}_{1(23)}^2 - [\mathcal{C}_{12}^2 + \mathcal{C}_{13}^2] = \mathcal{C}_{1(23)}^2$$



Vector

$$P_{R/L} = \frac{1 \pm \gamma^5}{2}$$

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1 \gamma_\mu (c_L P_L + c_R P_R) \psi_0] [\bar{\psi}_3 \gamma^\mu (d_L P_L + d_R P_R) \psi_2] \quad c_L, c_R, d_L, d_R \in \mathbb{R}$$

$$\Rightarrow |\Psi\rangle = M_{LL} | - + - \rangle + M_{LR} | - - + \rangle + M_{RL} | + + - \rangle + M_{RR} | + - + \rangle$$

$$\begin{aligned} \propto & c_L d_L s_{\frac{\theta_3}{2}} \left[c_{\frac{\theta}{2}} c_{\frac{\theta_2}{2}} + e^{i\phi} s_{\frac{\theta}{2}} s_{\frac{\theta_2}{2}} \right] | - + - \rangle + c_L d_R s_{\frac{\theta_2}{2}} \left[c_{\frac{\theta}{2}} c_{\frac{\theta_3}{2}} + e^{i\phi} s_{\frac{\theta}{2}} s_{\frac{\theta_3}{2}} \right] | - - + \rangle \\ & + c_R d_L s_{\frac{\theta_2}{2}} \left[c_{\frac{\theta}{2}} s_{\frac{\theta_3}{2}} - e^{i\phi} s_{\frac{\theta}{2}} c_{\frac{\theta_3}{2}} \right] | + + - \rangle + c_R d_R s_{\frac{\theta_3}{2}} \left[c_{\frac{\theta}{2}} s_{\frac{\theta_2}{2}} - e^{i\phi} s_{\frac{\theta}{2}} c_{\frac{\theta_2}{2}} \right] | + - + \rangle \end{aligned}$$

❖ Individual 2-party entanglement:

$$\mathcal{C}_{12} = \mathcal{C}_{13} = 0, \quad \mathcal{C}_{23} = 2|M_{LL}M_{LR}^* + M_{RL}M_{RR}^*| \quad \leftarrow \text{vanish if } d_L d_R = 0$$

❖ one-to-other entanglement:

$$\mathcal{C}_{2(13)} = \mathcal{C}_{3(12)} = 2\sqrt{(|M_{LL}|^2 + |M_{RL}|^2)(|M_{LR}|^2 + |M_{RR}|^2)} \quad \leftarrow \text{vanish if } c_L c_R = d_L d_R = 0$$

$$\mathcal{C}_{1(23)} = 2|M_{RR}M_{LL} - M_{LR}M_{RL}| \quad \leftarrow \text{vanish if } c_L c_R d_L d_R = 0$$

❖ 3-tangle

$$\tau = \mathcal{C}_{1(23)}^2 - [\mathcal{C}_{12}^2 + \mathcal{C}_{13}^2] = \mathcal{C}_{1(23)}^2$$




➡ All entanglements **vanish** for **weak decays**










$$c_R = d_R = 0$$

F_3 for Vector

$$\mathbf{n} = \mathbf{e}_y$$

$$\mathbf{n} = \frac{1}{\sqrt{3}}(\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)$$





[KS, M.Spannowsky
2310.01477]

Tensor

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$c \equiv c_M + ic_E = e^{i\omega_1}$$
$$d \equiv d_M + id_E = e^{i\omega_2}$$

Tensor


$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$c \equiv c_M + ic_E = e^{i\omega_1}$$

$$d \equiv d_M + id_E = e^{i\omega_2}$$

$$\Rightarrow |\Psi\rangle = M_R|+++\rangle + M_L|---\rangle$$

$$\propto c^*d^*[2e^{i\phi}s\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2} + c\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2}]|+++\rangle + cd[-e^{i\phi}s\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2} + 2c\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2}]|---\rangle$$



Tensor

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$c \equiv c_M + ic_E = e^{i\omega_1}$$


$$d \equiv d_M + id_E = e^{i\omega_2}$$

$$\Rightarrow |\Psi\rangle = M_R|+++\rangle + M_L|---\rangle$$

$$\propto c^*d^*[2e^{i\phi}s\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2} + c\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2}]|+++\rangle + cd[-e^{i\phi}s\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2} + 2c\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2}]|---\rangle$$

♣ $|\Psi\rangle$ interpolates **product states** and the **maximally entangled** state:

$$(M_RM_L = 0) \quad |\pm\pm\pm\rangle \longleftrightarrow |\text{GHZ}\rangle = \frac{1}{\sqrt{2}}(|+++\rangle + |--\rangle) \quad (M_R = M_L)$$



Tensor

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$c \equiv c_M + ic_E = e^{i\omega_1}$$

$$d \equiv d_M + id_E = e^{i\omega_2}$$

$$\Rightarrow |\Psi\rangle = M_R|+++\rangle + M_L|---\rangle$$

$$\propto c^*d^*[2e^{i\phi}s_{\frac{\theta}{2}}s_{\frac{\theta_2}{2}}s_{\frac{\theta_3}{2}} + c_{\frac{\theta}{2}}s_{\frac{\theta_3-\theta_2}{2}}]|+++\rangle + cd[-e^{i\phi}s_{\frac{\theta}{2}}s_{\frac{\theta_3-\theta_2}{2}} + 2c_{\frac{\theta}{2}}s_{\frac{\theta_2}{2}}s_{\frac{\theta_3}{2}}]|---\rangle$$

♣ $|\Psi\rangle$ interpolates **product states** and the **maximally entangled** state:

$$(M_RM_L = 0) \quad |\pm\pm\pm\rangle \longleftrightarrow |\text{GHZ}\rangle = \frac{1}{\sqrt{2}}(|+++\rangle + |--\rangle) \quad (M_R = M_L)$$

♣ **No** individual 2-party entanglements:

$$\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{23} = 0$$

Tensor

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$c \equiv c_M + ic_E = e^{i\omega_1}$$

$$d \equiv d_M + id_E = e^{i\omega_2}$$

$$\Rightarrow |\Psi\rangle = M_R|+++\rangle + M_L|---\rangle$$

$$\propto c^*d^*[2e^{i\phi}s\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2} + c\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2}]|+++\rangle + cd[-e^{i\phi}s\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2} + 2c\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2}]|---\rangle$$

♣ $|\Psi\rangle$ interpolates **product states** and the **maximally entangled** state:

$$(M_RM_L = 0) \quad |\pm\pm\pm\rangle \longleftrightarrow |\text{GHZ}\rangle = \frac{1}{\sqrt{2}}(|+++\rangle + |--\rangle) \quad (M_R = M_L)$$

♣ **No** individual 2-party entanglements:

3-tangle

$$\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{23} = 0 \longrightarrow \tau = \mathcal{C}_{i(jk)}^2 - [\underbrace{\mathcal{C}_{ij}^2}_0 + \underbrace{\mathcal{C}_{ik}^2}_0] = \mathcal{C}_{i(jk)}^2$$

Tensor

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$c \equiv c_M + ic_E = e^{i\omega_1}$$

$$d \equiv d_M + id_E = e^{i\omega_2}$$

$$\Rightarrow |\Psi\rangle = M_R|+++\rangle + M_L|---\rangle$$

$$\propto c^*d^*[2e^{i\phi}s\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2} + c\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2}]|+++\rangle + cd[-e^{i\phi}s\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2} + 2c\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2}]|---\rangle$$

♣ $|\Psi\rangle$ interpolates **product states** and the **maximally entangled** state:

$$(M_RM_L = 0) \quad |\pm\pm\pm\rangle \longleftrightarrow |\text{GHZ}\rangle = \frac{1}{\sqrt{2}}(|+++\rangle + |--\rangle) \quad (M_R = M_L)$$

♣ **No** individual 2-party entanglements:

3-tangle

$$\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{23} = 0 \longrightarrow \tau = \mathcal{C}_{i(jk)}^2 - [\mathcal{C}_{ij}^2 + \mathcal{C}_{ik}^2] = \mathcal{C}_{i(jk)}^2$$

$$\begin{array}{cc} \parallel & \parallel \\ 0 & 0 \end{array}$$

♣ one-to-other entanglements are **universal**:

$$\mathcal{C}_{1(23)} = \mathcal{C}_{2(13)} = \mathcal{C}_{3(12)} = 2|M_RM_L|$$

$$F_3 = 4|M_RM_L|^2$$

Tensor

$$\mathcal{L}_{\text{int}} = [\bar{\psi}_1(c_M + ic_E\gamma_5)\sigma^{\mu\nu}\psi_0][\bar{\psi}_3(d_M + id_E\gamma_5)\sigma_{\mu\nu}\psi_2]$$

$$\begin{aligned} c &\equiv c_M + ic_E = e^{i\omega_1} \\ d &\equiv d_M + id_E = e^{i\omega_2} \end{aligned}$$

$$\Rightarrow |\Psi\rangle = M_R|+++\rangle + M_L|---\rangle$$

$$\propto c^*d^*[2e^{i\phi}s\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2} + c\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2}]|+++\rangle + cd[-e^{i\phi}s\frac{\theta}{2}s\frac{\theta_3-\theta_2}{2} + 2c\frac{\theta}{2}s\frac{\theta_2}{2}s\frac{\theta_3}{2}]|---\rangle$$

❖ $|\Psi\rangle$ interpolates **product states** and the **maximally entangled** state:

$$(M_RM_L = 0) \quad |\pm\pm\pm\rangle \longleftrightarrow |\text{GHZ}\rangle = \frac{1}{\sqrt{2}}(|+++\rangle + |--\rangle) \quad (M_R = M_L)$$

❖ **No** individual 2-party entanglements:

3-tangle

$$\mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{23} = 0 \longrightarrow \tau = \mathcal{C}_{i(jk)}^2 - [\mathcal{C}_{ij}^2 + \mathcal{C}_{ik}^2] = \mathcal{C}_{i(jk)}^2$$

$$\begin{array}{cc} \parallel & \parallel \\ 0 & 0 \end{array}$$




❖ one-to-other entanglements are **universal**:

$$\left. \begin{aligned} \mathcal{C}_{1(23)} = \mathcal{C}_{2(13)} = \mathcal{C}_{3(12)} &= 2|M_RM_L| \\ F_3 &= 4|M_RM_L|^2 \end{aligned} \right\} \begin{array}{l} \text{independent of the coupling} \\ \text{structure (CP phases)} \\ \omega_1, \omega_2 \end{array}$$









F_3 for Tensor

$$\mathbf{n} = \mathbf{e}_y$$

$$\mathbf{n} = \frac{1}{\sqrt{3}}(\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)$$




[KS, M.Spannowsky
2310.01477]




outcome :

$a, a' = +1$ or -1





$c, c' = +1$ or -1




outcome :

$b, b' = +1$ or -1



3-partite Bell inequality



- **Completely** Hidden Variable: $\langle ABC \rangle_{\text{CHV}} = \int p(\lambda)[a(\lambda)b(\lambda)c(\lambda)]d\lambda$

- Mermin correlation: $\mathcal{B}_M = ABC' + AB'C + A'BC - A'B'C'$

- **Mermin inequalities:** $\langle \mathcal{B}_M \rangle_{\text{CHV}} \leq 2 \quad \langle \mathcal{B}_M \rangle_{\text{QM}} \leq 4 \quad [\text{Mermin '90}]$

- **Partially** Hidden Variable: $\langle ABC \rangle_{\text{PHV}} = \int p(\lambda)[ab(\lambda)c(\lambda)]d\lambda, \int p(\lambda)[a(\lambda)bc(\lambda)]d\lambda, \dots$










$$\mathcal{B}_S = ABC + ABC' + AB'C + A'BC$$

- Svetlichny correlation:










$$-A'B'C' - A'B'C - A'BC' - AB'C'$$

- **Svetlichny inequalities:** $\langle \mathcal{B}_S \rangle_{\text{PHV}} \leq 4 \quad \langle \mathcal{B}_S \rangle_{\text{QM}} \leq 4\sqrt{2} \quad [\text{Svetlichny '87}]$

Vector



Tensor



Summary

- Quantum Theoretic quantities (e.g. entanglement and Bell non-locality) are useful in testing the SM and QM at high-energies and provides a new perspective to QFT.
 - Entanglement in $pp \rightarrow t\bar{t}$ at the LHC has recently been measured. Entanglement and Bell violation may be measured also in the other channels, e.g. $pp \rightarrow WW, ZZ, H/Z \rightarrow \tau^+\tau^-$.
 - We demonstrated that the entanglement and Bell violation in $H \rightarrow \tau^+\tau^-$ can be measured in good precision at future e^+e^- colliders.
-

- Entanglement and non-locality in the 3-qubit system from a 3-body decay are studied.

Future works and directions

- Effect of **masses** in the final particles
- More **spin structures**: $SFFV, VVFF, SFVF_{3/2}, SVVT \dots$
- How entanglement is created and evolves in the particle physics interactions/measurements?
- Can we constrain new physics from the general properties of entanglement?

Thank you for listening!