


# **SUSY searches at the LHC**

**Marie-Hélène Genest**


**RPP 2013, January 17<sup>th</sup> 2013**









# Data delivered



Peak instantaneous luminosity :  
 $7.33 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$



(generated 2013-01-12 08:22 including fill 3453)



(generated 2013-01-12 08:22 including fill 3453)



# SUSY searches @ the LHC

Broadly and deeply cover the SUSY signature space

*General strategy to search for SUSY, based on phenomenology oriented searches :*

1. Strong production in a R-parity conserving (RPC) scenario
2. Natural spectrum in a RPC scenario
3. Low effective couplings leading to long-lived SUSY particles
4. Prompt R-parity violating (RPV) scenarios
5. MSSM extensions
6. Other searches



# SUSY direct searches so far

## ATLAS :

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults>

- **Full 2011 data (4.8 /fb, 7 TeV) :**
  - 24 papers
  - 6 conference notes
- **2012 Data Analyses ( $\leq 13$  /fb, 8 TeV) :**
  - 14 conference notes

## CMS :

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>

- **Full 2011 data (4.7-4.9 /fb, 7 TeV) :**
  - 16 papers
  - 7 conference notes
- **2012 Data Analyses ( $\leq 12$  /fb, 8 TeV) :**
  - 1 paper
  - 7 conference notes



# SUSY searches : strategy


Broadly and deeply cover the SUSY signature space

1. Strong production in a R-parity conserving (RPC) scenario




# Strong production in RPC

Inclusive **jets** +  $E_{\text{miss}}$  + X ( $\gamma$ ,  $\ell$ , more jets... depending on NLSP)



$$R = (-1)^{(L+3B+2J)} \quad \text{where} \quad \begin{cases} L = \text{leptonic number} \\ B = \text{baryonic number} \\ J = \text{spin} \end{cases} \quad \begin{array}{l} R = -1 \text{ for sparticles} \\ R = +1 \text{ for SM particles} \end{array}$$

- Lightest sparticle (LSP) stable (WIMP candidate)
- Pair produced sparticles
- Cascade decay down to the LSP



## Search for large and small $\Delta M$

$$\begin{aligned} \text{MET} &\sim \Delta M \\ H_T &= \sum p_T(\text{jet}) [+ p_T(l, \gamma)] \\ M_{\text{Eff}} &= \text{MET} + H_T = 2 M_{\text{SUSY}} \end{aligned}$$




For a given mass, the production of strongly interacting sparticles would dominate





# Inclusive gluino and squarks @ 8 TeV

jets +  $E_T^{\text{miss}}$  signature (veto  $e, \mu$ )



- 12 signal regions to probe different production mechanisms and SUSY mass scales
- Main background: leptonic  $W$ +jets/ $t\bar{t}$ ,  $Z(\nu\nu)$ +jets, Multijets


The 3-jet medium SR as example :






# A more compressed scenario

Models with compressed MSUGRA scenarios  $\Delta M/M_{\text{SUSY}}$  from 0.85 to 0.15







→ The signal regions with the softer cuts allow to go to lower  $\Delta M/M_{\text{SUSY}}$





# Using $\alpha_T$

$$\alpha_T = \frac{E_T^{j2}}{\sqrt{2E_T^{j1}E_T^{j2}(1 - \cos \Delta\phi)}} = \frac{\sqrt{E_T^{j2}/E_T^{j1}}}{\sqrt{2(1 - \cos \Delta\phi)}}$$




Probed squark up to  $\sim 800$  GeV



# Using the razor

- Used in the search for the pair production of two heavy particles, each decaying to an unseen particle plus a visible one
- Idea: move from the lab frame to the CM frame by looking for the boost that makes two jets to be of equal momentum and use this momentum to estimate the mass scale



$$M_R = \sqrt{(|\vec{p}_{j1}| + |\vec{p}_{j2}|)^2 - (p_z^{j1} + p_z^{j2})^2}$$



Peaks at

$$M_\Delta = \frac{M_S^2 - M_{\text{LSP}}^2}{M_S}$$

$$M_T^R = \sqrt{\frac{E_T^{\text{miss}}(p_T^{j1} + p_T^{j2}) - \vec{E}_T^{\text{miss}} \cdot (\vec{p}_T^{j1} + \vec{p}_T^{j2})}{2}}$$

Edge at  $M_\Delta$

$$R = \frac{M_T^R}{M_R}$$





# Using the razor





## 4 exclusive boxes:

1<sup>st</sup>: MU-TAU  $\tau \geq 1 \& \mu \geq 1 \& 0 e$

2<sup>nd</sup>: MU all the other events w/  $\mu \geq 1$

3<sup>rd</sup>: ELE-TAU  $\tau \geq 1 \& e \geq 1 \& 0 \mu$

4<sup>th</sup>: ELE all the other events w/  $e \geq 1$



Limit for tau-enriched  
SUSY events





# Gauge-mediated SUSY breaking (GGM)

Neutralino NLSP (bino or admixture) : photon-based signature


PLB 718 (2012) 411

$2 \gamma + E_T^{\text{miss}}$




$\gamma + \text{b-jets} + E_T^{\text{miss}}$

**Higgsino-like NLSP**




$\gamma + \text{lepton} + E_T^{\text{miss}}$

**Wino-like NLSP**




1211.1167, submitted to PLB

GGM: higgsino-like neutralino,  $\tan\beta = 1.5$ ,  $|\mu| < 0$



ATLAS-CONF-2012-144

GGM: wino-like NLSP





# SUSY searches : strategy

Broadly and deeply cover the SUSY signature space


## 1. Strong production in a R-parity conserving (RPC) scenario

*Inclusive searches have set stringent limits on strongly produced sparticles (1st, 2nd generation squarks, gluinos) [less stringent in case of very compressed scenarios]*

## 2. Natural spectrum in a RPC scenario



# 2- Natural SUSY




L. Hall, LBL workshop 10/2011



Dedicated search program for “3rd generation SUSY”: direct production or gluino-mediated production of sbottom/stop pairs




# Gluino-mediated 3<sup>rd</sup> generation



High jet multiplicities (6-9 jets)  
+ cut on  $E_T^{\text{miss}}/\sqrt{H_T}$




3 b-jets +  $E_T^{\text{miss}}$





# Gluino-mediated 3<sup>rd</sup> generation







# Direct sbottom @ 8 TeV

2 b-jets +  $E_T^{\text{miss}}$



The search exploits the 2-body kinematics



Different signal regions for different  $\Delta m$   
 $\Delta m = m(\tilde{b}_1) - m(\tilde{\chi}_1^0)$



3 leptons + jets +  $E_T^{\text{miss}}$







# Direct stop searches

Several decay modes are possible, depending on the couplings and the SUSY particle mass hierarchy


High cross sections, very similar to SM background

$$\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm \rightarrow b W^\pm \tilde{\chi}_0$$

$$\tilde{t}_1 \rightarrow c \tilde{\chi}_0$$



Low cross section (2 pb or less), high mass:  
Mostly stop  $\rightarrow$  top + LSP



$b + \tilde{\chi}^\pm$  and, where kinematically allowed,  $t + \tilde{\chi}^0$   
Need powerful discriminating variables to reject top BG


Mass ranges ,  $\Delta M$  (stop – neutralino),  $\Delta M$  (stop-chargino),  $\Delta M$ (chargino-neutralino)  
all play a crucial role in the search optimization



# Direct stop searches


- [1] [arxiv:1208.1447](#) (0-lepton 7 TeV)
- [2] [arxiv:1208.2590](#) (1-lepton 7 TeV)
- [3] [arxiv:1209.4186](#) (2-lepton 7 TeV)
- [4] [ATLAS-CONF-2012-166](#) (1-lepton 8 TeV)
- [5] [arxiv:1208.4305](#) (very light stop: 2-lepton 7 TeV)
- [6] [arxiv:1209.2102](#) (light stop: 1/2-lepton, bjets 7 TeV)
- [7] [ATLAS-CONF-2012-167](#) (2-lepton 8 TeV)
- [8] [ATLAS-CONF-2013-001](#) (0-lepton, bb+MET 8 TeV)

Status: December 2012





# 2- Natural SUSY




L. Hall, LBL workshop 10/2011

Dedicated search program for  
“Electroweak SUSY”: direct  
production neutralinos,  
charginos, sleptons





# Electroweak production : chargino/neutralino

3 leptons +  $E_T^{\text{miss}}$



Three-lepton events with an  $ee$  or  $\mu\mu$  OSSF dilepton pair, where the third lepton is either an electron or a muon



Chargino-neutralino limits  
extended up to  $\sim 650 \text{ GeV}$

$M_T \text{ (GeV)}$	$E_T^{\text{miss}} \text{ (GeV)}$	$M_{\ell\ell} < 75 \text{ GeV}$		$75 \text{ GeV} < M_{\ell\ell} < 105 \text{ GeV}$	
		total bkg	observed	total bkg	observed
$> 160$	50 – 100	$2.1 \pm 0.5$	4	$3.3 \pm 0.5$	3
	100 – 150	$1.7 \pm 0.4$	0	$1.8 \pm 0.2$	1
	150 – 200	$0.8 \pm 0.3$	1	$0.63 \pm 0.16$	1
	$> 200$	$0.25 \pm 0.20$	0	$0.58 \pm 0.19$	1





# Electroweak production : sleptons


2 leptons +  $E_T^{\text{miss}}$

- Reduce the WW background by using its endpoint in transverse mass,  $m_{T2}$  (at  $\sim 90$  GeV)

$$m_{T2} = \min_{\vec{q}_T^{(1)} + \vec{q}_T^{(2)} = \vec{p}_T} (\max(m_T(\vec{p}_T^{(1)}, \vec{q}_T^{(1)}), m_T(\vec{p}_T^{(2)}, \vec{q}_T^{(2)})))$$



## First limits on sleptons since LEP





# SUSY searches : strategy

Broadly and deeply cover the SUSY signature space

1. Strong production in a R-parity conserving (RPC) scenario
2. Natural spectrum in a RPC scenario


Comprehensive program for the third generation sector in place with limits starting to bite into naturalness – need to continue to cover the full phase space. EW searches also underway with first limits on direct slepton since LEP.

3. Low effective couplings leading to long-lived SUSY particles




# R-parity violation and long-lived sparticles

- R-parity violation (RPV):  $W = W_{MSSM} + \underbrace{\lambda_{ijk} L_i L_j \bar{E}_k + \lambda'_{ijk} L_i Q_j \bar{D}_k}_{\text{Lepton Number Violation (LFV)}} + \underbrace{\kappa_i L_i H_u + \lambda''_{ijk} \bar{U}_i \bar{D}_j \bar{D}_k}_{\text{Baryon Number Violation (BNV)}}$
- RPV can lead to a displaced vertex if  $\lambda, \lambda', \lambda''$  is very small
- A long-lived (LL) particle can also occur in RPC :
  - $\Delta M(\tilde{\chi}_1^\pm, \tilde{\chi}_1^0) \sim 100 \text{ MeV}$  (eg. in AMSB) : disappearing track
  - LL gluino due to the very heavy squarks mediating its decay : R-hadron
  - Weak coupling NLSP-gravitino in GMSB : LL slepton






*Challenging analyses requiring dedicated developments (re-tracking, trigger, ...)*





# Displaced vertex

- RPV with  $\lambda'_{2ij} \neq 0$  : sparticle decay gives a multi-track vertex with a high- $p_T$  muon, a few mm to  $\sim 10$  cm from the IP
- Dedicated tracking to increase signal efficiency
- Remove vertices reco'd in regions of high-density material
- Background-free analysis in  $M_{\text{vertex}} / N_{\text{track}}$  plane



Sample	$m_{\tilde{q}}$ [GeV]	$m_{\tilde{\chi}_1^0}$ [GeV]
MH	700	494
ML	700	108
HH	1500	494




# LL chargino : disappearing track



- In jet (from ISR) +  $E_T^{\text{miss}}$  events, search for high- $p_T$  isolated tracks that stop in outer TRT

$$pp \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_1^0 j, \quad pp \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_1^\mp j$$

$$\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm \quad \text{branching ratio set to 100\%}$$



For  $\Delta m = 160$  (170) MeV, the chargino mass limit is set at 103 (85) GeV






# R-hadron / long-lived slepton

- Selection based on good quality, isolated high- $p_T$  track
- Use the time of flight and  $dE/dx$  measurement to get  $\beta$ ,  $\beta\gamma$

Three analyses :

- Full-detector
- MS-agnostic (ignore MS)
- ID-only

} cover the lack of knowledge of R-hadron interactions with the detector and the lifetimes for which they would not reach the calorimeters



Exclude directly produced LL sleptons up to 278 GeV and R-hadrons containing a gluino up to 985 GeV (generic interaction model)



# SUSY searches : strategy

Broadly and deeply cover the SUSY signature space

*The ATLAS SUSY group has a well-defined strategy to search for SUSY, based on phenomenology oriented searches :*

1. Strong production in a R-parity conserving (RPC) scenario
2. Natural spectrum in a RPC scenario
3. Low effective couplings leading to long-lived SUSY particles
4. Prompt RPV scenarios
5. MSSM extensions






# Leptonic RPV

- Search for a tau sneutrino decaying into  $e\mu$ ,  $e\tau$  or  $\mu\tau$  ( $\lambda'_{311}$ ,  $\lambda_{i3k}$ )
- Limit on  $\sigma \times \text{BR}$  as a function of the sneutrino mass for each channel

- 4 leptons +  $E_T^{\text{miss}}$


$$\begin{aligned}
 pp &\rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^-, \\
 \tilde{\chi}_1^\pm &\rightarrow W^\pm \tilde{\chi}_1^0, \\
 \tilde{\chi}_1^0 &\rightarrow \nu_{i/j} \ell_{i/j}^\pm \ell_k^\mp.
 \end{aligned}$$






# Hadronic RPV & scalar gluon

- RPV gluino decay into three quarks
- Resolved analysis with 6 jets
- Boosted analyses for low-mass gluinos



Resolved analysis : exclude up to 666 GeV  
Boosted analysis : exclude up to 255 GeV

- Massive coloured scalar (sgluon) with  $R=1$  (beyond MSSM)
- Pair production: 2 resonances  $M1, M2$  reconstructed with  $\geq 4$  high- $p_T$  jets



Exclude scalar gluons for masses from 150 to 287 GeV



# 1. Inclusive search


# 2. Natural spectrum

# 3. Long-lived sparticles

# 4. Prompt RPV

# 5. MSSM extension

ATLAS SUSY Searches\* - 95% CL Lower Limits (Status: Dec 2012)



\*Only a selection of the available mass limits on new states or phenomena shown.  
All limits quoted are observed minus 1 $\sigma$  theoretical signal cross section uncertainty.



# SUSY searches : strategy

Broadly and deeply cover the SUSY signature space

*The ATLAS SUSY group has a well-defined strategy to search for SUSY, based on phenomenology oriented searches :*


1. Strong production in a R-parity conserving (RPC) scenario
2. Natural spectrum in a RPC scenario
3. Low effective couplings leading to long-lived SUSY particles
4. Prompt RPV scenarios
5. MSSM extensions
6. Other searches



# General search

- Did we miss anything? Clean up with a general search for new physics
- All event topologies involving electrons, photons, muons, jets, b-tagged jets and missing transverse momentum in a single analysis (655 channels defined)
- Scan the effective mass distribution of each final state for deviations from the Standard Model prediction (note : BG from MC only)

Distribution of the p-values :



- Consistent with the expectation from toy experiments
- No event class found with a p-value smaller than  $10^{-3}$
- No big signal hidden in the previously unexplored channels




# Indirect search : $B_s \rightarrow \mu\mu$


SM prediction:

$$\text{SM } B(B_s \rightarrow \mu\mu) = (3.2 \pm 0.2) \times 10^{-9}$$


Buras et al. arXiv:1012.1447




Branching ratio very sensitive to new physics



$$BR(B_s \rightarrow \mu^+ \mu^-) \propto \tan^6 \beta / m_A^4$$





Fit performed in 8 (for 2011) + 7 (for 2012) BDT bins





# Indirect search $B_s \rightarrow \mu\mu$



Combining 2011+2012 data

Bkg only hypothesis p-value is  $5 \times 10^{-4}$  corresponding to  $3.5 \sigma$

$$\mathcal{B}(B_s \rightarrow \mu^+\mu^-) = 3.2_{-1.2}^{+1.4}(\text{stat})_{-0.3}^{+0.5}(\text{syst}) \times 10^{-9}$$

First evidence of the decay  $B_s \rightarrow \mu^+\mu^-$

Consistent with the SM!

Submitted to PRL [arXiv:1211.2674](https://arxiv.org/abs/1211.2674)



# Conclusion

- Strong and diverse program for SUSY searches
- 2012 data analyses well under way, complete dataset being analysed now
- Goals :
  - Extend inclusive searches, also for compressed spectra
  - Continue the stop search, covering all signatures
  - Expand gaugino/slepton searches
  - Continue developing innovative searches for RPV & long-lived signatures



# Additional material



# What do the various lines mean ?

## ❑ Exclusion limits : a new standard ATLAS/CMS procedure (>June 2012)

- Ease the life of theorist by separating the signal theoretical and experimental systematics

**Expected limit:**



- **Central value:** all uncertainties included in the fit as nuisance parameters, except theoretical signal uncertainties (PDF, scales)


- **$\pm 1\sigma$  band** :  $\pm 1\sigma$  results of the fit

**Observed limit:**



- **Central value:** Idem as for expected limit
- **$\pm 1\sigma$  band** : re-run and increase/decrease the signal cross section by the theoretical signal uncertainties (PDF, scales)

**Excluded Model Cross section (SMS)** ←



- ➔ Number quoted in paper correspond to observed -1  $\sigma$  observed (conservative)




# ATLAS 0-lepton @ 8 TeV

	Requirement	Channel				
		A 2-jets	B 3-jets	C 4-jets	D 5-jets	E 6-jets
Trigger	$E_T^{\text{miss}} [\text{GeV}] >$	160				
	$p_T(j_1) [\text{GeV}] >$	130				
Pile-up	$p_T(j_2) [\text{GeV}] >$	60				
	$p_T(j_3) [\text{GeV}] >$	–	60	60	60	60
	$p_T(j_4) [\text{GeV}] >$	–	–	60	60	60
	$p_T(j_5) [\text{GeV}] >$	–	–	–	60	60
	$p_T(j_6) [\text{GeV}] >$	–	–	–	–	60
QCD	$\Delta\phi(\text{jet}, \mathbf{E}_T^{\text{miss}})_{\text{min}} [\text{rad}] >$	0.4 ( $i = \{1, 2, (3)\}$ )		0.4 ( $i = \{1, 2, 3\}$ ), 0.2 ( $p_T > 40 \text{ GeV jets}$ )		
	$E_T^{\text{miss}} / m_{\text{eff}}(Nj) >$	0.3/0.4/0.4 (2j)	0.25/0.3/– (3j)	0.25/0.3/0.3 (4j)	0.15 (5j)	0.15/0.25/0.3 (6j)
SUSY mass scale	$m_{\text{eff}}(\text{incl.}) [\text{GeV}] >$	1900/1300/1000	1900/1300/–	1900/1300/1000	1700/–/–	1400/1300/1000

12 Signal regions split in tight / medium / loose requirements




# 0-lepton @ 8 TeV





Z+jets BG estimation

$t\bar{t}$  BG estimation




3-jet channel example



W+jets BG estimation


QCD BG estimation





# Direct stop searches @ 7 TeV



- Five searches with 5 fb<sup>-1</sup> (+ 1 in natural GMSB with 2 fb<sup>-1</sup>) @ 7 TeV





# General search


$$p = A \int_0^\infty db G(b; N_{SM}, \delta N_{SM}) \sum_{i=N_{obs}}^\infty \frac{e^{-b} b^i}{i!}$$



With SUSY, many event classes with small p-value would be expected.





# SUSY Higgs sector



- $\phi \rightarrow \tau\tau$  Signal searched for in three channels:  $e\mu$  ;  $e\tau_{\text{had}}$  ;  $\mu\tau_{\text{had}}$

Selection :

- Veto events with additional isolated leptons
- MET pointing in the direction of the visible products
- At most 1 jet with  $p_T > 30$  GeV
- Two categories (jet  $p_T > 20$  GeV) :
  - no b-tag jet
  - at least one b-tagged jet



- Stop mixing parameter  $X_t = 2M_{\text{SUSY}}$
- Higgsino mass  $\mu = 200$  GeV
- Gluino mass  $M_g = 800$  GeV
- Stop and sbottom trilinear couplings  $A_b = A_t$